

**Endothelial NADPH oxidase-2 promotes interstitial cardiac fibrosis and diastolic dysfunction through pro-inflammatory effects and endothelial-mesenchymal transition.**

Colin E Murdoch PhD, Sanjay Chaubey MBBCh, Lingfang Zeng PhD, Bin Yu PhD, Aleksander Ivetic PhD, Simon J Walker BSc, \*Davy Vanhoutte PhD, \*Stephane Heymans MD, PhD, David J Grieve PhD, Alison C Cave PhD, Alison C Brewer PhD, Min Zhang MD, PhD, Ajay M Shah MD, FMedSci.

King's College London British Heart Foundation Centre of Excellence, Cardiovascular Division, London, UK; and \*Cardiovascular Research Institute Maastricht (CARIM), Maastricht University, The Netherlands.

**Brief title:** Endothelial NOX2 enhances cardiac fibrosis

**Correspondence:** Professor Ajay M Shah, Cardiovascular Division, The James Black Centre, King's College London, 125 Coldharbour Lane, London SE5 9NU, UK. Tel: +44 2078485189; Fax: +44 2078485193; E-mail: [ajay.shah@kcl.ac.uk](mailto:ajay.shah@kcl.ac.uk)

**Disclosures:** none

**Funding:** British Heart Foundation (RG/08/011/25922, RE/08/003); Fondation Leducq Transatlantic Network of Excellence Award; the Department of Health via a National Institute for Health Research (NIHR) Biomedical Research Centre award to Guy's & St Thomas' NHS Foundation Trust in partnership with King's College London and King's College Hospital NHS Foundation Trust

## **ABSTRACT**

**Objectives.** To investigate the effect of endothelial dysfunction on development of cardiac hypertrophy and fibrosis.

**Background.** Endothelial dysfunction accompanies cardiac hypertrophy and fibrosis but its contribution to these conditions is unclear. Increased NADPH oxidase-2 (NOX2) activation causes endothelial dysfunction.

**Methods.** Transgenic mice with endothelial-specific NOX2 overexpression (TG) and wild-type littermates received chronic AngII infusion (1.1mg/kg/day, 2 weeks) to induce hypertrophy and fibrosis.

**Results.** TG had similar systolic hypertension and hypertrophy to wild-type but developed greater cardiac fibrosis and evidence of isolated left ventricular diastolic dysfunction ( $P<0.05$ ). TG myocardium had more inflammatory cells and VCAM-1-positive vessels than wild-type after AngII (both  $P<0.05$ ). TG microvascular endothelial cells (EC) treated with AngII recruited 2-fold more leukocytes than wild-type in an *in vitro* adhesion assay ( $P<0.05$ ). However, inflammatory cell NOX2 *per se* was not essential for the pro-fibrotic effects of AngII. TG mice showed a higher level of endothelial-mesenchymal transition (EMT) than wild-type after AngII infusion. In cultured EC treated with AngII, NOX2 enhanced EMT as assessed by the relative expression of fibroblast versus endothelial-specific markers.

**Conclusion.** AngII-induced endothelial NOX2 activation has profound pro-fibrotic effects in the heart *in vivo*, leading to a diastolic dysfunction phenotype. Endothelial NOX2 enhances EMT and has pro-inflammatory effects. This may be an important mechanism underlying cardiac fibrosis and diastolic dysfunction during increased renin-angiotensin activation.

**Key Words:** angiotensin II, diastolic dysfunction, endothelium, endothelial-mesenchymal transition, NADPH oxidase

## **Abbreviations**

AngII, angiotensin II;  $\alpha$ SMA,  $\alpha$ -smooth muscle actin; BM, bone marrow; CMEC, coronary microvascular endothelial cells; EC, endothelial cell; EMT, endothelial-mesenchymal transition; NOX, NADPH oxidase; ROS, reactive oxygen species; TG, transgenic; VCAM-1, vascular cell adhesion molecule-1

## **INTRODUCTION**

Endothelial dysfunction accompanies conditions such as hypertension that predispose to adverse cardiac remodeling, and predicts future cardiovascular morbidity.(1) As well as modulating vascular function, the endothelium has direct effects on cardiac function.(2-4) Pioneering work by Brutsaert and colleagues(3) demonstrated that cardiac endothelial cells directly influence myocardial contraction, a concept later confirmed in humans *in vivo*.(5) The importance of endothelial-cardiomyocyte crosstalk has been demonstrated in several disease settings.(3,6) However, the role of endothelial dysfunction in the pathogenesis of adverse cardiac remodeling caused by increased renin-angiotensin activation remains unclear.

A major driver of endothelial dysfunction is the reactive oxygen species (ROS)-generating enzyme, NADPH oxidase-2 (NOX2), which is activated by angiotensin II (AngII) and other agonists.(7) NOX2 is expressed in endothelium as well as cardiomyocytes, adventitial fibroblasts, inflammatory cells and conduit vascular smooth muscle cells. Vascular NOX2 expression and activity are increased in hypertension where it modulates intracellular signaling pathways that promote vascular remodeling, contributes to vasodilator dysfunction through superoxide-mediated inactivation of nitric oxide, and may have pro-inflammatory effects.(7)

To investigate the pathophysiologic roles of endothelial NOX2, we generated a transgenic mouse model with endothelial-specific NOX2 overexpression.(8) This model has a 2-fold increase in endothelial NOX2 protein levels and displays augmented NOX2 activation in response to AngII. Here, we used this model to investigate the effects of endothelial NOX2 activation on the development of cardiac hypertrophy, fibrosis and dysfunction during chronic AngII elevation.

## **METHODS**

*Detailed methods are provided in the online supplement.*

Animal experiments complied with UK Home Office regulations. Male transgenic mice with endothelium-targeted NOX2 overexpression (TG) (8) were compared with wild-type littermates. Chimeric mice were generated by bone marrow (BM) transplantation.(9) AngII (1.1mg/kg/day) or vehicle were infused via osmotic minipumps. Blood pressure was measured by radiotelemetry or tail-cuff plethysmography.(10) Left ventricular pressure-volume relations and echocardiography were performed as described.(11) Cardiac fibrosis was quantified in Picrosirius Red-stained paraffin sections while immunostaining was performed in cryosections. LV homogenate NADPH oxidase activity was measured using lucigenin-enhanced chemiluminescence.(12)

*In vitro* leukocyte adhesion to coronary microvascular endothelial cells (CMEC) from TG or wild-type mouse hearts was studied in a flow chamber,(13) with or without AngII (0.1µmol/L, 4h). To assess endothelial-mesenchymal transition (EMT), cultured human aortic EC were infected with Ad.βGal or Ad.NOX2 virus, then treated with 0.1µmol/L AngII or vehicle for up to 5 days. Co-culture experiments with EC and fibroblasts were performed in Transwell dishes.

Data are expressed as mean±SEM. Comparisons were made by repeated measures ANOVA, 2-way ANOVA with Bonferroni post-hoc testing, 1-way ANOVA or Student's t-test as appropriate. P<0.05 was considered significant.

## **RESULTS**

### ***Effects of endothelial NOX2 overexpression on AngII-induced hypertension and cardiac hypertrophy.***

Chronic AngII infusion for 14 days caused similar systolic hypertension in TG and wild-type mice (Supplementary Fig. 1A). Left ventricular NADPH oxidase activity increased in both AngII-treated groups, with a significantly greater rise in TG (Supplementary Fig. 1B, supplementary Fig. 2). AngII infusion induced a similar increase in left ventricle/body weight

ratio or cardiomyocyte cross-sectional area in TG and wild-type mice (Supplementary Fig. 1C,D).

***Endothelial NOX2 overexpression potentiates AngII-induced cardiac fibrosis.***

The extent of AngII-induced fibrosis was significantly (~2-fold) greater in TG than wild-type hearts (Supplementary Fig. 1E). TG also had significantly higher pro-collagen I mRNA expression than wild-type (Supplementary Fig. 1F). Representative examples of the fibrosis and myocyte hypertrophy are shown in Fig. 1.

***LV diastolic dysfunction with preserved systolic function in endothelial NOX2 TG.***

TG had significantly lower echocardiographic left ventricular end-diastolic dimensions than wild-type after chronic AngII infusion (Supplementary Fig. 3). However, there were no significant differences in indices such as interventricular septal thickness or ejection fraction. More detailed assessment was undertaken by pressure-volume analysis. Systolic function assessed by end-systolic elastance or LV  $dP/dt_{max}/EDV$  was similar in AngII-treated wild-type and TG groups (Supplementary Fig. 4A,B). Left ventricular end-diastolic volume increased in wild-type mice after AngII but decreased in TG (Supplementary Fig. 4C). Left ventricular end-systolic volume tended to increase in wild-type after AngII treatment but fell significantly in TG (Supplementary Fig. 4D). As a result, both stroke volume and stroke work were significantly lower in AngII-treated TG versus wild-type (Supplementary Fig. 4E, Table 1). Isovolumic left ventricular relaxation time-constant and left ventricular end-diastolic pressure were similar in wild-type and TG groups (Table 1). However, left ventricular diastolic stiffness assessed by the end-diastolic pressure volume relationship was significantly higher in AngII-treated TG versus wild-type hearts (Supplementary Fig. 4F). Although there was no difference in systolic BP, arterial elastance was significantly higher in AngII-treated TG (Table 1). Supplementary Fig. 4G,H show representative steady-state pressure-volume loops and the response to preload reduction in wild-type and TG. These indicate that TG

hearts were smaller and stiffer than wild-type after chronic AngII, indicating a phenotype of isolated diastolic left ventricular dysfunction with preserved systolic function.

***Endothelial NOX2 enhances myocardial inflammatory cell infiltration.***

To investigate mechanisms underlying the increased fibrosis in TG, we assessed myocardial inflammatory cell infiltration. TG hearts had significantly more CD45<sup>+</sup> inflammatory cells and Mac3<sup>+</sup> cells (macrophages) after AngII infusion than wild-type (Fig. 2, Supplementary Fig. 5). AngII treatment increased connective tissue growth factor and osteopontin mRNA levels but levels were similar in wild-type and TG (Supplementary Fig. 6). There was no difference between groups in TGF $\beta$ , TNF $\alpha$ , MCP-1 or endothelin-1 mRNA levels.

To confirm that the effects of endothelial NOX2 on inflammatory infiltration and fibrosis were ROS-dependent, mice were treated with the antioxidant N-acetylcysteine concurrent with AngII infusion. BP and cardiac hypertrophy were similar in wild-type and TG treated with N-acetylcysteine (Supplementary Fig. 7). N-acetylcysteine markedly reduced CD45<sup>+</sup> cell number in both wild-type and TG AngII-treated groups, and prevented the development of fibrosis (Supplementary Fig. 7).

***NOX2 enhances endothelial activation and leukocyte-endothelial interaction.***

We assessed the interaction between inflammatory cells and NOX2-overexpressing endothelial cells in an *in vitro* adhesion assay under flow conditions. Leukocyte attachment to TG and wild-type CMEC was low at baseline, but after AngII treatment (0.1  $\mu$ M, 4 h) significantly more leukocytes attached to TG CMEC (Fig. 2C). TG CMEC had significantly higher vascular cell adhesion molecule-1 (VCAM-1) protein expression than wild-type after AngII treatment (Fig. 2D). In line with this, LV sections of TG treated with chronic AngII had more VCAM-1 positive blood vessels than wild-type (Fig. 2E, Supplementary Fig. 8). There was no difference in capillary density between TG and wild-type hearts (Fig. 2F).

***Inflammatory cell NOX2 is not essential for AngII-induced cardiac fibrosis.***

Since NOX2 is abundantly expressed in inflammatory cells, we investigated whether NOX2 in these cells was required for the pro-fibrotic response to AngII. We used BM transplantation to create chimeric mice with different NOX2 genotypes in BM and resident tissue cells: (a) wild-type recipient mice with wild-type BM (WT:WT); (b) wild-type recipients with BM from global NOX2 knockout mice (KO:WT); (c) global NOX2 knockout mice with wild-type BM (WT:KO); (d) NOX2 TG with TG BM (TG:TG); (e) NOX2 TG with wild-type BM (WT:TG); and (f) wild-type mice with TG BM (TG:WT). Successful alteration of NOX2 genotype after BM transplantation was confirmed by assessing ROS production in recipient BM and peripheral blood mononuclear cells (Supplementary Fig. 9A).

AngII infusion caused similar increases in BP and hypertrophy among groups (Supplementary Fig. 9B,C). AngII-stimulated cardiac fibrosis was significantly lower in global NOX2 KO versus wild-type, as reported previously(12) (Supplementary Fig. 10). WT:WT and KO:WT mice both had similar fibrosis to un-transplanted wild-type mice (Supplementary Fig. 10A,B). In contrast, WT:KO mice had significantly less fibrosis, similar to un-transplanted knockout mice (Supplementary Fig. 10A,B). Thus, knockout BM cells did not significantly alter AngII-induced fibrosis. Fibrosis in NOX2 TG:TG was similar to that in WT:TG while transplantation of TG BM to wild-type mice did not increase the level of fibrosis to that in the TG:TG group (Supplementary Fig. 10A right-hand panels, Supplementary Fig. 10B). Thus, TG BM cells did not significantly alter AngII-induced fibrosis. Taken together, these results indicate that NOX2 in BM cells is dispensable for the pro-fibrotic effects of AngII and it is NOX2 in the resident tissue cells that is essential. Furthermore, endothelium-targeted NOX2 overexpression does not significantly alter the effects of BM cells as compared to wild-type.

***NOX2 promotes endothelial-mesenchymal transition (EMT).***

Recent studies indicate that EMT is an important contributor to cardiac fibrosis during chronic pressure overload.(14) We therefore investigated whether enhancing NOX2 activity in endothelial cells affected EMT. Hearts of AngII-treated TG had significantly higher levels of fibroblast-specific  $\alpha$ -smooth muscle actin ( $\alpha$  SMA) and collagen-1 than wild-type while levels of endothelial-specific CD31 were significantly lower (Fig. 3A). Myocardial sections of AngII-treated TG showed increased evidence of EMT, as indicated by co-localization of endothelial-specific and mesenchymal markers, compared with wild-type (Fig. 3B).

To further assess the effects of endothelial NOX2, we studied cultured human aortic EC with adenoviral-mediated overexpression of NOX2 or  $\beta$ -galactosidase control, and treated with AngII. AngII caused a modest increase in fibroblast-specific markers (FSP-1,  $\alpha$  SMA) and a concomitant decrease in endothelial-specific markers (CD31, CD144) in control EC. This transition from endothelial- to fibroblast-specific expression was enhanced in NOX2-overexpressing cells (Fig. 4). NOX2 overexpression *per se* caused a small switch from endothelial to fibroblast-specific expression in the absence of AngII. Supplementary Fig. 11A shows representative examples of co-expression of endothelial-specific and mesenchymal markers in NOX2-overexpressing cells after AngII. Cellular morphology was significantly different after 5 days of AngII treatment, with NOX2-overexpressing cells showing a more elongated fibroblast phenotype (Supplementary Fig. 11B).

#### ***Paracrine effects of endothelial cells on fibroblasts?***

It is feasible that NOX2 elevation in EC affects fibroblasts through the direct effects of ROS or other released factors. To assess this possibility, we performed experiments in which human aortic EC overexpressing NOX2 or  $\beta$ -galactosidase were co-cultured with fibroblasts (see Supplementary Methods). Experiments were performed with and without AngII. No difference was found between groups in the fibroblast expression of pro-collagen I mRNA or in markers of myofibroblast transformation (Supplementary Fig. 12).



## **DISCUSSION**

We investigated the specific effects of endothelial NOX2 activation on AngII-induced cardiac remodeling, taking advantage of a recently developed mouse model with endothelium-targeted overexpression of NOX2.(8) The 2-fold increase in endothelial NOX2 protein in this model has no basal effects but enhances AngII-stimulated increases in NOX2 activity. In the present study, we found that *in vivo* AngII-induced endothelial NOX2 activation (a) enhances the development of cardiac fibrosis independent of cardiomyocyte hypertrophy, and leads to a phenotype of left ventricular diastolic dysfunction with preserved systolic function; (b) promotes myocardial inflammatory cell infiltration through increased endothelial expression of VCAM-1 (i.e. endothelial activation), although NOX2 in inflammatory cells is dispensable for the pro-fibrotic effects; (c) enhances EMT which contributes to the cardiac fibrosis. Collectively, these data suggest that NOX2-driven endothelial dysfunction and activation may be an important mechanism that promotes cardiac fibrosis and left ventricular diastolic dysfunction in pathologic settings of renin-angiotensin system activation.

### ***Endothelium-cardiomyocyte crosstalk***

The cardiac endothelium forms a strategic interface between circulating blood and myocardial tissue. Crosstalk between cardiac EC and cardiomyocytes plays important roles in normal cardiac development.(3,4) The cardiac endothelium also influences contractile function in the adult heart, for example through paracrine factors such as nitric oxide and endothelin.(2) Reciprocal signaling between cardiomyocytes and the myocardial microvasculature has been found to be important during chronic pressure overload. For example, signaling from EC to cardiomyocytes through the neuregulin 1-ErbB2 axis promotes cardiomyocyte survival during pressure overload,(15) and angiogenic factors released by cardiomyocytes in the chronically overloaded heart regulate myocardial capillary

density.(11,16) While these studies support the potential for endothelial dysfunction to affect cardiomyocyte growth, survival and contractile function, the effects on cardiac fibrosis in the heart under stress are unclear.

### ***Pro-inflammatory effects of endothelial NOX2***

Here, we demonstrate that endothelial dysfunction has a profound ROS-dependent impact on the development of cardiac fibrosis, independent of cardiomyocyte hypertrophy. One mechanism underlying these effects may be the pro-inflammatory properties of activated endothelium. The healthy endothelium is anti-inflammatory and anti-thrombotic but in pathologic settings such as renin-angiotensin system activation, it becomes activated and promotes increased interaction with circulating inflammatory cells.(17) Previous studies suggested that NOX2 is involved in cytokine- and AngII-induced EC activation and the expression of adhesion molecules such as VCAM-1,(18,19) and in enhanced monocyte binding to ECs under oscillatory shear stress.(20) Here, we found that NOX2-overexpressing CMEC had a larger increase in VCAM-1 expression after AngII stimulation than wild-type, which was associated with greater endothelial-leukocyte adhesion in an *in vitro* flow assay. Consistent with a similar pro-inflammatory action *in vivo*, TG myocardium had more VCAM-1-positive blood vessels and inflammatory cells after AngII treatment than wild-type. These results clearly indicate that AngII-induced endothelial NOX2 activation augments an inflammatory response in the heart *in vivo*. Whether other cell types (e.g. vascular smooth muscle cells, pericytes) are also involved is an interesting question.

In addition to EC, NOX2 is abundant in inflammatory cells where it is suggested to contribute to oxidative stress, for example in atherosclerosis.(21) NOX2 activation in inflammatory cells is involved in AngII-induced vascular hypertrophy.(22) It was therefore possible that NOX2 activation in infiltrating inflammatory cells (as well as the endothelium) might be important in the development of cardiac fibrosis. To dissect the specific contribution

of inflammatory cell NOX2 activation, we generated chimeric mice in which the BM comprised NOX2-competent or -deficient cells. These series of studies demonstrated that NOX2 activation in infiltrating inflammatory cells is dispensable for the development of AngII-induced fibrosis and that it is NOX2 in recipient heart cells that is essential for the profibrotic effects. This result suggests that although NOX2 may be important for certain functions of inflammatory cells (e.g. phagocytosis, LDL oxidation in atherosclerosis [21]), it is not essential for all functions and is dispensable in the setting of AngII-induced cardiac pathology.

### ***Role of EMT***

Another mechanism by which endothelial dysfunction might promote cardiac fibrosis is through EMT. The transition of epithelial and endothelial cells to a mesenchymal phenotype is well recognized to be important in cardiac development. In a landmark study, Zeissberg et al.(14) showed that EMT makes a significant contribution to pathologic cardiac fibrosis during chronic pressure overload. EC production of endothelin-1 has been reported to contribute to EMT in the diabetic heart.(23) We therefore investigated whether endothelial NOX2 activation affects EMT. We found evidence of significantly increased EMT in TG hearts after chronic AngII as compared with wild-type. More definitive evidence that endothelial NOX2 promotes EMT was obtained in cultured EC where NOX2 significantly enhanced the transition from endothelial to fibroblast phenotype after AngII treatment. These results strongly suggest that NOX2-dependent increase in EMT may be a crucial mechanism that enhances cardiac fibrosis in response to chronic AngII treatment.

We also tested the possibility that paracrine effects of EC on fibroblasts, for example through the direct effects of ROS released by EC, may contribute to fibrosis. *In vitro* co-culture experiments with EC and fibroblasts did not provide evidence in support of this

possibility. However, a possible contribution of such a mechanism *in vivo* cannot be excluded.

### ***Isolated diastolic dysfunction***

A striking functional consequence of endothelial NOX2 overexpression on the response to chronically elevated AngII was a phenotype of LV diastolic dysfunction with preserved systolic function, resulting in smaller and stiffer hearts and reduced stroke volumes in TG compared with wild-type. This was documented in TG by the gold-standard method of *in vivo* pressure-volume analysis, which demonstrated significantly increased left ventricular diastolic passive stiffness in the AngII-treated TG versus wild-type group whereas load-independent indices of systolic function were similar in the two groups. We found no significant difference between the groups in parameters of active left ventricular relaxation (e.g. the isovolumic relaxation time-constant), suggesting that cardiomyocyte relaxation was probably similar in the two groups. This contractile dysfunction phenotype is consistent with, and fully explainable by, the increased fibrosis observed in TG independent of differences in cardiomyocyte hypertrophy. The use of an endothelium-targeted transgenic model allowed us to study the role of endothelial dysfunction/activation independent of other changes that occur in the remodeling heart during increased renin-angiotensin system activation. As mentioned earlier, NOX2 is also expressed in cardiomyocytes and previous studies in cultured cells and *in vivo* models have shown that cardiomyocyte NOX2 activation may contribute to the development of AngII-induced cardiac hypertrophy.(12) No difference was found in extent of hypertrophy between wild-type and TG in the current study, consistent with the fact that NOX2 was overexpressed specifically in EC. Therefore, the phenotype of diastolic left ventricular dysfunction with preserved systolic function found in the current model was the consequence specifically of endothelial dysfunction/activation.

The clinical syndrome of heart failure with preserved ejection fraction is currently receiving significant attention due to the high prevalence of the condition, controversy regarding its etiology and pathogenesis, and the lack of effective therapies.(24) Whilst simple extrapolations cannot be made from the current study in a murine model to the clinical setting, the occurrence of a contractile dysfunction phenotype that apparently closely resembles the cardiac dysfunction found in human heart failure with preserved ejection fraction is of great interest. It is notable that previous human studies suggested that inflammation contributes to enhanced fibrosis in such patients,(25) consistent with the findings of increased inflammatory cell infiltration, fibrosis and diastolic dysfunction in our study. The current results raise the intriguing possibility that endothelial dysfunction/activation and downstream pathways such as inflammation and EMT may be important mechanisms contributing to the development of heart failure with preserved ejection fraction. If so, this could provide new therapeutic options to tackle this prevalent condition.

## REFERENCES

1. Heitzer T, Schlinzig T, Krohn K, Meinertz T, Munzel T. Endothelial dysfunction, oxidative stress, and risk of cardiovascular events in patients with coronary artery disease. *Circulation* 2001;104:2673-8.
2. Shah AM. Paracrine modulation of heart cell function by endothelial cells. *Cardiovasc Res* 1996;31:847-67.
3. Brutsaert DL. Cardiac endothelial-myocardial signaling: its role in cardiac growth, contractile performance, and rhythmicity. *Physiological Rev* 2003;83:59-115.
4. Hsieh PC, Davis ME, Lisowski LK, Lee RT. Endothelial-cardiomyocyte interactions in cardiac development and repair. *Annu Rev Physiol* 2006;68:51-66.
5. Paulus WJ, Vantrimpont PJ, Shah AM. Paracrine coronary endothelial control of left ventricular function in humans. *Circulation* 1995;92:2119-26.
6. MacCarthy PA, Grieve DJ, Li JM, Dunster C, Kelly FJ, Shah AM. Impaired endothelial regulation of ventricular relaxation in cardiac hypertrophy: role of reactive oxygen species and NADPH oxidase. *Circulation* 2001;104:2967-74.
7. Lassegue B, San Martin A, Griendling KK. Biochemistry, physiology, and pathophysiology of NADPH oxidases in the cardiovascular system. *Circ Res* 2012;110:1364-90.
8. Murdoch CE, Alom-Ruiz SP, Wang M et al. Role of endothelial Nox2 NADPH oxidase in angiotensin II-induced hypertension and vasomotor dysfunction. *Basic Res Cardiol* 2011;106:527-38.
9. Foteinos G, Hu Y, Xiao Q, Metzler B, Xu Q. Rapid endothelial turnover in atherosclerosis-prone areas coincides with stem cell repair in apolipoprotein E-deficient mice. *Circulation* 2008;117:1856-63.

10. Ray R, Murdoch CE, Wang M et al. Endothelial Nox4 NADPH oxidase enhances vasodilatation and reduces blood pressure in vivo. *Arterioscler Thromb Vasc Biol* 2011;31:1368-76.
11. Zhang M, Brewer AC, Schroder K et al. NADPH oxidase-4 mediates protection against chronic load-induced stress in mouse hearts by enhancing angiogenesis. *Proc Natl Acad Sci U S A* 2010;107:18121-6.
12. Bendall JK, Cave AC, Heymes C, Gall N, Shah AM. Pivotal role of a gp91(phox)-containing NADPH oxidase in angiotensin II-induced cardiac hypertrophy in mice. *Circulation* 2002;105:293-6.
13. Burns SO, Killock DJ, Moulding DA et al. A congenital activating mutant of WASp causes altered plasma membrane topography and adhesion under flow in lymphocytes. *Blood* 2010;115:5355-65.
14. Zeisberg EM, Tarnavski O, Zeisberg M et al. Endothelial-to-mesenchymal transition contributes to cardiac fibrosis. *Nat Med* 2007;13:952-61.
15. Lemmens K, Segers VF, Demolder M, De Keulenaer GW. Role of neuregulin-1/ErbB2 signaling in endothelium-cardiomyocyte cross-talk. *J Biol Chem* 2006;281:19469-77.
16. Shiojima I, Sato K, Izumiya Y et al. Disruption of coordinated cardiac hypertrophy and angiogenesis contributes to the transition to heart failure. *J Clin Invest* 2005;115:2108-18.
17. Frey RS, Ushio-Fukai M, Malik AB. NADPH oxidase-dependent signaling in endothelial cells: role in physiology and pathophysiology. *Antiox Redox Signal* 2009;11:791-810.

18. Wang HD, Xu S, Johns DG et al. Role of NADPH oxidase in the vascular hypertrophic and oxidative stress response to angiotensin II in mice. *Circ Res* 2001;88:947-53.
19. Li JM, Fan LM, Christie MR, Shah AM. Acute tumor necrosis factor alpha signaling via NADPH oxidase in microvascular endothelial cells: role of p47phox phosphorylation and binding to TRAF4. *Mol Cell Biol* 2005;25:2320-30.
20. Hwang J, Saha A, Boo YC et al. Oscillatory shear stress stimulates endothelial production of O<sub>2</sub><sup>-</sup> from p47phox-dependent NAD(P)H oxidases, leading to monocyte adhesion. *J Biol Chem* 2003;278:47291-8.
21. Cathcart MK. Regulation of superoxide anion production by NADPH oxidase in monocytes/macrophages: contributions to atherosclerosis. *Arterioscler Thromb Vasc Biol* 2004;24:23-8.
22. Liu J, Yang F, Yang XP, Jankowski M, Pagano PJ. NAD(P)H oxidase mediates angiotensin II-induced vascular macrophage infiltration and medial hypertrophy. *Arterioscler Thromb Vasc Biol* 2003;23:776-82.
23. Widyantoro B, Emoto N, Nakayama K et al. Endothelial cell-derived endothelin-1 promotes cardiac fibrosis in diabetic hearts through stimulation of endothelial-to-mesenchymal transition. *Circulation* 2010;121:2407-18.
24. Lam CS, Brutsaert DL. Endothelial dysfunction: a pathophysiologic factor in heart failure with preserved ejection fraction. *J Am Coll Cardiol* 2012;60:1787-9.
25. Westermann D, Lindner D, Kasner M et al. Cardiac inflammation contributes to changes in the extracellular matrix in patients with heart failure and normal ejection fraction. *Circ Heart Fail* 2011;4:44-52.



## FIGURE LEGENDS

### **Figure 1. Representative myocardial sections showing cardiac hypertrophy and fibrosis.**

Top, cardiomyocyte borders outlined using wheat-germ agglutinin. Scale bars 50  $\mu\text{m}$ .

Bottom, Picrosirius Red-stained sections. Scale bars 100  $\mu\text{m}$ . Sal = saline.

### **Figure 2. Inflammatory cell infiltration *in vivo* and endothelial-leukocyte interaction *in vitro*.**

(A,B) Mean number of CD45<sup>+</sup> and Mac3<sup>+</sup> cells/field, respectively. (C) Number of leukocytes adherent to WT or TG EC with or without AngII treatment (0.1 $\mu\text{M}$ , 4 hr) in an *in vitro* assay. (D) VCAM-1 protein levels in WT and TG EC treated with AngII.

Representative Western blots at the top, mean data below. (E) Number of VCAM-1 positive vessels in myocardial sections from WT and TG. (F) Capillary density in myocardial sections from WT and TG (capillaries/ $\text{mm}^2$ ). \*  $P < 0.05$  by 2-way ANOVA; n=6-8/group.

### **Figure 3. Enhanced endothelial-mesenchymal transition in AngII-treated TG hearts.**

(A) Representative Western blots and mean data for CD31, smooth muscle actin (SMA) and collagen 1 (Coll) levels in cardiac homogenates from AngII-treated WT or TG mice. Actin was used as a loading control. \*,  $P < 0.05$ ; n=4/group. (B) Myocardial sections co-stained for CD31 (red) and SMA (green) and imaged by confocal microscopy. Yellow color indicates co-localisation of endothelial and mesenchymal markers. Scale bar 10  $\mu\text{m}$ . Panels to the right show pixel-by-pixel fluorescence intensities of the two fluorophores along the line indicated in the top panel (TG AngII) and confirm a significant overlapping pattern between CD31 and SMA, indicative of endothelial-mesenchymal transition.

### **Figure 4. Effect of NOX2 on endothelial-mesenchymal transition in cultured endothelial cells.**

(A) Representative Western blots for CD144, CD31, FSP1, smooth muscle actin (SMA) and NOX2 levels in EC overexpressing NOX2 or  $\beta$ -gal and treated with AngII (100 nmol/L) or vehicle (Con). GAPDH was a loading control. (B) Mean data for CD144 and SMA protein levels. Similar results were observed for CD31 and FSP1. \*,  $P < 0.05$ ; n=4/group.

**Table 1. Cardiac function assessed by LV pressure-volume loops in WT and TG mice.**

	WT Saline	WT ANGII	TG Saline	TG ANGII
Heart rate (bpm)	552±9	509±17	543±14	543±12
LVEDP (mmHg)	7.5±0.7	5.0±0.8	6.9±1.1	8.1±2.3
dP/dtmax (mmHg/min)	7804±459	8348±365	7497±529	8098±435
dP/dtmin (mmHg/min)	-6497±406	-6654±417	-6600±544	-6843±604
Stroke volume (μL)	16.7±0.9	22.8±1.7**	15.4±1	16.3±1.6**
End systolic volume (μL)	17.8±1.1	21.9±3.6	17.4±1.4	13.7±0.9‡‡
End diastolic volume (μL)	29.5±1.7	39.9±3.7**	29.0±1.4	25.3±1.5‡‡‡
Ejection fraction (%)	53±1	57±5	52±4	60±4
Cardiac output (mL/min/g)	105.5±8.1	112.0±11.5	91.2±5.0	82.3±8.2‡
Stroke work (mmHg.μl/g)	14472±1280	17216±1888	12365±865	11569±1324‡
Ea (mmHg/μL)	5.7±0.2	4.6±0.4	5.9±0.4	7.0±1.0‡‡
Tau-Weiss (ms)	7.4±0.4	6.2±0.6	7.4±0.7	7.8±1.2
dP/dtmax/EDV	261.9±39.6	258.8±32.0	273.9±73.0	334.8±71.1
Ees	4.28±0.38	8.91±0.69***	4.78±0.40	8.91±1.28***
EDPVR	0.22±0.03	0.25±0.04	0.20±0.02	0.46±0.11** ‡
Number	10	10	9	8

Ea, arterial elastance; EDV, end-diastolic volume; Ees, end-systolic elastance; EDPVR, end-diastolic pressure volume relation. \*\*,\*\*\* = P<0.01, 0.001 respectively for effect of AngII within the group. #,##,### = P<0.05, 0.01, 0.001 respectively for comparison of TG AngII group versus WT AngII group.




Fig. 1




Fig. 2




Fig. 3




Fig. 4

## SUPPLEMENTAL MATERIAL

### **Endothelial NADPH oxidase-2 promotes interstitial cardiac fibrosis and diastolic dysfunction through pro-inflammatory effects and endothelial-mesenchymal transition.**

Colin E Murdoch PhD, Sanjay Chaubey MBBCh, Lingfang Zeng PhD, Bin Yu PhD, Aleksander Ivetic PhD, \*Davy Vanhoutte PhD, \*Stephane Heymans MD, PhD, David J Grieve PhD, Alison C Cave PhD, Alison C Brewer PhD, Min Zhang MD, PhD, Ajay M Shah MD, FMedSci.

## **METHODS**

### ***Animal studies.***

All procedures were performed in accordance with the 'Guidance on the Operation of the Animals' (Scientific Procedures) Act, 1986 (UK Home Office). Transgenic mice with endothelium-targeted overexpression of Nox2 (TG) and global Nox2 knockout mice, both on a C57/BL6 background, were described previously.<sup>1,2</sup> Male Nox2 TG were compared with matched wild-type littermates (WT).

Chimeric mice were generated by bone marrow (BM) transplantation as previously described.<sup>3</sup> 6-8 week old mice received a lethal dose of whole-body irradiation (10Gy, 15min) and 24 hours later,  $0.5 \times 10^7$  BM cells isolated from donor mice were administered by tail vein injection. Donor mice were killed by cervical dislocation, their femurs were removed aseptically and a single cell suspension was prepared by flushing the marrow cavities. Cells were washed twice in HBSS and resuspended  $1 \times 10^7$  cells per ml. Mice were allowed to recover for at least 4 weeks before further study.

AngII (1.1mg/kg/day) or saline vehicle was infused via osmotic minipumps (Model 1002, Alzet, Cupertino, CA) that were implanted subcutaneously under isoflurane anesthesia. Osmotic pumps containing saline were used as controls.

Blood pressure was measured either by ambulatory radiotelemetry (PA-C10, Data Sciences International) or tail-cuff plethysmography, as described previously.<sup>4</sup> Tail-cuff

plethysmography was performed after 3 training sessions in conscious restrained mice, using a Kent Scientific system (XBP 1000; Torrington, CT).

LV pressure-volume relations were measured with a 1.4F micromanometer/microconductance catheter system (SPR-839, Millar Instruments, Houston, TX) introduced retrogradely into the LV via the right carotid artery under 2% isoflurane anesthesia.<sup>5</sup> Body temperature was maintained at 37°C to ensure heart rates of approximately 500 bpm. After stabilization, pressure volume loops were obtained via a Aria pressure-volume system (Millar, Houston, Tex) coupled to a Powerlab/8SP with Chart Software (ADInstruments, UK). Analysis was conducted using Chart and pressure volume analysis software (PVAN 3.3 Millar Instruments). The correction factor ( $\alpha$ ) was obtained from the ratio of cardiac output obtained from echocardiographic measurements (product of aortic blood flow and aortic cross sectional area) and the catheter cardiac output. Parallel conductance was obtained from a series of loops obtained during iv bolus injection of hypertonic saline via the jugular vein.

Echocardiography was performed under 2% isoflurane anesthesia at heart rates >400 bpm, using either a Sonos 5500 system with a 15 MHz linear transducer (Philips, Bothell, WA) or a Vevo 2100 with a 30MHz linear array transducer (Visualsonics, Toronto, CA). Temperature was maintained at 37°C and heart rates were over 400 bpm. Two-dimensional and M-mode images were obtained in parasternal long axis and short axis views respectively, the latter being measured at the level of the papillary muscles.

### ***Histology and immunohistochemistry.***

Hearts were arrested in diastole by perfusion with 5M KCL at 115 mmHg. Tissue was fixed in PFA (1%) and underwent normal histological preparation into paraffin blocks and onto slides. 6  $\mu$ m sections were then deparaffinized and rehydrated through a series of EtOH concentrations. Cardiomyocyte area and cardiac fibrosis were quantified in sections stained with wheat-germ agglutinin and Picrosirius Red respectively.<sup>5</sup> The collagen content was quantified as the percentage of total LV area under polarized light. Cardiomyocyte membranes and capillaries were co-stained using Wheat Germ Agglutinin (WGA) and Isolectin B4, conjugated to TRITC and FITC respectively (Vector RL-1022; Vector, USA). Confocal images were obtained on a Leica laser scanning confocal microscope (TCS-SP5) at



640x magnification. The numbers of capillaries per mm<sup>2</sup> were calculated from 10 different fields (~90 capillaries/field) using AxioVision v4.6 (Carl Zeiss, Germany).

Cryosections were used for immunostaining with antibodies against CD45 (BD, UK), Mac3 (Abcam), CD31/PECAM-1 (Chemicon International),  $\alpha$ -SMA (Abcam) and VCAM-1 (R&D). 4'-6-Diamidino-2-phenylindole (DAPI; Sigma) was used for nuclear staining. Sections were then incubated with HRP-labeled secondary antibody and streptavidin-HRP (Perkin Elmer) signal amplification was used. Visualization was conducted by incubation with diaminobenzidine (DAKO kit SK4100, Vector Laboratories) followed by Harris counterstaining. CD45 or MAC3-positive cells were counted using semi-automated software (AxioVision v4.6, Carl Zeiss, Germany) and expressed as counts per mm<sup>2</sup> averaged over 10 fields. VCAM-1 positive blood vessels were counted and expressed as % of the total number of blood vessels per heart section.

#### ***Leukocyte adhesion assay.***

Coronary microvascular endothelial cells (CMEC) were isolated from hearts of 6-8 week old mice and used at passages 2-3.<sup>4</sup> A flow chamber (Glycotech, Maryland) was placed onto confluent CMEC which had been incubated with or without AngII (0.1 $\mu$ M, 4h). BM cells isolated from WT mice and labeled with Celltracker™ (Invitrogen) were added to the perfusate which superfused at a rate of 1.5 dynes/cm<sup>2</sup> for 30 min.<sup>6</sup> Non-bound leukocytes were washed away before visualization using an inverted microscope. The number of recruited leukocytes was counted in 8-10 fields from 3 separate isolations.

#### ***EndMT assays.***

Human aortic endothelial cells (HAEC) were purchased from ATCC and maintained in gelatine-coated flasks in medium M199 supplemented with 10% FBS (Invitrogen), 1 ng/ml  $\beta$ -endothelial cell growth factor (Sigma), 3  $\mu$ g/ml endothelial cell growth supplement from bovine neural tissues (Sigma), 2.5  $\mu$ g/ml thymidine (Sigma), 10 U/ml heparin (Sigma) and 100 U/ml penicillin-streptomycin in a 5% CO<sub>2</sub> humidified incubator. HAECs were infected with Ad. $\beta$ Gal or Ad.NoX2 virus at MOI=20 for 6 hr in the presence of complete growth medium, then incubated with human endothelial serum free medium (Invitrogen) in the absence or presence of 100 nmol/L AngII. The medium was refreshed every day.

Immunofluorescent staining was performed 48 hr post virus infection while Western blotting was performed 5 days post infection.

Cell slides were fixed with methanol at room temperature for 20min; washed twice with PBS, 5min each; permeabilized with 0.01% Triton X-100/PBS for 15 min, and then washed with PBS three times for 5min each. Slides were blocked with diluted normal donkey serum (1:20 in PBS) at room temperature for 30 min, incubated with rabbit anti-VE-cadherin (1:100 in diluted donkey serum, 2158, Cell Signaling) and mouse anti-alpha smooth muscle actin (1:100 in diluted donkey serum, ab683, Abcam) at 4°C overnight, followed by washing three times with PBS, 10min each. The slides were then incubated with donkey anti-Rabbit IgG-Alexa569 (1:1000 in diluted donkey serum, DAKO) and donkey anti-mouse IgG-Alexa488 (1:1000 in diluted donkey serum, DAKO) at room temperature for 45 min, followed by washing three times with PBS, 10 min each. The slides were counterstained with DAPI at room temperature for 5min, followed by washing twice with PBS, 5min each. Fluorescent mounting medium and coverslip were applied and images were taken under an Olympus 1X81-2 fluorescent microscope with Velocity software by PerkinElmer.

### ***Endothelial cell-fibroblast co-culture***

HAECs were infected with Ad.βGal or Ad.NoX2 virus (MOI=20) as described above for EMT assays. 24 hrs post-infection, the HAECs were detached with trypsin and seeded into the inserts of 100 mm<sup>2</sup> Transwell culture dishes (Sigma) in EC medium. In parallel, human F153 fibroblasts (ATCC) were seeded in 100 mm<sup>2</sup> dishes in F-12/DMEM medium supplemented with 10% FBS and 100 U/ml penicillin-streptomycin. Six millilitre human endothelial serum free medium (SFM, Life Technologies) with or without 100 nmol/L AngII was added into the HAECs-containing inserts and F153-containing dishes, respectively. One HAEC-containing insert was put into one F153-containing dish to form a co-culture system. 24 hrs later, the cells were collected separately. One third of the F153 cells were used to extract cellular total RNA for qPCR analyses with primer sets of 5'-GCTCGTGGAAATGATGGTGC-3' and 5'-ACCCTGGGGACCTTCAGAG-3' for pro-collagen I and 5'-CACAACCTGGGACGACATGGAG-3' and 5'-TTCATGAGGTAGTCAGTCTGG-3' for β-actin as loading control. Two-thirds of the cells were lysed for Western blot analysis.

### ***ROS measurement***

NADPH oxidase activity in homogenised left ventricle (50µg) was assessed by lucigenin (5 µM)-enhanced chemiluminescence on a plate luminometer (Lucy 1, Rosys Anthos, Wals, Austria) as previously described.<sup>7</sup> All samples were analyzed in triplicate and the level of superoxide production was calculated from the average area under the curve. In some experiments, one of the following inhibitors was pre-incubated with the cell homogenate for 20 minutes prior to addition of NADPH: the flavoprotein inhibitor, diphenyleneiodonium (DPI, 10 µM); a NOS inhibitor, N<sup>G</sup>-Nitro-L-arginine methyl ester hydrochloride (L-NAME, 100 µM); a mitochondrial electron transport chain inhibitor, rotenone (50 µM); a superoxide scavenger, tiron (10 mM); a putative NADPH oxidase inhibitor, apocynin (10 µM); or a xanthine oxidase inhibitor, oxypurinol (100 µM).

ROS production in dihydroethidine-loaded BM mononuclear cells was quantified by flow cytometry.

### ***Real-time RT-PCR***

Quantification of mRNA expression was conducted using the Applied Biosystems 7000 sequence detection system (Applied Biosystems, UK) with SYBR Green and the comparative Ct method, with GAPDH levels used for normalization. RNA was isolated using an SV total RNA isolation kit (Promega, UK) and cDNA synthesised using avian myeloblastosis virus reverse transcriptase (AMV; Promega, UK) at 42°C for 90 minutes. Forward and reverse primer sequences were as follow (all 5'-3'): GAPDH: 5'-CGTGCCGCCTGGAGAA, 5'-CCCTCAGATGCCTGCTTCAC; Actin, 5'-CTGGCACCCAGCACAATG, 5'-GCCGATCCACACGGAGTACT; Procollagen I, 5'-CCTCAGGGTATTGCTGGACAAC, 5'-TTGATCCAGAAGGACCTTGTTG; MCP1, 5'-TGTAGTTTTTGTCCACCAAGCTCAAG, 5'-GTAGGTTCTGATCTCATTTGGTTCC

### ***Western blot***

Cells (75ml flask) were scratched off in the presence of medium and collected by centrifugation at 1000rpm at 4°C for 5min. The cell pellets were washed twice with ice cold PBS and resuspended in 200µl lysis buffer (20mM Tris-Cl, pH7.5, 120mM NaCl, 1mM EDTA, 1% Triton X-100 plus protease inhibitors) followed by sonication for 6 sec and incubation on ice for 45 min. Supernatant was recovered after spinning down at 10,000rpm at 4°C for

5min and protein concentration was detected using Bio-Rad reagents. 25 µg cell lysate was applied to SDS-PAGE, followed by transfer to PVDF membrane and incubation with primary antibodies against FSP1 (ab41532, Abcam), α-SMA (ab683, Abcam), PECAM-1 (CD31, sc-1506, Santa Cruz), CD144/ VE-cadherin (Santa Cruz ), Nox2 (BD), GAPDH (sc-25778, Santa Cruz) and HRP-conjugated secondary antibody (DAKO). The bound antibodies were revealed by ECL reagents (DAKO) and X-ray film.

### **Statistics**

Data are expressed as mean±SEM. Comparisons were made by 2-way ANOVA with Bonferroni post-hoc testing, 1-way ANOVA or Student's t-test as appropriate. Analyses were performed on GraphPad Prism (v4.03 for Windows, San Diego, CA). P<0.05 was considered significant.

### **References**

1. Murdoch CE, Alom-Ruiz SP, Wang M, Zhang M, Walker S, Yu B, Brewer A, Shah AM. Role of endothelial Nox2 NADPH oxidase in angiotensin ii-induced hypertension and vasomotor dysfunction. *Basic Res Cardiol.* 2011;106:527-538
2. Pollock JD, Williams DA, Gifford MA, Li LL, Du X, Fisherman J, Orkin SH, Doerschuk CM, Dinauer MC. Mouse model of X-linked chronic granulomatous disease, an inherited defect in phagocyte superoxide production. *Nature Genetics.* 1995;9:202-209
3. Foteinos G, Hu Y, Xiao Q, Metzler B, Xu Q. Rapid endothelial turnover in atherosclerosis-prone areas coincides with stem cell repair in apolipoprotein E-deficient mice. *Circulation.* 2008;117:1856-1863
4. Ray R, Murdoch CE, Wang M, Santos CX, Zhang M, Alom-Ruiz S, Anilkumar N, Ouattara A, Cave AC, Walker SJ, Grieve DJ, Charles RL, Eaton P, Brewer AC, Shah AM. Endothelial Nox4 NADPH oxidase enhances vasodilatation and reduces blood pressure in vivo. *Arterioscler Thromb Vasc Biol.* 2011;31:1368-1376
5. Zhang M, Brewer AC, Schroder K, Santos CX, Grieve DJ, Wang M, Anilkumar N, Yu B, Dong X, Walker SJ, Brandes RP, Shah AM. NADPH oxidase-4 mediates protection against chronic

load-induced stress in mouse hearts by enhancing angiogenesis. *Proc Natl Acad Sci U S A*. 2010;107:18121-18126

6. Burns SO, Killock DJ, Moulding DA, Metelo J, Nunes J, Taylor RR, Forge A, Thrasher AJ, Ivetic A. A congenital activating mutant of WASP causes altered plasma membrane topography and adhesion under flow in lymphocytes. *Blood*. 2010;115:5355-5365

7. Bendall JK, Cave AC, Heymes C, Gall N, Shah AM. Pivotal role of a gp91(phox)-containing NADPH oxidase in angiotensin II-induced cardiac hypertrophy in mice. *Circulation*. 2002;105:293-296

**Supplementary Figure 1. Effect of endothelial NOX2 on AngII-induced cardiac hypertrophy and fibrosis.** (A) Systolic blood pressure (SBP) after saline or AngII (1.1mg/kg/day) infusion. Average SBP  $\pm$ SEM over a 12-hour dark period (active) is shown. \*,  $P < 0.05$  AngII vs saline by repeated measures ANOVA.  $n = 6$ /group. (B) LV NADPH oxidase activity in wild-type and TG treated with saline or AngII. (C) LV/body weight (BW) ratio. (D) Cardiomyocyte area. (E) % fibrosis. (F) LV collagen I mRNA levels. \*  $P < 0.05$ , \*\*\*  $P < 0.001$  by 2-way ANOVA;  $n = 6-8$ /group.

**Supplementary Figure 2: NADPH-dependent superoxide generation in LV homogenates.** Lucigenin (5 $\mu$ M)-enhanced chemiluminescence was measured in LV homogenate from WT and TG mice with saline or AngII infusion. The chemiluminescence signal was measured over 15 min in the presence or absence of DPI (10  $\mu$ M), L-NAME (100  $\mu$ M), Tiron (20 mM), oxypurinol (100  $\mu$ M), rotenone (100  $\mu$ M) or apocynin (100  $\mu$ M). Results were quantified as area under the curve and presented as arbitrary integrated light units. Data is shown for the effects of inhibitors in the AngII-treated TG group. The inhibitors had similar effects in the other 3 groups. \*\*,  $P < 0.01$  by 1-way ANOVA followed by Dunnett's post hoc test, comparing to control.  $n \geq 6$ /group.

**Supplementary Figure 3: Echocardiographic indices in WT and TG mice treated with AngII or saline.** (A) Interventricular septal diameter. (B) LV end diastolic diameter (LVEDD). (C) LV

end systolic diameter (LVESD). (D) Ejection fraction. (E) Heart rate (HR). \*\*,  $P < 0.05$  by 2-way ANOVA.  $n = 12/\text{group}$ .

**Supplementary Figure 4. Left ventricular function assessed by pressure-volume analysis.**

(A) Ees, end-systolic elastance. (B) LV  $dp/dt_{\max}$  normalized by end-diastolic volume (EDV). (C) LV EDV. (D) LV end-systolic volume (ESV). (E) Stroke work (SW). (F) End-diastolic pressure volume relation (EDPVR). (G) Representative steady-state pressure-volume loops from wild-type (WT) and TG mice. (H) Representative pressure-volume loops during acute preload reduction in AngII-treated WT and TG. Dashed lines at top left and bottom right of the loops show the ESPVR and EDPVR respectively. \*  $P < 0.05$  AngII vs. respective saline group; #  $P < 0.05$  WT AngII vs. TG AngII by 2-way ANOVA;  $n = 12/\text{group}$ .

**Supplementary Figure 5. Inflammatory cell infiltration *in vivo*.** (A,B) Representative sections stained for  $CD45^+$  and  $Mac3^+$  cells, respectively. Scale bars 100  $\mu\text{m}$ .

**Supplementary Figure 6: Expression of inflammatory cytokines and pro-fibrotic factors in WT and TG heart.** mRNA levels of the genes shown were measured by real-time PCR in LV from WT and TG treated with saline or AngII.  $\beta$ -actin levels were used for normalization. \*  $P < 0.05$  compared with respective saline control.  $n \geq 6/\text{group}$ .

**Supplementary Figure 7: Effects of N-acetyl cysteine (NAC) on blood pressure, cardiac hypertrophy and fibrosis.** (A) Systolic blood pressure (SBP) measured by tail cuff plethysmography. (B) Cardiac fibrosis in LV sections. (C) LV/BW ratio. (D)  $CD45^+$  cells in LV sections. NAC (5g/L) was administered in the drinking water for the whole duration of 2 week AngII infusion. \*  $P < 0.05$  vs saline treatment by repeated measure ANOVA.  $n = 6/\text{group}$  for BP and morphometric data;  $n = 4/\text{group}$  for histological data.

**Supplementary Figure 8: Expression of VCAM-1 in LV sections.** Representative examples of VCAM-1 positive blood vessels in LV sections from WT and TG mice chronically treated with AngII.

**Supplementary Figure 9: Effects of bone marrow (BM)-derived cells on blood pressure and cardiac hypertrophy.** Bone marrow chimeric mice were generated as described in the text.

(A) Superoxide production measured by flow cytometry in dihydroethidine (DHE)-loaded BM cells from chimeric mice that previously received bone marrow transfer (BMT) of WT cells (left) or Nox2 knockout (KO) cells KO (right). Phorbol ester (PMA) was used to stimulate ROS production. (B) Systolic blood pressure (SBP) measured by tail cuff plethysmography in the different experimental groups. (C) Cardiac hypertrophy assessed by LV/body weight (BW) ratio. \*,  $P < 0.05$  for the comparisons shown;  $n = 6$ /group.

**Supplementary Figure 10. Effect of NOX2 in bone marrow-derived inflammatory cells on AngII-induced fibrosis.**


(A) Representative Picrosirius Red-stained LV sections. Scale bars 100  $\mu\text{m}$ . (B,C) Bottom, mean % fibrosis. See text for details of experimental groups. Horizontal lines above the columns in panel B denote significance at  $P < 0.05$  level (\*) by 1-way ANOVA;  $n \geq 5$ /group.

**Supplementary Figure 11. Effect of NOX2 on endothelial-mesenchymal transition in cultured endothelial cells.**


(A) Representative confocal images of EC stained for CD144 and SMA. Arrow indicates cell expressing both endothelial and mesenchymal markers. (B) Representative phase contrast images of EC after AngII treatment. Arrow indicates cell with typical elongated fibroblastic phenotype.

**Supplementary Figure 12: Effects of endothelial NOX2 overexpression on fibroblasts in Transwell co-culture.**

Endothelial cells were infected with Ad.NOX2 or Ad. $\beta$ -Gal and experiments were performed with or without AngII (100 nmol/L). (A) Fibroblast pro-collagen 1 mRNA expression levels. (B) Representative Western blots showing markers of myofibroblast transformation, smooth muscle actin (SMA) and SM22. (C, D) Mean data for experiment shown in panel B.  $n = 4$ /group.




Supplementary Fig. 1




Supplementary Fig. 2







Supplementary Fig. 3



Supplementary Fig. 4




Supplementary Fig. 5




Supplementary Fig. 6


A




B




C






D





Supplementary Fig. 7




Supplementary Fig. 8




Supplementary Fig. 9



Supplementary Fig. 10



Supplementary Fig. 11



Supplementary Fig. 12



AMERICAN COLLEGE OF CARDIOLOGY FOUNDATION  
AUTHOR RESPONSIBILITIES: RELATIONSHIP WITH INDUSTRY AND FINANCIAL DISCLOSURE

In addition to disclosing relationship with industry and/or financial disclosures on the *cover letter* and *title* page of your manuscript, we ask that each author sign this form.

**To be published in the following journal (please check):**

*Journal of the American College of Cardiology(JACC)*  *JACC: Cardiovascular Interventions*  *JACC: Cardiovascular Imaging*  *JACC: Heart Failure*

Article entitled: Endothelial NADPH oxidase-2 promotes interstitial cardiac fibrosis and diastolic dysfunction through pro-inflammatory effects and endothelial-mesenchymal transition

Manuscript number: JACC110813-4952D

Corresponding Author: Prof. Shah

Corresponding author's printed name: Prof Ajay Shah

**\*Corresponding Author's Responsibility**

To avoid delay in the publication of your manuscript, **complete and submit this form as quickly as possible.**

After you have submitted your electronic form, you will receive an email confirmation that includes a PDF copy. Please print and retain for your files. **If for some reason you are unable to complete this form electronically**, please contact the JACC office directly or print this form, fill in the spaces manually, and email, fax, or post mail a copy to:

*Journal of the American College of Cardiology(JACC)*  
*JACC: Cardiovascular Interventions*  
*JACC: Cardiovascular Imaging*  
*JACC: Heart Failure*  
3655 Nobel Drive, Suite 630  
San Diego, California 92122-1005  
Fax: (858) 558-3148  
Email: [jaccsd@acc.org](mailto:jaccsd@acc.org) [jaccint@acc.org](mailto:jaccint@acc.org) [jaccimg@acc.org](mailto:jaccimg@acc.org) [jacchf@acc.org](mailto:jacchf@acc.org)

Complete the section below, type your name into the Type Name box and input the date. *By submitting this form, you attest to being an author on the paper.*

Type Name: Colin Murdoch Date: 12/05/13

Title and Company (if employer representative): King's College London

I have **no relationship with industry** to disclose.

I have **no financial information** to disclose.

local\_p\_id: 348571

time: 1386283826

ip address: 77.98.27.182





AMERICAN COLLEGE OF CARDIOLOGY FOUNDATION  
AUTHOR RESPONSIBILITIES: RELATIONSHIP WITH INDUSTRY AND FINANCIAL DISCLOSURE

In addition to disclosing relationship with industry and/or financial disclosures on the *cover letter* and *title* page of your manuscript, we ask that each author sign this form.

**To be published in the following journal (please check):**

*Journal of the American College of Cardiology(JACC)*  *JACC: Cardiovascular Interventions*  *JACC: Cardiovascular Imaging*  *JACC: Heart Failure*

Article entitled: Endothelial NADPH oxidase-2 promotes interstitial cardiac fibrosis and diastolic dysfunction through pro-inflammatory effects and endothelial-mesenchymal transition

Manuscript number: JACC110813-4952DR

Corresponding Author: Prof. Shah

Corresponding author's printed name: Prof Shah

**\*Corresponding Author's Responsibility**

To avoid delay in the publication of your manuscript, **complete and submit this form as quickly as possible.**

After you have submitted your electronic form, you will receive an email confirmation that includes a PDF copy. Please print and retain for your files. **If for some reason you are unable to complete this form electronically**, please contact the JACC office directly or print this form, fill in the spaces manually, and email, fax, or post mail a copy to:

*Journal of the American College of Cardiology(JACC)*  
*JACC: Cardiovascular Interventions*  
*JACC: Cardiovascular Imaging*  
*JACC: Heart Failure*  
3655 Nobel Drive, Suite 630  
San Diego, California 92122-1005  
Fax: (858) 558-3148  
Email: [jaccsd@acc.org](mailto:jaccsd@acc.org) [jaccint@acc.org](mailto:jaccint@acc.org) [jaccimg@acc.org](mailto:jaccimg@acc.org) [jacchf@acc.org](mailto:jacchf@acc.org)

Complete the section below, type your name into the Type Name box and input the date. *By submitting this form, you attest to being an author on the paper.*

Type Name: Mr Sanjay Chaubey Date: 8/12/13

Title and Company (if employer representative):

I have **no relationship with industry** to disclose.

I have **no financial information** to disclose.

local\_p\_id: 348572

time: 1386509954

ip address: 80.47.139.139



AMERICAN COLLEGE OF CARDIOLOGY FOUNDATION  
AUTHOR RESPONSIBILITIES: RELATIONSHIP WITH INDUSTRY AND FINANCIAL DISCLOSURE

In addition to disclosing relationship with industry and/or financial disclosures on the *cover letter* and *title* page of your manuscript, we ask that each author sign this form.

**To be published in the following journal (please check):**

*Journal of the American College of Cardiology(JACC)*  *JACC: Cardiovascular Interventions*  *JACC: Cardiovascular Imaging*  *JACC: Heart Failure*

Article entitled: Endothelial NADPH oxidase-2 promotes interstitial cardiac fibrosis and diastolic dysfunction through pro-inflammatory effects and endothelial-mesenchymal transition

Manuscript number: JACC110813-4952D

Corresponding Author: Prof. Shah

Corresponding author's printed name: Ajay Shah

**\*Corresponding Author's Responsibility**

To avoid delay in the publication of your manuscript, **complete and submit this form as quickly as possible.**

After you have submitted your electronic form, you will receive an email confirmation that includes a PDF copy. Please print and retain for your files. **If for some reason you are unable to complete this form electronically**, please contact the JACC office directly or print this form, fill in the spaces manually, and email, fax, or post mail a copy to:

*Journal of the American College of Cardiology(JACC)*  
*JACC: Cardiovascular Interventions*  
*JACC: Cardiovascular Imaging*  
*JACC: Heart Failure*  
3655 Nobel Drive, Suite 630  
San Diego, California 92122-1005  
Fax: (858) 558-3148  
Email: [jaccsd@acc.org](mailto:jaccsd@acc.org) [jaccint@acc.org](mailto:jaccint@acc.org) [jaccimg@acc.org](mailto:jaccimg@acc.org) [jacchf@acc.org](mailto:jacchf@acc.org)

Complete the section below, type your name into the Type Name box and input the date. *By submitting this form, you attest to being an author on the paper.*

Type Name: Lingfang Zeng Date: 06 Dec 2013

Title and Company (if employer representative): Senior Lecturer, KCL

I have **no relationship with industry** to disclose.

I have **no financial information** to disclose.

local\_p\_id: 348573

time: 1386321651

ip address: 159.92.203.113



AMERICAN COLLEGE OF CARDIOLOGY FOUNDATION  
AUTHOR RESPONSIBILITIES: RELATIONSHIP WITH INDUSTRY AND FINANCIAL DISCLOSURE

In addition to disclosing relationship with industry and/or financial disclosures on the *cover letter* and *title* page of your manuscript, we ask that each author sign this form.

**To be published in the following journal (please check):**

*Journal of the American College of Cardiology(JACC)*  *JACC: Cardiovascular Interventions*  *JACC: Cardiovascular Imaging*  *JACC: Heart Failure*

Article entitled: Endothelial NADPH oxidase-2 promotes interstitial cardiac fibrosis and diastolic dysfunction through pro-inflammatory effects and endothelial-mesenchymal transition

Manuscript number: JACC110813-4952DR

Corresponding Author: Prof. Shah

Corresponding author's printed name: Ajay Shah

**\*Corresponding Author's Responsibility**

To avoid delay in the publication of your manuscript, **complete and submit this form as quickly as possible.**

After you have submitted your electronic form, you will receive an email confirmation that includes a PDF copy. Please print and retain for your files. **If for some reason you are unable to complete this form electronically**, please contact the JACC office directly or print this form, fill in the spaces manually, and email, fax, or post mail a copy to:

*Journal of the American College of Cardiology(JACC)*  
*JACC: Cardiovascular Interventions*  
*JACC: Cardiovascular Imaging*  
*JACC: Heart Failure*  
3655 Nobel Drive, Suite 630  
San Diego, California 92122-1005  
Fax: (858) 558-3148  
Email: [jaccsd@acc.org](mailto:jaccsd@acc.org) [jaccint@acc.org](mailto:jaccint@acc.org) [jaccimg@acc.org](mailto:jaccimg@acc.org) [jacchf@acc.org](mailto:jacchf@acc.org)

Complete the section below, type your name into the Type Name box and input the date. *By submitting this form, you attest to being an author on the paper.*

Type Name: Bin Yu Date: 01/02/14

Title and Company (if employer representative):

I have **no relationship with industry** to disclose.

I have **no financial information** to disclose.

local\_p\_id: 328554

time: 1391233779

ip address: 5.108.101.121



AMERICAN COLLEGE OF CARDIOLOGY FOUNDATION  
AUTHOR RESPONSIBILITIES: RELATIONSHIP WITH INDUSTRY AND FINANCIAL DISCLOSURE

In addition to disclosing relationship with industry and/or financial disclosures on the *cover letter* and *title* page of your manuscript, we ask that each author sign this form.

**To be published in the following journal (please check):**

*Journal of the American College of Cardiology(JACC)*  *JACC: Cardiovascular Interventions*  *JACC: Cardiovascular Imaging*  *JACC: Heart Failure*

Article entitled: Endothelial NADPH oxidase-2 promotes interstitial cardiac fibrosis and diastolic dysfunction through pro-inflammatory effects and endothelial-mesenchymal transition

Manuscript number: JACC110813-4952D

Corresponding Author: Prof. Shah

Corresponding author's printed name: Prof. Ajay Shah

**\*Corresponding Author's Responsibility**

To avoid delay in the publication of your manuscript, **complete and submit this form as quickly as possible.**

After you have submitted your electronic form, you will receive an email confirmation that includes a PDF copy. Please print and retain for your files. **If for some reason you are unable to complete this form electronically**, please contact the JACC office directly or print this form, fill in the spaces manually, and email, fax, or post mail a copy to:

*Journal of the American College of Cardiology(JACC)*  
*JACC: Cardiovascular Interventions*  
*JACC: Cardiovascular Imaging*  
*JACC: Heart Failure*  
3655 Nobel Drive, Suite 630  
San Diego, California 92122-1005  
Fax: (858) 558-3148  
Email: [jaccsd@acc.org](mailto:jaccsd@acc.org) [jaccint@acc.org](mailto:jaccint@acc.org) [jaccimg@acc.org](mailto:jaccimg@acc.org) [jacchf@acc.org](mailto:jacchf@acc.org)

Complete the section below, type your name into the Type Name box and input the date. *By submitting this form, you attest to being an author on the paper.*

Type Name: Aleksandar Ivetic Date: 06 December 2013

Title and Company (if employer representative): King's College London

I have **no relationship with industry** to disclose.

I have **no financial information** to disclose.

local\_p\_id: 348574

time: 1386290490

ip address: 91.85.76.101



AMERICAN COLLEGE OF CARDIOLOGY FOUNDATION  
AUTHOR RESPONSIBILITIES: RELATIONSHIP WITH INDUSTRY AND FINANCIAL DISCLOSURE

In addition to disclosing relationship with industry and/or financial disclosures on the *cover letter* and *title* page of your manuscript, we ask that each author sign this form.

**To be published in the following journal (please check):**

*Journal of the American College of Cardiology(JACC)*  *JACC: Cardiovascular Interventions*  *JACC: Cardiovascular Imaging*  *JACC: Heart Failure*

Article entitled: Endothelial NADPH oxidase-2 promotes interstitial cardiac fibrosis and diastolic dysfunction through pro-inflammatory effects and endothelial-mesenchymal transition

Manuscript number: JACC110813-4952DRR

Corresponding Author: Prof. Shah

Corresponding author's printed name: A. M. Shah

**\*Corresponding Author's Responsibility**

To avoid delay in the publication of your manuscript, **complete and submit this form as quickly as possible.**

After you have submitted your electronic form, you will receive an email confirmation that includes a PDF copy. Please print and retain for your files. **If for some reason you are unable to complete this form electronically**, please contact the JACC office directly or print this form, fill in the spaces manually, and email, fax, or post mail a copy to:

*Journal of the American College of Cardiology(JACC)*  
*JACC: Cardiovascular Interventions*  
*JACC: Cardiovascular Imaging*  
*JACC: Heart Failure*  
3655 Nobel Drive, Suite 630  
San Diego, California 92122-1005  
Fax: (858) 558-3148  
Email: [jaccsd@acc.org](mailto:jaccsd@acc.org) [jaccint@acc.org](mailto:jaccint@acc.org) [jaccimg@acc.org](mailto:jaccimg@acc.org) [jacchf@acc.org](mailto:jacchf@acc.org)

Complete the section below, type your name into the Type Name box and input the date. *By submitting this form, you attest to being an author on the paper.*

Type Name: simon walker Date: 27th Feb 2014

Title and Company (if employer representative):

I have **no relationship with industry** to disclose.

I have **no financial information** to disclose.

local\_p\_id: 159640

time: 1393495648

ip address: 159.92.53.195



AMERICAN COLLEGE OF CARDIOLOGY FOUNDATION  
AUTHOR RESPONSIBILITIES: RELATIONSHIP WITH INDUSTRY AND FINANCIAL DISCLOSURE

In addition to disclosing relationship with industry and/or financial disclosures on the *cover letter* and *title* page of your manuscript, we ask that each author sign this form.

**To be published in the following journal (please check):**

*Journal of the American College of Cardiology(JACC)*  *JACC: Cardiovascular Interventions*  *JACC: Cardiovascular Imaging*  *JACC: Heart Failure*

Article entitled: Endothelial NADPH oxidase-2 promotes interstitial cardiac fibrosis and diastolic dysfunction through pro-inflammatory effects and endothelial-mesenchymal transition

Manuscript number: JACC110813-4952D

Corresponding Author: Prof. Shah

Corresponding author's printed name: Ajay Shah

**\*Corresponding Author's Responsibility**

To avoid delay in the publication of your manuscript, **complete and submit this form as quickly as possible.**

After you have submitted your electronic form, you will receive an email confirmation that includes a PDF copy. Please print and retain for your files. **If for some reason you are unable to complete this form electronically**, please contact the JACC office directly or print this form, fill in the spaces manually, and email, fax, or post mail a copy to:

*Journal of the American College of Cardiology(JACC)*  
*JACC: Cardiovascular Interventions*  
*JACC: Cardiovascular Imaging*  
*JACC: Heart Failure*  
3655 Nobel Drive, Suite 630  
San Diego, California 92122-1005  
Fax: (858) 558-3148  
Email: [jaccsd@acc.org](mailto:jaccsd@acc.org) [jaccint@acc.org](mailto:jaccint@acc.org) [jaccimg@acc.org](mailto:jaccimg@acc.org) [jacchf@acc.org](mailto:jacchf@acc.org)

Complete the section below, type your name into the Type Name box and input the date. *By submitting this form, you attest to being an author on the paper.*

Type Name: Davy Vanhoutte Date: 12/05/2013

Title and Company (if employer representative): PhD

I have **no relationship with industry** to disclose.

I have **no financial information** to disclose.

local\_p\_id: 319586

time: 1386286324

ip address: 205.142.197.33



AMERICAN COLLEGE OF CARDIOLOGY FOUNDATION  
AUTHOR RESPONSIBILITIES: RELATIONSHIP WITH INDUSTRY AND FINANCIAL DISCLOSURE

In addition to disclosing relationship with industry and/or financial disclosures on the *cover letter* and *title* page of your manuscript, we ask that each author sign this form.

**To be published in the following journal (please check):**

*Journal of the American College of Cardiology(JACC)*  *JACC: Cardiovascular Interventions*  *JACC: Cardiovascular Imaging*  *JACC: Heart Failure*

Article entitled: Endothelial NADPH oxidase-2 promotes interstitial cardiac fibrosis and diastolic dysfunction through pro-inflammatory effects and endothelial-mesenchymal transition

Manuscript number: JACC110813-4952DR

Corresponding Author: Prof. Shah

Corresponding author's printed name: Prof Shah

**\*Corresponding Author's Responsibility**

To avoid delay in the publication of your manuscript, **complete and submit this form as quickly as possible.**

After you have submitted your electronic form, you will receive an email confirmation that includes a PDF copy. Please print and retain for your files. **If for some reason you are unable to complete this form electronically**, please contact the JACC office directly or print this form, fill in the spaces manually, and email, fax, or post mail a copy to:

*Journal of the American College of Cardiology(JACC)*  
*JACC: Cardiovascular Interventions*  
*JACC: Cardiovascular Imaging*  
*JACC: Heart Failure*  
3655 Nobel Drive, Suite 630  
San Diego, California 92122-1005  
Fax: (858) 558-3148  
Email: [jaccsd@acc.org](mailto:jaccsd@acc.org) [jaccint@acc.org](mailto:jaccint@acc.org) [jaccimg@acc.org](mailto:jaccimg@acc.org) [jacchf@acc.org](mailto:jacchf@acc.org)

Complete the section below, type your name into the Type Name box and input the date. *By submitting this form, you attest to being an author on the paper.*

Type Name: Heymans Stephane Date: 31/01/2014

Title and Company (if employer representative):

I have **no relationship with industry** to disclose.

I have **no financial information** to disclose.

local\_p\_id: 348576

time: 1391171873

ip address: 87.64.74.11



AMERICAN COLLEGE OF CARDIOLOGY FOUNDATION  
AUTHOR RESPONSIBILITIES: RELATIONSHIP WITH INDUSTRY AND FINANCIAL DISCLOSURE

In addition to disclosing relationship with industry and/or financial disclosures on the *cover letter* and *title* page of your manuscript, we ask that each author sign this form.

**To be published in the following journal (please check):**

*Journal of the American College of Cardiology(JACC)*  *JACC: Cardiovascular Interventions*  *JACC: Cardiovascular Imaging*  *JACC: Heart Failure*

Article entitled: Endothelial NADPH oxidase-2 promotes interstitial cardiac fibrosis and diastolic dysfunction through pro-inflammatory effects and endothelial-mesenchymal transition

Manuscript number: JACC110813-4952D

Corresponding Author: Prof. Shah

Corresponding author's printed name: Ajay M Shah

**\*Corresponding Author's Responsibility**

To avoid delay in the publication of your manuscript, **complete and submit this form as quickly as possible.**

After you have submitted your electronic form, you will receive an email confirmation that includes a PDF copy. Please print and retain for your files. **If for some reason you are unable to complete this form electronically**, please contact the JACC office directly or print this form, fill in the spaces manually, and email, fax, or post mail a copy to:

*Journal of the American College of Cardiology(JACC)*  
*JACC: Cardiovascular Interventions*  
*JACC: Cardiovascular Imaging*  
*JACC: Heart Failure*  
3655 Nobel Drive, Suite 630  
San Diego, California 92122-1005  
Fax: (858) 558-3148  
Email: [jaccsd@acc.org](mailto:jaccsd@acc.org) [jaccint@acc.org](mailto:jaccint@acc.org) [jaccimg@acc.org](mailto:jaccimg@acc.org) [jacchf@acc.org](mailto:jacchf@acc.org)

Complete the section below, type your name into the Type Name box and input the date. *By submitting this form, you attest to being an author on the paper.*

Type Name: David J Grieve Date: 5th December 2013

Title and Company (if employer representative):

I have **no relationship with industry** to disclose.

I have **no financial information** to disclose.

local\_p\_id: 262660

time: 1386276638

ip address: 31.54.173.237





AMERICAN COLLEGE OF CARDIOLOGY FOUNDATION  
AUTHOR RESPONSIBILITIES: RELATIONSHIP WITH INDUSTRY AND FINANCIAL DISCLOSURE

In addition to disclosing relationship with industry and/or financial disclosures on the *cover letter* and *title* page of your manuscript, we ask that each author sign this form.

**To be published in the following journal (please check):**

*Journal of the American College of Cardiology(JACC)*  *JACC: Cardiovascular Interventions*  *JACC: Cardiovascular Imaging*  *JACC: Heart Failure*

Article entitled: Endothelial NADPH oxidase-2 promotes interstitial cardiac fibrosis and diastolic dysfunction through pro-inflammatory effects and endothelial-mesenchymal transition

Manuscript number: JACC110813-4952D

Corresponding Author: Prof. Shah

Corresponding author's printed name: A. Shah

**\*Corresponding Author's Responsibility**

To avoid delay in the publication of your manuscript, **complete and submit this form as quickly as possible.**

After you have submitted your electronic form, you will receive an email confirmation that includes a PDF copy. Please print and retain for your files. **If for some reason you are unable to complete this form electronically**, please contact the JACC office directly or print this form, fill in the spaces manually, and email, fax, or post mail a copy to:

*Journal of the American College of Cardiology(JACC)*  
*JACC: Cardiovascular Interventions*  
*JACC: Cardiovascular Imaging*  
*JACC: Heart Failure*  
3655 Nobel Drive, Suite 630  
San Diego, California 92122-1005  
Fax: (858) 558-3148  
Email: [jaccsd@acc.org](mailto:jaccsd@acc.org) [jaccint@acc.org](mailto:jaccint@acc.org) [jaccimg@acc.org](mailto:jaccimg@acc.org) [jacchf@acc.org](mailto:jacchf@acc.org)

Complete the section below, type your name into the Type Name box and input the date. *By submitting this form, you attest to being an author on the paper.*

Type Name: Alison Cave Date: 4th December 2024

Title and Company (if employer representative):

I have **no relationship with industry** to disclose.

I have **no financial information** to disclose.

local\_p\_id: 360058

time: 1386281746

ip address: 81.106.105.39



AMERICAN COLLEGE OF CARDIOLOGY FOUNDATION  
AUTHOR RESPONSIBILITIES: RELATIONSHIP WITH INDUSTRY AND FINANCIAL DISCLOSURE

In addition to disclosing relationship with industry and/or financial disclosures on the *cover letter* and *title* page of your manuscript, we ask that each author sign this form.

**To be published in the following journal (please check):**

*Journal of the American College of Cardiology(JACC)*  *JACC: Cardiovascular Interventions*  *JACC: Cardiovascular Imaging*  *JACC: Heart Failure*

Article entitled: Endothelial NADPH oxidase-2 promotes interstitial cardiac fibrosis and diastolic dysfunction through pro-inflammatory effects and endothelial-mesenchymal transition

Manuscript number: JACC110813-4952D

Corresponding Author: Prof. Shah

Corresponding author's printed name: Ajay Shah

**\*Corresponding Author's Responsibility**

To avoid delay in the publication of your manuscript, **complete and submit this form as quickly as possible.**

After you have submitted your electronic form, you will receive an email confirmation that includes a PDF copy. Please print and retain for your files. **If for some reason you are unable to complete this form electronically**, please contact the JACC office directly or print this form, fill in the spaces manually, and email, fax, or post mail a copy to:

*Journal of the American College of Cardiology(JACC)*  
*JACC: Cardiovascular Interventions*  
*JACC: Cardiovascular Imaging*  
*JACC: Heart Failure*  
3655 Nobel Drive, Suite 630  
San Diego, California 92122-1005  
Fax: (858) 558-3148  
Email: [jaccsd@acc.org](mailto:jaccsd@acc.org) [jaccint@acc.org](mailto:jaccint@acc.org) [jaccimg@acc.org](mailto:jaccimg@acc.org) [jacchf@acc.org](mailto:jacchf@acc.org)

Complete the section below, type your name into the Type Name box and input the date. *By submitting this form, you attest to being an author on the paper.*

Type Name: Alison Brewer Date: 6/12/2013

Title and Company (if employer representative):

I have **no relationship with industry** to disclose.

I have **no financial information** to disclose.

local\_p\_id: 348578

time: 1386319882

ip address: 46.65.70.43



AMERICAN COLLEGE OF CARDIOLOGY FOUNDATION  
AUTHOR RESPONSIBILITIES: RELATIONSHIP WITH INDUSTRY AND FINANCIAL DISCLOSURE

In addition to disclosing relationship with industry and/or financial disclosures on the *cover letter* and *title* page of your manuscript, we ask that each author sign this form.

**To be published in the following journal (please check):**

*Journal of the American College of Cardiology(JACC)*  *JACC: Cardiovascular Interventions*  *JACC: Cardiovascular Imaging*  *JACC: Heart Failure*

Article entitled: Endothelial NADPH oxidase-2 promotes interstitial cardiac fibrosis and diastolic dysfunction through pro-inflammatory effects and endothelial-mesenchymal transition

Manuscript number: JACC110813-4952D

Corresponding Author: Prof. Shah

Corresponding author's printed name: Ajay M Shah

**\*Corresponding Author's Responsibility**

To avoid delay in the publication of your manuscript, **complete and submit this form as quickly as possible.**

After you have submitted your electronic form, you will receive an email confirmation that includes a PDF copy. Please print and retain for your files. **If for some reason you are unable to complete this form electronically**, please contact the JACC office directly or print this form, fill in the spaces manually, and email, fax, or post mail a copy to:

*Journal of the American College of Cardiology(JACC)*  
*JACC: Cardiovascular Interventions*  
*JACC: Cardiovascular Imaging*  
*JACC: Heart Failure*  
3655 Nobel Drive, Suite 630  
San Diego, California 92122-1005  
Fax: (858) 558-3148  
Email: [jaccsd@acc.org](mailto:jaccsd@acc.org) [jaccint@acc.org](mailto:jaccint@acc.org) [jaccimg@acc.org](mailto:jaccimg@acc.org) [jacchf@acc.org](mailto:jacchf@acc.org)

Complete the section below, type your name into the Type Name box and input the date. *By submitting this form, you attest to being an author on the paper.*

Type Name: Min Zhang Date: 06 Dec. 2013

Title and Company (if employer representative):

I have **no relationship with industry** to disclose.

I have **no financial information** to disclose.

local\_p\_id: 348579

time: 1386290367

ip address: 117.136.44.1



**AMERICAN COLLEGE OF CARDIOLOGY FOUNDATION  
CORRESPONDING AUTHOR RESPONSIBILITY AND COPYRIGHT TRANSFER  
AGREEMENT**

**Journal** Journal of the American College of Cardiology  
**Manuscript #** JACC110813-4952D  
**Title** Endothelial NADPH oxidase-2 promotes interstitial cardiac fibrosis and diastolic dysfunction through pro-inflammatory effects and endothelial-mesenchymal transition  
**Authors** Colin Murdoch, Sanjay Chaubey, Lingfang Zeng, Bin Yu, Aleksandar Ivetic, Simon Walker, Davy Vanhoutte, Stephane Heymans, David Grieve, Alison Cave, Alison Brewer, Min Zhang, and Ajay Shah  
**Date** December 06, 2013 10:54 GMT  
**IP Address** 94.2.190.208

Journal publishers and authors share a common interest in the protection of copyright: authors principally because they want their creative works to be protected from plagiarism and other unlawful uses, publishers because they need to protect their work and investment in the production, marketing and distribution of the published version of the article. In order to do so effectively, publishers request a formal written transfer of copyright from the author(s) for each article published. Publishers and authors are also concerned that the integrity of the official record of publication of an article (once refereed and published) be maintained, and in order to protect that reference value and validation process, we ask that authors recognize that distribution (including through the Internet/WWW or other on-line means) of the authoritative version of the article as published is best administered by the Publisher.

To avoid any delay in the publication of your image, please read the terms of this agreement, complete all required information, and electronically submit your form as quickly as possible.

**\*For corresponding author only**

- I will obtain written permission from all individuals who are listed in the Acknowledgment section of the manuscript because readers may infer their endorsement of data and conclusions.  
**(Acknowledgments no longer need to be sent to the Editorial Office.)**

I certify that:

1. All persons who have made substantial contributions in the manuscript (e.g., data collection, analysis, or writing or editing assistance), but who do not fulfill authorship criteria, are named with their specific contributions in the Acknowledgment section of the manuscript;
2. All persons named in the Acknowledgment section have provided the corresponding author with written permission to be named in the manuscript; and
3. If an Acknowledgment section is not included, no other persons have made substantial contributions to the manuscript.

Corresponding Author's Name: Ajay M Shah

**To be published in the following journal (please check):**

*Journal of the American College of Cardiology (JACC)*  *JACC: Cardiovascular Interventions*  *JACC: Cardiovascular Imaging*  *JACC: Heart Failure*

Article entitled: Endothelial NADPH oxidase-2 promotes interstitial cardiac fibrosis and diastolic dysfunction through pro-inflammatory effects and endothelial-mesenchymal transition

Manuscript number: JACC110813-4952D
Corresponding author's printed name: Prof. Shah
First author's printed name: Colin Murdoch

I hereby assign to the American College of Cardiology Foundation the copyright in the manuscript identified above and any supplemental tables, illustrations or other information submitted therewith (the "article") in all forms and media (whether now known or hereafter developed), throughout the world, in all languages, for the full term of copyright and all extensions and renewals thereof, effective when and if the article is accepted for publication. This transfer includes the right to adapt the presentation of the article for use in conjunction with computer systems and programs, including reproduction or publication in machine-readable form and incorporation in electronic retrieval systems.

Authors retain or are hereby granted (without the need to obtain further permission) rights to use the article for traditional scholarship communications, for teaching, and for distribution within their institution, as set out in the General Terms of Publication (see note 1), and also agree to the other terms.

- I am the sole author of the manuscript
- I am one author signing on behalf of all co-authors of the manuscript
- The manuscript is a 'work made for hire' and I am signing as an authorized representative of the employing company
- I am a US Government employee and there is no copyright to transfer but I affirm the author warranties (see notes 3 & 4)
- I am a co-author who is not a US Government employee but whose co-authors are government employees (see note 4)
- I am an employee of the UK
- Canadian or Australian Government claiming Crown Copyright but I affirm the author warranties (see note 5)
- I am a co-author who is not claiming Crown Copyright but whose co-authors are employees of the UK
- Canadian or Australian Government (see note 5)

## FUNDING

- The underlying research reported in the article was funded by the US National Institute of Health**
- The underlying research reported in the article was performed by a Howard Hughes Medical Institute investigator**

Elsevier will automatically submit your paper to PubMed Central if the research was funded by the NIH (as indicated by checking the appropriate box above)

Please mark **one or more** of the above boxes (as appropriate) and then input your name and date.

Type Name: Ajay M Shah

Title and Company (if employer representative):

Date: 6 Dec 2013

Data Protection: By submitting this form you are consenting that the personal information provided herein may be used by the American College of Cardiology Foundation, Elsevier, and its affiliated companies worldwide to contact you concerning the publishing of your article and occasionally for marketing purposes. We respect your privacy. If you **do** not wish to receive news, promotions and special offers about our products and services, then please mark this box. See also the Elsevier website at <http://www.elsevier.com> and click *Privacy Policy*.

Please complete this electronic file by filling out all required criteria and clicking the "Submit" button at the bottom of this page. You will receive a PDF version of your completed electronic form via email. Please print

this copy and retain for your files.

*Journal of the American College of Cardiology (JACC)*  
*JACC: Cardiovascular Interventions*  
*JACC: Cardiovascular Imaging*  
*JACC: Heart Failure* 3655 Nobel Drive, Suite 630  
San Diego, California 92122-1005  
Fax: (858) 558-3148  
Email: [jaccsd@acc.org](mailto:jaccsd@acc.org)  
[jaccint@acc.org](mailto:jaccint@acc.org)  
[jaccimg@acc.org](mailto:jaccimg@acc.org)  
[jacchf@acc.org](mailto:jacchf@acc.org)

## **General Terms of Publication**

### **1. As an author you (or your employer or institution) may do the following:**

- make copies (print or electronic) of the article for your own personal use, including for your own classroom teaching use;
- make copies and distribute such copies (including through e-mail) of the article to research colleagues, for the personal use by such colleagues (but not commercially or systematically, e.g. via an e-mail list or list server);
- post a pre-print version of the article on Internet websites including electronic pre-print servers, and to retain indefinitely such version on such servers or sites;
- post a revised personal version of the final text of the article (to reflect changes made in the peer review and editing process) on your personal or institutional website or server, with a link to the journal homepage (on [elsevier.com](http://elsevier.com));
- present the article at a meeting or conference and to distribute copies of the article to the delegates attending such meeting;
- for your employer, if the article is a 'work for hire', made within the scope of your employment, your employer may use all or part of the information in the article for other intra-company use (e.g. training);
- retain patent and trademark rights and rights to any process or procedure described in the article;
- include the article in full or in part in a thesis or dissertation (provided that this is not to be published commercially);
- use the article or any part thereof in a printed compilation of your works, such as collected writings or lecture notes (subsequent to publication of the article in the journal); and
- prepare other derivative works, to extend the article into book-length form, or to otherwise re-use portions or excerpts in other works, with full acknowledgement of its original publication in the journal.

All copies, print or electronic, or other use of the paper or article must include the appropriate bibliographic citation for the article's publication in the journal.

#### **• Requests from third parties**

Requests for all uses not included above, including the authorization of third parties to reproduce or otherwise use all or part of the article (including figures and tables), should be referred to the Elsevier Global Rights Department by going to our website at <http://www.elsevier.com/locate/permissions> and selecting 'Permissions'.

#### **• Author warranties**

- The article you have submitted to the journal for review is original, has been written by the stated authors and has not been published elsewhere.
- The article is not currently being considered for publication by any other journal and will not be submitted for such review while under review by the journal.

- The article contains no libellous or other unlawful statements and does not contain any materials that violate any personal or proprietary rights of any other person or entity.
- You have obtained written permission from copyright owners for any excerpts from copyrighted works that are included and have credited the sources in your article.
- If the article was prepared jointly with other authors, you have informed the co-author(s) of the terms of this copyright transfer and that you are signing on their behalf as their agent, and represent that you are authorized to do so.

• **US Government employees**

- If all co-authors are US Government employees there is no copyright to transfer. Please sign the form, to confirm the author warranties.
- If there is a number of co-authors, of which at least one is a US Government employee (and this work was prepared in such capacity) and at least one is not a government employee, the non-government author should sign this form, indicating transfer of those rights which such author has (also on behalf of any other non-government co-authors).

• **Crown Copyright**

- UK Government employee authors may elect to transfer copyright.
- UK Government employees wishing to claim Crown Copyright should mark the appropriate box overleaf, sign the form to affirm the author warranties and attach the completed authorization form as per HMSO guidelines at <http://www.hmso.gov.uk/copyright/guidance/articles/htm>.
- The work of Canadian or Australian Government employees is automatically subject to Crown Copyright. Please mark the appropriate box and sign the form to affirm the author warranties.
- If there is a number of co-authors, of which at least one is claiming Crown Copyright and at least one is not an employee of the UK, Canadian or Australian Government, the non-government author should sign this form, indicating transfer of those rights which such author has (also on behalf of any other non-government co-authors).

• **Elsevier's AiP (Articles in Press) service**

Elsevier may choose to publish an abstract or portions of the paper before we publish it in the journal. Please contact our Production department immediately if you do not want us to make any such prior publication for any reason, including disclosure of a patentable invention.

**\*If you have checked a box indicating that you are NOT signing on behalf of all co-authors on the manuscript, the Editorial Office will be in contact to collect the remaining signatures. If you checked the box indicating that you are signing on behalf of all co-authors, please complete this form and submit. Nothing further will be needed.**

I am not signing on behalf of all co-authors of the manuscript.

I am one author signing on behalf of all co-authors of the manuscript.

**Name(printed):** Ajay M Shah

**Title (if employer representative):**

**Date:** 6 Dec, 2013

local\_p\_id: 33472

time: 1386327399

ip address: 94.2.190.208



AMERICAN COLLEGE OF CARDIOLOGY FOUNDATION  
AUTHOR RESPONSIBILITIES: RELATIONSHIP WITH INDUSTRY AND FINANCIAL DISCLOSURE

In addition to disclosing relationship with industry and/or financial disclosures on the *cover letter* and *title* page of your manuscript, we ask that each author sign this form.

**To be published in the following journal (please check):**

*Journal of the American College of Cardiology(JACC)*  *JACC: Cardiovascular Interventions*  *JACC: Cardiovascular Imaging*  *JACC: Heart Failure*

Article entitled: Endothelial NADPH oxidase-2 promotes interstitial cardiac fibrosis and diastolic dysfunction through pro-inflammatory effects and endothelial-mesenchymal transition

Manuscript number: JACC110813-4952D

Corresponding Author: Prof. Shah

Corresponding author's printed name: Ajay M Shah

**\*Corresponding Author's Responsibility**

To avoid delay in the publication of your manuscript, **complete and submit this form as quickly as possible.**

After you have submitted your electronic form, you will receive an email confirmation that includes a PDF copy. Please print and retain for your files. **If for some reason you are unable to complete this form electronically**, please contact the JACC office directly or print this form, fill in the spaces manually, and email, fax, or post mail a copy to:

*Journal of the American College of Cardiology(JACC)*  
*JACC: Cardiovascular Interventions*  
*JACC: Cardiovascular Imaging*  
*JACC: Heart Failure*  
3655 Nobel Drive, Suite 630  
San Diego, California 92122-1005  
Fax: (858) 558-3148  
Email: [jaccsd@acc.org](mailto:jaccsd@acc.org) [jaccint@acc.org](mailto:jaccint@acc.org) [jaccimg@acc.org](mailto:jaccimg@acc.org) [jacchf@acc.org](mailto:jacchf@acc.org)

Complete the section below, type your name into the Type Name box and input the date. *By submitting this form, you attest to being an author on the paper.*

Type Name: Ajay M Shah Date: 6 Dec, 2013

Title and Company (if employer representative):

I have **no relationship with industry** to disclose.

I have **no financial information** to disclose.

local\_p\_id: 33472

time: 1386327443

ip address: 94.2.190.208