

1 Title Page

2 Use of a saliva-based diagnostic test to inform on tapeworm infection in horses in the UK

3 Kirsty L. Lightbody<sup>1</sup>, Jacqueline B. Matthews<sup>2</sup>, Jeremy G. Kemp-Symonds<sup>3</sup>, Peter A. Lambert<sup>4</sup> and  
4 Corrine J. Austin<sup>1</sup>

5 <sup>1</sup>Austin Davis Biologics, Unit 1, Denfield Lodge, Lower St, Great Addington, Northamptonshire, UK,  
6 NN14 4BL

7 <sup>2</sup>Moredun Research Institute, Pentlands Science Park, Edinburgh, Scotland, UK, EH23 0PZ,

8 <sup>3</sup>Bransby Horses, Bransby, Lincoln UK, LN1 2PH

9 <sup>4</sup>School of Health and Life Sciences, Aston University, Birmingham, UK, B4 7ET

10

11 Keywords

12 Anoplocephala; Cestodes; EquiSal<sup>®</sup>; Targeted selective treatment

13

14 Author's declaration of interests

15 The authors Corrine J. Austin and Kirsty L. Lightbody report an affiliation to a commercial entity with  
16 a financial interest in materials discussed in this manuscript.

17

18 Ethical animal research

19 Saliva samples were collected as part of a routine targeted deworming programme.

20

21 Source of funding

22 None

23

24 Acknowledgements

25 The authors would like to thank the staff at Bransby Horses for collecting the saliva samples and  
26 administering anthelmintic treatments, as well as Michelle Billington for help collating the data. Philip  
27 Renshaw and Stuart Trevallion at Austin Davis Biologics performed the EquiSal® Tapeworm tests and  
28 reported results.

29

30 Authorship

31 Kirsty L. Lightbody analysed the data, drafted and revised the manuscript. Jacqueline B. Matthews  
32 critically reviewed and revised the manuscript. Jeremy G. Kemp-Symonds, Bransby Horses Veterinary  
33 Consultant, contributed towards acquisition of the samples, prescribed anthelmintics and critically  
34 reviewed and revised the manuscript. Peter A. Lambert contributed towards data analysis and  
35 critically reviewed and revised the manuscript. Corrine J. Austin analysed the data and critically  
36 reviewed and revised the manuscript.

37

38 Summary

39 Background

40 Anthelmintic resistance combined with limited chemotherapeutic options has prompted a change in  
41 approaches to control of equine helminth infections. Targeted selective treatment (TST) strategies  
42 utilise diagnostics to reduce anthelmintic use by treating individuals with worm burdens or egg  
43 shedding levels above a set threshold. Whilst faecal egg count analysis has limitations for informing

44 tapeworm treatment, a commercially available saliva-based diagnostic test accurately diagnoses  
45 horses with tapeworm infection.

#### 46 Objectives

47 Evaluation of a saliva-based diagnostic test to identify horses naturally-infected with tapeworm and  
48 assess the impact of using the test to inform anthelmintic administration.

#### 49 Study design

50 Retrospective longitudinal study.

#### 51 Methods

52 Saliva was collected from horses (n=237) at a UK welfare charity from autumn 2015-autumn 2016.  
53 Horses diagnosed as positive for tapeworm infection using the EquiSal® Tapeworm test were  
54 anthelmintic treated according to weight. The number of horses that received anthelmintic treatment  
55 based on the test result was compared to an all-group treatment approach and the reduction in  
56 anthelmintic usage calculated. Incoming horses were also tested (n=143) and the information used to  
57 inform quarantine treatments.

#### 58 Results

59 In autumn 2015, 85% of 237 horses tested received no anthelmintic and the majority (71%) of these  
60 remained below the treatment threshold throughout the study. Of the 69 horses that received  
61 treatment, seven required treatment following three subsequent tests, whilst >50% of horses  
62 administered with anthelmintic fell below the treatment threshold at the following test. No increase  
63 in tapeworm prevalence within the 237 horses was observed during the study despite a substantial  
64 reduction in the application of anti-tapeworm treatments. A total of 41% of incoming horses required  
65 anti-cestode treatment.

#### 66 Main Limitations

67 Other management practices were not included in the analysis.

## 68 Conclusions

69 Compared to an all-group treatment strategy, the diagnostic-led approach used here considerably  
70 reduced application of anti-cestode anthelmintics. This could reduce selection pressure for  
71 anthelmintic resistance.

72

## 73 Introduction

74 Horses are exposed to a range of parasitic helminths when grazing. These include tapeworms  
75 (*Anoplocephala perfoliata*, *Anoplocephala magna*, *Paranoplocephala mamilliana*) and roundworms  
76 (cyathostomins, large strongyles, *Parascaris equorum*). Studies in the UK and Ireland have shown *A.*  
77 *perfoliata* to be present in 51-69% of horses examined [1, 2, 3, 4]. These parasites have been  
78 associated with colic, weight loss and colitis [5, 6, 7]. Previously, frequent all-group administration of  
79 anthelmintics was used to control equine helminths; however, interval treatment-based practices  
80 strongly selected for anthelmintic resistance, especially in cyathostomins and *P. equorum* [8, 9, 10]. A  
81 limited number of anthelmintic classes are available for treating equine helminths and with no new  
82 chemical classes in development, care must be taken to preserve efficacy of the currently-effective  
83 products [9, 10]. Targeted selective treatment (TST) strategies aim to reduce use of anthelmintics by  
84 exploiting diagnostics (for example, faecal egg count [FEC] analysis) to identify animals that require  
85 treatment to reduce shedding of worm eggs in faeces. In the UK, this has become relatively  
86 commonplace in worm control programmes [11, 12] and is of value for detecting nematode eggs but  
87 not cestode eggs; the excretion of *A. perfoliata* eggs is intermittent and eggs are unevenly distributed  
88 in faeces resulting in low sensitivity of coprological analysis [13]. To address this, a serum-based test  
89 to diagnose tapeworm was developed and has been commercially available for over a decade [14].  
90 More recently, a non-invasive saliva ELISA test (EquiSal<sup>®</sup> Tapeworm<sup>1</sup>) [4], based on the same antigens,

91 was developed to facilitate uptake of testing by horse owners. The EquiSal® Tapeworm test, validated  
92 by comparing antigen-specific antibody levels in 104 equine saliva samples with tapeworm burdens  
93 post-mortem, differentiated 'low' burdens (0 tapeworm) from 'moderate/high' burdens (>1  
94 tapeworm) with 83% sensitivity and 85% specificity [4]. The pathological effects of tapeworm correlate  
95 with burden intensity and lesions are more severe in horses with >20 tapeworms [2, 15, 16]. In a  
96 previous study, all horses with a clinically-relevant 'high' burden (>20 tapeworms) were accurately  
97 diagnosed using this test [4]. In practice, diagnosis is based on a 'saliva score', with a score of <-0.09  
98 assigned as 'low' and >0.6 assigned as 'moderate/high'. Anti-tapeworm treatment is recommended  
99 when a saliva score = or > -0.09 is obtained. Reported here, is the first study evaluating the utility of  
100 this test in naturally-infected horses and the impact that use of the test has on the number of  
101 praziquantel administrations applied over 11 months.

## 102 Materials and Methods

### 103 Sample population

104 Saliva samples (n=1,000) were collected from horses as part of a site-wide targeted treatment  
105 programme for tapeworm at a UK welfare charity. Samples were predominantly obtained in  
106 October/November 2015 ('autumn 2015', n=305), April/May 2016 ('spring 2016', n=328) and  
107 August/September 2016 ('autumn 2016', n=367) and were collected from horses of both sexes and  
108 various breeds over a wide age range. The horses were grazed in 28 fields and only stabled under  
109 exceptional circumstances. The fields were maintained by a variety of means, including manure  
110 collection in some fields and resting of paddocks. During the study, between autumn 2015 and autumn  
111 2016, a total of 81 horses left the premises and 143 horses were introduced. This included 40 horses  
112 leaving before testing in spring 2016 and a further 41 leaving before testing in autumn 2016. Eighty-  
113 eight new horses arrived between autumn 2015 and spring 2016 and 85 of these were tested in spring  
114 2016. A further 53 new horses arrived between spring 2016 and autumn 2016 and two horses returned  
115 to re-join the population. Saliva samples were collected using EquiSal® saliva collection kits<sup>1</sup> as per

116 manufacturer's instructions. The samples were stabilised in preservative buffer (1x PBS, 0.05% Tween  
117 20, 0.05% 5-bromo-5-nitro-1,3-dioxane (BND) and 0.05% sodium azide) and centrifuged at 3000 x g  
118 for 5 min prior to analysis.

#### 119 Diagnostic testing

120 EquiSal<sup>®</sup> Tapeworm testing was carried out at Austin Davis Biologics<sup>1</sup> as described by Lightbody *et al.*  
121 [4], where each saliva sample was analysed in a test that utilised three ELISAs; the 'Specific' ELISA to  
122 detect IgG(T) specific to excretory/secretory 12/13 kDa tapeworm antigens, the 'Non-Specific Binding'  
123 (NSB) ELISA to determine levels of non-specific binding and the 'Total' ELISA to measure the total  
124 amount of IgG(T) in the sample [4].

#### 125 Anthelmintic administration

126 Horses were treated for *Anoplocephala* infection based on the saliva score derived from the EquiSal<sup>®</sup>  
127 Tapeworm test, above. Horses diagnosed as 'low' (<-0.09) received no anti-cestode treatment and  
128 those diagnosed above the 1+ burden cut-off, indicated by a 'borderline' (-0.09-0.6) or  
129 'moderate/high' (>0.6) saliva score, were administered with praziquantel-based products such as  
130 Equimax<sup>2</sup>, Equest Pramox<sup>3</sup>, Noropraz<sup>4</sup> and EquiTape<sup>5</sup> by attending staff. Horses were weighed using a  
131 weighbridge and were dosed with between 1-2.5 mg of praziquantel/kg as per manufacturers'  
132 instructions on the product used.

#### 133 Analysis

134 Microsoft<sup>®</sup> Office Excel 2013<sup>6</sup> was used to record EquiSal<sup>®</sup> Tapeworm test dates and saliva scores for  
135 each horse. Horses were organised according to the number of tests carried out then sorted according  
136 to test dates. New arrivals, as well as horses that had left the premises, were identified. For the  
137 purposes here, only data from permanent residents between autumn 2015 and autumn 2016,  
138 referred to as the 'constant herd', were analysed using scatter plots in Microsoft<sup>®</sup> Office Excel 2013.  
139 EquiSal<sup>®</sup> Tapeworm saliva scores for individuals were plotted over the testing period and the number

140 of anti-cestode anthelmintic administrations calculated based on the number of borderline or  
141 moderate/high diagnoses.

## 142 Results

143 The constant herd (n=237) comprised horses tested on all three occasions (n=215) and horses tested  
144 in autumn 2015 and autumn 2016, but not spring 2016 (n=22). Within this population were 113  
145 geldings (48%) and 124 mares (52%) (Figure 1, panel A) with ages ranging from 1 year to 37 years with  
146 38 (16%) aged between 1 to 5 years, 63 (27%) aged between 5 to 10 years, 36 (15%) aged between 11  
147 to 15 years, 41 (17%) aged between 16 to 20 years, 38 (16%) aged between 21 to 25 years, 18 (8%)  
148 aged between 26 to 30 years and three (1%) aged over 30 years (Figure 1, panel B). The constant herd  
149 included a wide range of breeds including Thoroughbreds (TBs), Arabs and crosses (n=24), Cobs and  
150 crosses (n=65), Native ponies and crosses (n=139), Warmbloods and crosses (n=5) and Draught horses  
151 and crosses (n=4) (Figure 1, panel C).

152 Between autumn 2015 and autumn 2016, a total of 1,000 saliva samples were tested (Table 1). This  
153 included 305 horses tested between October and December 2015 (autumn 2015), 328 tested between  
154 March and June 2016 (spring 2016) and 367 tested between August and October 2016 (autumn 2016).  
155 Analysis of the test outputs revealed that 71% of the constant herd (168 horses) received no anti-  
156 cestode treatment between autumn 2015 and autumn 2016. Only 69 horses (29% of the constant  
157 herd) were administered a praziquantel-based anthelmintic during this period (Figure 2).

158 Testing in autumn 2015 showed that 202 horses (85%) from the constant herd fell below the  
159 treatment threshold and were diagnosed as having a low burden (below 1+ tapeworm cut-off) (Table  
160 2). Analysis of saliva from the remaining 35 horses (15% of the constant herd) provided values  
161 categorised as borderline (n=17) or moderate/high (n=18) (above 1+ tapeworm cut-off). All of these  
162 were administered a praziquantel-based anthelmintic (Table 2).

163 When re-tested in spring 2016, 184 horses (86%) of the 215 constant herd horses sampled were  
164 diagnosed as low and did not require treatment (Table 2). Thirty-one horses (14%) were administered  
165 a praziquantel-based anthelmintic as they fell in the borderline (n=13) or moderate/high (n=18)  
166 categories (Table 2). Of the 31 horses identified as having scores above the treatment threshold, 15  
167 had previously been below the treatment threshold (Figure 3, Panel A). The remaining 16 horses (7%  
168 of the constant herd) had previously been treated in autumn 2015, following borderline (n=4) or  
169 moderate/high (n=12) test results. The results obtained in spring 2016 indicated that 17 (52%) of 33  
170 constant herd horses that received anti-cestode treatment in autumn 2015, fell below the treatment  
171 threshold and did not require further treatment (Figure 3, Panel B). Of the sixteen horses diagnosed  
172 above the treatment threshold in autumn 2015 and spring 2016, 12 had a lower saliva score in spring  
173 2016 following anthelmintic treatment in autumn 2015 (Figure 3, Panel C). Only four horses had a  
174 higher saliva score in spring 2016 compared to autumn 2015 (Figure 3, Panel C).

175 In autumn 2016, 204 (86%) of the 237 constant herd horses were diagnosed as low (Table 2). Thirty-  
176 three horses (14% of the constant herd) had scores above the treatment threshold and were  
177 diagnosed as borderline (n=16) or moderate/high (n=17) (Table 2). Nineteen horses requiring anti-  
178 cestode treatment in autumn 2016 had previously been diagnosed as low in autumn 2015 and spring  
179 2016 and had received no prior treatment, and two horses had been treated in autumn 2015, but  
180 were below the treatment threshold in spring 2016 (Figure 4, Panel A). Of the constant herd horses  
181 that were administered a praziquantel-based anthelmintic in spring 2016 (n=31), 19 horses (61%) were  
182 diagnosed as low and did not require further treatment in autumn 2016 (Figure 4, Panel B). Twelve  
183 horses were diagnosed as borderline or moderate/high in autumn 2016 following treatment in spring  
184 2016 and included seven horses with lower saliva scores compared to the previous test and five with  
185 increased saliva scores (Figure 4, Panel C).

186 Overall, 168 horses (71% of the constant herd) remained below the treatment threshold at all three  
187 test time points. Thirty-six horses (15% of the constant herd) were diagnosed as low in autumn 2016,



188 following a borderline or moderate/high result in autumn 2015 and/or spring 2016 (Figure 5, Panel A).  
189 The 33 horses (14% of the constant herd) with saliva scores above the treatment threshold in autumn  
190 2016 included nineteen horses diagnosed as low in autumn 2015 and spring 2016 (Figure 5, Panel B)  
191 and 14 horses diagnosed as borderline or moderate/high in autumn 2015 and/or spring 2016 (Figure  
192 5, Panel C). In total, only seven horses (3% of the constant herd) were diagnosed as borderline or  
193 moderate/high in all three tests (Figure 5, Panel D). This included four horses categorised as  
194 moderate/high in all tests, two categorised as moderate/high in autumn 2015 and spring 2016 and  
195 borderline in autumn 2016 and one categorised as moderate/high in autumn 2015 and borderline in  
196 spring and autumn 2016.

197 The gender, age and breed of horses in the constant herd and the number of borderline or  
198 moderate/high diagnoses obtained between autumn 2015 to autumn 2016 are shown in Table 3.  
199 Geldings and mares reported a similar proportion of borderline or moderate/high diagnoses (32/113  
200 vs. 37/124), however younger horses required more treatments with 21/38 (55%) of 1-5 year old  
201 horses being diagnosed as borderline or moderate/high on at least one occasion. In addition, 10/38  
202 (26%) of the 1-5 year old horses required multiple treatments following borderline or moderate/high  
203 diagnoses in two or three tests. Cobs and Cob crosses received the most treatments, with 32/65 (49%)  
204 reporting at least one borderline or moderate/high result.

205 Over the study, 141 new horses arrived and two horses returned to the site. Analysis of the test results,  
206 indicated that 35 (40%) of the 88 horses that arrived from autumn 2015 to spring 2016 and 23 (42%)  
207 of the 55 horses that arrived between spring 2016 and autumn 2016 were diagnosed in the borderline  
208 or moderate/high categories.

## 209 Discussion

210 As demonstrated for strongyle FEC-based TST-strategies [8, 11], the results here show that use of a  
211 diagnostic test to inform on the requirement to treat *Anoplocephala* infection, led to lower  
212 anthelmintic usage compared with an interval treatment strategy in which all horses were

213 administered with an anti-cestode product in the spring and autumn. In total, 99 doses of praziquantel  
214 product were administered to the constant herd using this approach. This represents an 86%  
215 reduction in anthelmintic administration during the study period compared to an interval treatment  
216 strategy based on two annual treatments for all horses. Despite the reduction in anthelmintic use,  
217 tapeworm infection prevalence did not increase during the course of the study.

218 The majority of horses from the constant herd, diagnosed as being below the treatment threshold in  
219 autumn 2015, fell below this level and were diagnosed as low in subsequent tests. The apparent over-  
220 dispersed distribution of infection here is similar to previous reports on tapeworm infection intensity  
221 [18, 19, 20], as well as for other types of equine helminths [21, 22]. The fact that 168 horses were  
222 diagnosed as low on all three occasions suggests that some horses control *Anoplocephala* burden  
223 level, similar to results reported for nematode infections, where some horses repeatedly have low FEC  
224 values, even in the absence of anthelmintic treatment [17, 23, 24], or they were not exposed to  
225 infection.

226 Praziquantel is rapidly absorbed following oral administration and the drug and its metabolites are  
227 predominantly excreted within 24 hr [25]. Despite the lack of persistence of the drug, most horses  
228 that received praziquantel treatment in autumn 2015 or spring 2016, showed lower saliva scores in  
229 the subsequent test; with 54% and 61% of the horses treated in autumn 2015 and spring 2016 falling  
230 below the treatment threshold at the following test. This indicates that treatments were likely to have  
231 been effective or to have lowered the burden sufficiently to reduce the stimulation of antigen-specific  
232 IgG(T). The remaining 46% of horses that received treatment in autumn 2015 tested above the  
233 treatment threshold again in spring 2016 and 39% of horses treated in spring 2016, tested above the  
234 treatment threshold again in autumn 2016. It is possible that these horses had become re-infected as  
235 a previous study [4] reported that saliva scores of eight horses fell below the treatment threshold  
236 within five weeks of treatment and the scores of three remaining horses fell below the treatment  
237 threshold within three months, indicating that, in a re-test at six months, saliva scores above the

238 threshold are suggestive of the acquisition of new infection. Notably, a substantial increase in  
239 tapeworm incidence was not observed in those horses diagnosed as low burden, as only 7% of  
240 untreated horses in autumn 2015 required treatment following testing in spring 2016 and similarly,  
241 only 11% of untreated horses in spring 2016 required treatment following testing in autumn 2016.

242 The patterns of infection and reinfection here highlight the value of biannual monitoring as the  
243 prevalence and frequency of tapeworm infections can be affected by factors such as grazing practice,  
244 pasture management, the presence of oribatid mite intermediate hosts, as well as climatic and  
245 environmental conditions [19, 26, 27, 28]. Prevalence and distribution of tapeworm infection will vary  
246 between seasons and between geographical locations; therefore regular biannual monitoring of  
247 individuals would identify those acquiring new infections allowing treatment at an early stage,  
248 reducing contamination of the pasture and limiting exposure of the rest of the herd, as well as  
249 preventing unnecessary use of anthelmintics. Regular monitoring will also identify those individuals  
250 which may be more prone to re-infection.

251 Over the study period, only seven horses (9% of the constant herd tested) received three doses of  
252 praziquantel-based anthelmintic due to borderline or moderate/high scores in all tests. This group  
253 included four Cob geldings, one Thoroughbred gelding, one Cob mare and a Native pony mare. In line  
254 with previous reports, no association between gender and tapeworm infection was observed in this  
255 study [19, 26]. Studies have also reported that tapeworm infection prevalence and intensity is not  
256 influenced by age or breed [2, 16, 18, 26]. Younger horses in this population (between 1-5 years)  
257 reported more borderline or moderate/high scores than older horses (>6 years) and although this may  
258 be suggestive of age-related exposure to infection or the development of acquired immunity [29], and  
259 is similar to the influence of age on strongyle infections [22, 30], additional research would be required  
260 to investigate and confirm age-related exposure as determined by saliva testing. The high number of  
261 Cobs and Cob crosses reporting borderline or moderate/high results throughout this study may be  
262 related to age rather than breed, as 40% of these horses were in the 1-5 year age group.

263 New horses joining the population may be responsible for pasture contamination. Here, 41% of new  
264 arrivals were diagnosed with a burden indicative of a requirement to treat. This proportion of test-  
265 positive horses is higher than that of the constant herd (approximately 15%) and may be reflective of  
266 horses of arriving from sites with unknown or questionable helminth control practices. This augments  
267 the need to enact appropriate quarantine procedures, either by testing or deworming, on all incoming  
268 individuals into populations.

269 The EquiSal<sup>®</sup> Tapeworm test [4] reports sensitivity and specificity of 83% and 85%, respectively, when  
270 a 1+ tapeworm cut-off is applied (low burden = 0 tapeworm, moderate/high burden = 1+ tapeworm).  
271 In comparison, coprological diagnosis of *Anoplocephala* infection is highly variable, in part due to the  
272 sporadic release of tapeworm eggs, poor distribution of eggs in faeces, the FEC technique used and  
273 the burden present [13, 20]. Modified FEC methods, such as the centrifugation/flotation technique,  
274 are most sensitive, reporting up to 61% sensitivity, however they are more time consuming and labour  
275 intensive than standard methods [7, 13, 18, 20, 31]. Sensitivity of the centrifugation/flotation  
276 technique can be increased to approximately 90% when diagnosing infections with >20 tapeworms  
277 present [18, 31]. When a 20+ tapeworm cut-off is applied to the EquiSal<sup>®</sup> Tapeworm test  
278 (low/moderate burden = 0-19 tapeworms, high burden = 20+ tapeworm), sensitivity is 86% [4].  
279 Overall, the EquiSal<sup>®</sup> Tapeworm test shows similar sensitivity to modified FEC analysis when  
280 diagnosing high (20+) tapeworm burdens however, the saliva-based test displays higher sensitivity  
281 when diagnosing low (1+) burdens, which may include infections with immature tapeworms that  
282 would not be identified using FEC methodologies [18, 20, 28]. If saliva testing is undertaken twice a  
283 year and/or at least four months after anti-cestode treatment, then it is likely that antibodies, if  
284 detected, are due a current infection rather than a historic infection as it was reported that antigen-  
285 specific salivary antibodies of 11 horses reduced below the 1+ tapeworm cut-off within three months  
286 of treatment [4]. However, the dynamics of tapeworm-specific antibodies in the saliva will depend on  
287 the individual and whether reinfection occurs.

288 The use of diagnostic tests to predict *Anoplocephala* burden and hence inform anti-cestode  
289 treatments will reduce the frequency of chemicals, such as praziquantel and pyrantel, used in practice.  
290 This theoretically could reduce selection pressure for the development of drug resistance not just in  
291 cestodes but in nematodes too, as some products contain a combination of praziquantel and  
292 macrocyclic lactones. Resistance to macrocyclic lactones in nematodes is already a concern as  
293 shortened egg reappearance periods (ERP) following ivermectin and moxidectin use have been  
294 reported [10, 32, 33]. Additionally, resistance to pyrantel, a broad spectrum anthelmintic used to treat  
295 tapeworm infections, has also been reported in nematodes [8, 32, 34]. Although anthelmintic  
296 resistance has yet to be documented for *A. perfoliata* and other tapeworm species, it is not  
297 unforeseeable that, with the widespread application of regular blanket treatments, resistance would  
298 be inevitable. This study demonstrates that the saliva-based EquiSal® Tapeworm test holds promise  
299 in reducing treatment frequency in practice, which could help protect efficacy of anti-cestode  
300 anthelmintics for the future.

301 Manufacturer's details

302 1 Austin Davis Biologics Ltd., Great Addington, Northamptonshire, UK

303 2 Virbac, Carros, France

304 3 Zoetis, London, UK

305 4 Norbrook Laboratories Ltd, Newry, UK

306 5 Bayer, Newbury, UK

307 6 Microsoft, Redmond, WA, USA

308 Figure Legends

309 **Figure 1.** Demographics of the constant herd horses (n=237), including gender (A), age (B) and breed  
310 (C).

311 **Figure 2.** Anti-cestode treatments administered to constant herd horses (n=237) between autumn  
312 2015 and autumn 2016. Ninety-nine doses of anthelmintic were administered to 69 horses, with 46  
313 treated once, 16 treated twice and seven treated three times.

314 **Figure 3.** EquiSal® Tapeworm test saliva scores for constant herd horses tested in autumn 2015 and  
315 spring 2016. (A) Untreated horses that changed from a low to a borderline or moderate/high (n=15).  
316 (B) Horses treated in autumn 2015 that changed from a borderline or moderate/high result to a low  
317 result (n=17). (C) Horses treated in autumn 2015 that remained in either borderline or moderate/high  
318 result categories (n=16).

319 **Figure 4.** EquiSal® Tapeworm test saliva scores for constant herd horses tested in autumn 2015, spring  
320 2016 and autumn 2016. (A) Untreated horses from spring 2016 that changed from low to a borderline  
321 or moderate/high result (n=21). (B) Horses treated in spring 2016 that changed from a borderline or  
322 moderate/high result to a low result (n=19). (C) Horses treated in spring 2016 that remained either in  
323 borderline or moderate/high result categories (n=12).

324 **Figure 5.** EquiSal® Tapeworm test saliva scores for constant herd horses tested in autumn 2015, spring  
325 2016 and autumn 2016. (A) Horses treated in either autumn 2015 and/or spring 2016 that changed  
326 from a borderline or moderate/high result to a low result in autumn 2016 (n=36). (B) Untreated horses  
327 from autumn 2015 and spring 2016 that changed from a low result to a borderline or moderate/high  
328 result in autumn 2016 (n=19) (C) Horses treated in autumn 2015 and/or spring 2016 that remained in  
329 either borderline or moderate/high results categories in autumn 2016 (n=14). (D) Horses treated in  
330 both autumn 2015 and spring 2016 that remained in either borderline or moderate/high results  
331 categories in autumn 2016 (n=7).

### 332 References

333 [1] Owen, R.A, Jagger, D.W. and Quan-Taylor R. (1988) Prevalence of *Anoplocephala perfoliata* in  
334 horses and ponies in Clwyd, Powys and adjacent English marches. *Vet Rec* **123**, 562-563

335 [2] Fogarty, U., del Piero, F., Purnell, R.E. and Mosurski, K.R. (1994) Incidence of Anoplocephala  
336 perfoliata in horses examined at an Irish abattoir. *Vet Rec* **134**, 515-518

337 [3] Pittaway, C.E., Lawson, A.L., Coles, G.C. and Wilson A.D. (2014) Systemic and mucosal IgE antibody  
338 responses of horses to infection with Anoplocephala perfoliata. *Vet Parasitol* **199**, 32-41

339 [4] Lightbody, K.L, Davis, P.J. and Austin, C.J. (2016) Validation of a novel saliva-based ELISA test for  
340 diagnosing tapeworm burden in horses. *Vet Clin Path* **45**, 335-346

341 [5] Cribb, N.C., Cote, N.M., Boure, L.P. and Peregrine A.S. (2006) Acute small intestinal obstruction  
342 associated with Parascaris equorum infection in young horses: 25 cases (1985-2004). *New Zeal Vet J*  
343 **54**, 338-343

344 [6] Corning, S. (2009) Equine cyathostomins: a review of biology, clinical significance and therapy.  
345 *Parasite Vector* **2**, S1

346 [7] Nielsen, M.K. (2016) Equine tapeworm infection: Disease, diagnosis and control. *Equine Vet Educ*  
347 **28**, 388-395

348 [8] Kaplan, R.M. (2002) Anthelmintic resistance in nematodes of horses. **33**, 491-507

349 [9] Nielsen, M.K. (2012) Sustainable equine parasite control: Perspectives and research needs. *Vet*  
350 *Parasitol* **185**, 32-44

351 [10] Matthews, J.B. (2014) Anthelmintic resistance in equine nematodes. *Int J Parasitol* **4**, 310-315

352 [11] Stratford, C.H., Lester, H.E., Morgan, E.R., Pickles, K.J., Relf, V., McGorum, B.C. and Matthews, J.B.  
353 (2014) A questionnaire study of equine gastrointestinal parasite control in Scotland. *Equine Vet J* **46**,  
354 25-31

355 [12] Easton, S., Pinchbeck, G.L., Tzelos, T., Bartley, D.J., Hotchkiss, E., Hodgkinson, J.E. and Matthews,  
356 J.B. (2016) Investigating interactions between UK horse owners and prescribers of anthelmintics. *Prev*  
357 *Vet Med* **135**, 17-27

- 358 [13] Abbott, J.B. and Barrett, E.J. (2008) The problem of diagnosing tapeworm infections in horses.  
359 *Equine Vet J* **40**, 5-6
- 360 [14] Proudman, C.J. and Trees, A.J. (1996) Correlation of antigen specific IgG and IgG(T) responses with  
361 *Anoplocephala perfoliata* infection intensity in the horse. *Parasite Immunol* **18**, 499-506
- 362 [15] Gasser, R.B., Williamson, R.M. and Beveridge I. (2005) *Anoplocephala perfoliata* of horses -  
363 significant scope for further research, improved diagnosis and control. *Parasitology* **131**, 1-13
- 364 [16] Edwards, G.B. (1999) The role of tapeworms in equine colic. *Pferdeheilkunde* **15**, 309-312
- 365 [17] Nielsen, M.K., Pfister, K. and von Samson-Himmelstjerna, G. (2014) Selective therapy in equine  
366 parasite control – Application and limitations. *Vet parasitol* **202**, 95-103
- 367 [18] Kjaer, L.N., Lungholt, M.M., Nielsen, M.K., Olsen, S.N. and Maddox-Hyttel, C. (2007) Interpretation  
368 of serum antibody response to *Anoplocephala perfoliata* in relation to parasite burden and faecal egg  
369 count. *Equine Vet J* **39**, 529-533
- 370 [19] Rehbein, S., Visser, M. and Winter, R. (2013) Prevalence, intensity and seasonality of  
371 gastrointestinal parasites in abattoir horses in Germany. *Parasitol Res* **112**, 407-413
- 372 [20] Tomczuk, K., Kostro, K., Szczepaniak, K.O., Grzybek, M., Studzinska, M., Demkowska-Kutrzepa, M.  
373 and Roczen-Karczmarz, M. (2014) Comparison of the sensitivity of coprological methods in detecting  
374 *Anoplocephala perfoliata* invasions. *Parasitol Res* **113**, 2401-2406
- 375 [21] Lester, H.E., Spanton, J., Stratford, C.H., Bartley, D.J., Morgan, E.R., Hodgkinson, J.E., Coumbe, K.,  
376 Mair, T., Swan, B., Lemon, G., Cookson, R. and Matthews J.B. (2013) Anthelmintic efficacy against  
377 cyathostomins in horses in Southern England. *Vet Parasitol* **197**, 189-196
- 378 [22] Relf, V.E., Morgan, E.R., Hodgkinson, J.E. and Matthews, J.B. (2013) Helminth egg excretion with  
379 regard to age, gender and management practices on UK Thoroughbred studs. *Parasitology* **140**, 641-  
380 652



- 381 [23] Gomez, H.H. and Georgi, J.R. (1991) Equine helminth infections: control by selective  
382 chemotherapy. *Equine Vet J* **23**, 198-200
- 383 [24] Nielsen, M.K., Haaning N. and Olsen S.N. (2006) Strongyle egg shedding consistency in horses on  
384 farms using selective therapy in Denmark. *Vet Parasitol* **135**, 333-335
- 385 [25] Steiner, K., Garbe, A., Diekmann, H.W. and Nowak, H. (1976) The fate of praziquantel in the  
386 organism I. Pharmacokinetics in animals. *Eur J Drug Metab Ph* **1**, 85-95
- 387 [26] Ihler, C.F., Rootwelt, V., Heyeraas, A. and Dolvik, N.I. (1995) The prevalence and epidemiology of  
388 *Anoplocephala perfoliata* infection in Norway. *Vet Res Commun* **19**, 487-494
- 389 [27] Meana, A., Pato, N.F., Martin, R., Mateos, A., Perez-Garcia, J. and Luzon, M. (2005)  
390 Epidemiological studies on equine cestodes in central Spain: infection pattern and population  
391 dynamics. *Vet Parasitol* **130**, 233-240
- 392 [28] Tomczuk, K., Kostro, K., Grzybek, M., Szczepaniak, K., Studzinska, M., Demkowska-Kutrzepa, M.  
393 and Roczen-Karczmarz, M. (2015) Seasonal changes of diagnostic potential in the detection of  
394 *Anoplocephala perfoliata* equine infections in the climate of Central Europe. *Parasitol Res* **114**, 767-  
395 772
- 396 [29] Proudman, C.J., Holmes, M.A., Sheoran, A.S., Edwards, S.E.R. and Trees, A.J. (1997)  
397 Immunoepidemiology of the equine tapeworm *Anoplocephala perfoliata*: age-intensity profile and  
398 age-dependancy of antibody subtype responses. *Parasitology* **114**, 89-94
- 399 [30] von Samson-Himmelstjerna, G., Traversa, D., Demeler, J., Rohn, K., Milillo, P., Schurmann, S., Lia,  
400 R., Perrucci, S., di Regalbono, A.F., Beraldo, P., Barnes, H., Cobb, R. and Boeckh, A. Effects of worm  
401 control practices examined by a combined faecal egg count and questionnaire survey on horse farms  
402 in Germany, Italy and the UK. *Parasite Vector*, **2**, S3

403 [31] Proudman, C.J. and Edwards, G.B. (1992) Validation of a centrifugation/flotation technique for  
404 the diagnosis of equine cestodiasis. *Vet Rec* **131**, 71-72

405 [32] Relf, V.E., Lester, H.E., Morgan, E.R., Hodgkinson, J.E. and Matthews, J.B. (2014) Anthelmintic  
406 efficacy of UK Thoroughbred stud farms. *Int J Parasitol* **44**, 507-514

407 [33] Daniels, S.P. and Proudman, C.J. (2016) Shortened egg reappearance after ivermectin or  
408 moxidectin use in horses in the UK. *Vet J* **218**, 36-39

409 [34] Kaplan, R.M., Klei, T.R., Lyons, E.T., Lester, G., Courtney, C.H., French, D.D., Tolliver, S.C.,  
410 Vidyashankar, A.N. and Zhao, Y. (2004) Prevalence of anthelmintic resistant cyathostomes on horse  
411 farms. *J Am Vet Med Assoc* **225**, 903-910

412

413

414

415

416

417

418

419

420

421

422

423 Tables

424 **Table 1.** Summary of numbers of horses analysed using the EquiSal® Tapeworm test between autumn  
425 2015 and autumn 2016.

	Autumn 2015 horse numbers	Spring 2016 horse numbers	Autumn 2016 horse numbers
Constant herd horses tested three times	215	215	215
Constant herd horses tested twice	22	0	22
Non-constant herd horses tested	68	113	130
Horses arrived	-	88	53
Horses returned	-	0	2
Horses departed	-	40	41

426

427

428

429

430

431

432

433

434

435

436 **Table 2.** EquiSal® Tapeworm test diagnosis and treatment recommendations. Below treatment  
 437 threshold = saliva score <-0.09 ('low'); above treatment threshold = saliva score =>-0.09 ('borderline'  
 438 or 'moderate/high').

	Autumn 2015		Spring 2016		Autumn 2016		
EquiSal® Tapeworm test Diagnosis	Constant herd horses (%)	Constant herd horses (%)	Constant herd horses (%)	Constant herd horses (%)	Constant herd horses (%)	Constant herd horses (%)	Treated for tapeworm
Low	202	85	184	86	204	86	No
Borderline + moderate/high	35	15	31	14	33	14	Yes
Total	237		215		237		

439

440

441

442

443

444

445

446

447

448

449 **Table 3.** Summary of the gender, age and breed of horses that remained low in all three EquiSal® tests  
 450 between autumn 2016 and spring 2016 (n=168) and horses that were diagnosed as borderline (B) or  
 451 moderate/high (M/H) in 1 test (n=46), 2 tests (n=16) and 3 tests (n=7).

	Category	B or M/H in 0 tests	B or M/H in 1 test	B or M/H in 2 tests	B or M/H in 3 tests
Gender	Gelding	81 (72%)	21 (19%)	6 (5%)	5 (4%)
	Mare	87 (70%)	25 (20%)	10 (8%)	2 (2%)
Age	1 to 5 years	17 (45%)	11 (29%)	6 (16%)	4 (10%)
	6 to 10 years	44 (70%)	14 (22%)	3 (5%)	2 (3%)
	11 to 15 years	31 (86%)	4 (11%)	1 (3%)	0 (0%)
	16 to 20 years	30 (73%)	9 (22%)	2 (5%)	0 (0%)
	21 to 25 years	28 (74%)	6 (16%)	4 (10%)	0 (0%)
	26 to 30 years	15 (83%)	2 (11%)	0 (0%)	1 (6%)
	30+ years	3 (100%)	0 (0%)	0 (0%)	0 (0%)
Breed	Thoroughbreds, Arabs and crosses	18 (75%)	5 (21%)	0 (0%)	1 (4%)
	Cobs and crosses	33 (51%)	18 (28%)	9 (14%)	5 (7%)
	Native ponies and crosses	110 (79%)	22 (16%)	6 (4%)	1 (1%)
	Warmbloods and crosses	3 (60%)	1 (20%)	1 (20%)	0 (0%)
	Draught horses and crosses	4 (100%)	0 (0%)	0 (0%)	0 (0%)

452




453


454

455

456

457





459

460

461

462

463

464

465




466

467




468





469

470









473