

Breastfeeding Duration and its Relation to Weight Gain, Eating Behaviours and Positive Maternal Feeding Practices in Infancy

Samantha L Rogers¹, PhD; Jackie Blissett, PhD²

Affiliations: ¹Centre for Research in Primary and Community Care, University of Hertfordshire, College Lane, Hatfield, AL10 9AB, UK.

²Centre for Technology Enabled Health Research, Faculty of Health and Life Sciences, Richard Crossman Building, Coventry University, Priory Street, Coventry, CV1 5FB, UK.

Email addresses: s.rogers7@herts.ac.uk; jacqueline.blissett@coventry.ac.uk

Address correspondence to: Dr Samantha Rogers, Centre for Research in Primary and Community Care, University of Hertfordshire, College Lane, Hatfield, AL10 9AB, UK, +44(0)1707 285215, s.rogers7@herts.ac.uk

Funding Source: This study was supported by the Economic and Social Research Council Studentship Award ES/G017786/1

Financial Disclosure: The authors have no financial relationships relevant to this article to disclose.

Conflict of Interest: The authors have no conflicts of interest to disclose.

Abstract

Research examining the relationship between breastfeeding and infant weight has generated conflicting results. Few studies account for significant covariates and many suffer methodological problems such as retrospective self-report. The current study aimed to investigate relationships between breastfeeding duration, infant weight and eating and positive maternal mealtime behaviours, whilst overcoming many of the limitations of previous research. Eighty-one women on low-risk maternity units gave informed consent and were visited at home at 1-week, 1-, 6- and 12-months postpartum. Infants included 45 males and 36 females (mean birth-weight 3.52 kg [SD 0.39]). Mothers and infants were weighed and measured and feeding information was recorded at each visit. Infant weight was converted to a standard deviation score (SDS^{*}), accounting for age and sex. Mothers reported infant eating behaviours at 12-months using the Children's Eating Behaviour Questionnaire and were observed feeding their infants solid food at home at 6- and 12-months. Partial correlations (covariates: maternal age, education, BMI, smoking during pregnancy, household income, infant birth weight SDS and age introduced to solid foods) revealed negative associations between breastfeeding duration and 1- to 6- and 1- to 12-month weight gain, and 6- and 12-month weight. Breastfeeding duration was also associated with a slower rate of infant eating and greater observed maternal vocalisations, appropriateness and sensitivity. Results support a dose-response relationship between breastfeeding and infant weight and suggest that breastfeeding may encourage the development of obesity-protective eating behaviours through learning to attend to internal hunger and satiety signals. Future research should investigate whether relationships between slowness in eating and weight extend to satiety responsiveness after infancy.

Key words: Breastfeeding duration; feeding practices; eating behaviours; weight gain; infancy

* **Abbreviations:** BMI – body mass index; CEBQ – Child Eating Behaviour Questionnaire; EPDS – Edinburgh Postnatal Depression Scale; SDS – standard deviation score

Introduction

For the last 30 years, research has investigated whether breastfeeding protects against rapid weight gain, overweight and obesity. Findings have revealed that breastfed infants gain less weight during the neonatal period than formula-fed infants (Heinig, Nommsen, Peerson, Lonnerdal, & Dewey, 1993), and that infants who gain less weight during this period have a reduced risk of becoming obese later in life (Stettler, Zemel, Kumanyika, & Stallings, 2002). Rapid weight gain in infancy is a risk factor for overweight/obesity in childhood and is associated with increased BMI and fat mass at 5- and 7- to 9-years (Sacco, de Castro, Euclides, Souza, & Rondo, 2013; Zhou et al., 2016). Although studies have also provided evidence for a dose-dependent protective effect of breastfeeding (Arenz, Ruckerl, Koletzko, & Von Kries, 2004; Hornell, Lagstrom, Lande, & Thorsdottir, 2013; Kramer, 1981; McCrory & Layte, 2012; Owen, Martin, Whincup, Smith, & Cook, 2005; Reynolds, Hennessy, & Polek, 2014; Skledar & Milosevic, 2015; Yan, Liu, Zhu, Huang, & Wang, 2014), there is considerable inconsistency in published findings.

Some studies claim the effect of breastfeeding on childhood obesity is small (Jiang & Foster, 2013; Umer et al., 2015) and others have found no effect at all (Davis et al., 2007; Martin et al., 2013; Novaes, Lamounier, Colosimo, Franceschini, & Priore, 2012; Oddy et al., 2004). It is possible that these null findings may have arisen due to a lack of control of covariates and a range of other methodological issues. There is wide variation between studies in the covariates accounted for; examples include infant birth weight, gender, gestational age, age introduced to solid food, maternal age, BMI, smoking status during pregnancy (and postnatally), maternal diabetes, postnatal depression, education and household income. Very few published studies account for all of these variables. Smithers, Kramer, and Lynch (2015) argue that poor

measurement (or lack of adjustment) of such factors can result in biased effects of breastfeeding being reported from longitudinal cohort studies. The current study attempted to address such issues by measuring the most common covariates not controlled for in other studies. Extensive demographic information was obtained and, if related to breastfeeding or infant weight, controlled for in subsequent analyses.

Methodological problems in this area involve (but are not limited to) retrospective data collection, inconsistent definitions of breastfeeding (including exclusivity and duration), small sample sizes or the same data from larger samples being used several times, and maternal self-report of infant/child height and weight and breastfeeding history. Michels et al. (2007), who did not find an association between breastfeeding and overweight, obtained their sample from the Nurses' Health Study II (whose children comprise the Growing Up Today Study). The same environmental and genetic information has therefore contributed to more than one sample and has been studied numerous times (Gillman et al., 2006; Gillman et al., 2001). Repeated use of the same cohort partially explains the occurrence of repeated findings both for and against the protective effect of breastfeeding on obesity. Furthermore, Michels et al. (2007) administered questionnaires to nurses' mothers asking if they breastfed their daughters and when breastfeeding stopped. Nurses' mothers were contacted when the nurses were aged between 37- and 44-years. The time elapsed since breastfeeding cessation suggests it is likely that mothers could not accurately recollect how they fed their infants and calls into question the accuracy of retrospective self-report. The current study attempted to improve such methodological issues by recruiting a new sample of participants and by avoiding the use of retrospective self-report of information.

Despite inconsistencies within the literature, the protective effect of breastfeeding is often demonstrated in large, methodologically rigorous studies. For example, a large, well-controlled study using multivariate analysis included 7,798 children in Ireland and controlled for socio-demographic factors, child birth weight, gender, physical activity and parental BMI. Results demonstrated that children who had been breastfed for 13- to 25-weeks had a 38% reduction in the risk of being obese at 9-years-of-age, compared to those never breastfed (McCrary & Layte, 2012). Furthermore, breastfeeding for at least 26-weeks was associated with a 51% reduction in obesity risk at 9-years-of-age. These results also supported the dose-dependent effect of breastfeeding for durations greater than 4-weeks (McCrary & Layte, 2012). Furthermore, the protective effect of breastfeeding is also illustrated by a meta-analysis, which found that a longer duration of breastfeeding was associated with a reduced risk of becoming overweight (Harder, Bergmann, Kallischnigg, & Plagemann, 2005). However, as with individual studies, systematic reviews and meta-analyses may also suffer limitations with respect to the potential bias due to confounding (Smithers et al., 2015).

In addition to the effect of breastfeeding on weight, it is also related to the development of healthy eating behaviours, such as increased consumption of fruits and vegetables (Kudlová & Schneidrová, 2012; Mennella, Jagnow, & Beauchamp, 2001). Breastfed infants are also found to be more responsive to satiety (Brown & Lee, 2012) and greater satiety responsiveness is related to a lower risk of being overweight in childhood (Webber, Hill, Saxton, Van Jaarsveld, & Wardle, 2009). Increased responsiveness to satiety may arise because breastfed infants may learn to better self-regulate their intake than formula-fed infants due to having more control over the size of the feed (Birch & Fisher, 1998) and the ever-changing fat content of the milk (Jenness, 1979; Nommsen, Lovelady, Heinig, Lönnerdal, & Dewey, 1991).

Maternal sensitivity is associated with breastfeeding, infant weight gain and eating behaviours and is a potential candidate to explain the mechanism of the protective effects of breastfeeding on obesity. Breastfeeding mothers may be more sensitive and responsive to the hunger and satiety signals communicated by their infant and demonstrate less controlling feeding practices than formula-feeding mothers. Shloim, Rudolf, Feltbower, Mohebati, and Hetherington (2015) observed mealtime interactions between mothers and infants and found that breastfeeding mothers were more in tune with their infants' signals during feeding. Breastfeeding mothers also provided a more favourable feeding environment and fed their infants more responsively than mothers who fed solids or milk from a bottle (Shloim et al., 2015). More sensitive and less controlling behaviours during feeding allow infants to self-regulate their energy intake and learn to respond to internal hunger and satiety cues (Brown & Lee, 2012; Taveras et al., 2006). However, much of the literature to date has relied on maternal report of feeding practices, with few prospective studies of breastfeeding outcomes examining observed sensitivity in solid feeding interactions.

A recent systematic review by Bergmeier, Skouteris, and Hetherington (2015) argued that much of the literature that has investigated relationships between maternal feeding practices and children's weight and eating behaviours has relied on unidirectional self-report methods. It is possible that such methods alone may be biased and capture intended, rather than actual, feeding behaviours (Bergmeier, Skouteris, & Hetherington, 2015). In support of this, Bergmeier, Skouteris, Haycraft, Haines, and Hooley (2015) found that maternal reported restriction was negatively associated with observed restriction during a mealtime observation, and reported pressure was only positively associated with observed pressure in mothers of girls, not boys.

Bergmeier, Skouteris, and Hetherington (2015) argued that longitudinal observational methods should be employed that examine the bi-directional dimensions of parent-child mealtime interactions.

Results of previous literature emphasise the importance of investigating the relationships between breastfeeding duration, infant weight gain and eating behaviours and observed maternal feeding behaviours in one study. Currently, there is no longitudinal study published that investigates all of these factors together over the first year of life. The aim of this study was to investigate the relationship between observed maternal feeding behaviour, breastfeeding duration and infant weight and eating behaviours during the first 12-months of life, in a sample of healthy infants of uncomplicated pregnancy, controlling for necessary covariates. Extensive demographic information was collected, which measured the most common covariates not controlled for in other studies and, if related to breastfeeding or infant weight, these were controlled for in subsequent analyses. It was hypothesised that infants breastfed for longer durations would: (1) show slower weight gain throughout the first year; (2) weigh less at 12-months; (3) demonstrate more obesity-protective eating behaviours at 12-months; and (4) have mothers who were observed to be more sensitive during feeding, than infants breastfed for shorter durations.

Materials and methods

The study protocol received full ethical approval from Birmingham East, North, and Solihull Research Ethics Committee, United Kingdom (reference number 10/H1206/67). Research and development approval was granted by Birmingham Women's National Health Service Foundation Trust (reference number 10/BWH/NO95).

Mothers were eligible to take part in the study if they had given birth on a low-risk maternity unit and if their infant was not born prematurely (prior to 36 weeks gestation) or small for gestational age (SGA). Premature and SGA infants were not included as these factors are associated with weight gain during the first 12-months of life. Mothers needed to be able to read and write English due to the requirement of completing questionnaires and the ability to communicate with the researcher. Midwives directed the researcher to women who met these criteria.

Two hundred and eighty-seven women were eligible to take part in the study and were approached after delivery on low-risk maternity units of Birmingham Women's Hospital. Of these, 81 mothers (28%) gave informed consent and agreed to be visited at home (mean age 29.42 years [SD 5.87]). Infants included 45 males and 36 females (mean birth-weight 3.52 kg [SD 0.39]).

Mothers and infants were visited at home at 1-week, 1-, 6- and 12-months postpartum. Demographics were reported at 1-week. Mothers and infants were weighed and measured at each visit. Mothers reported feeding information (exclusivity and duration of breastfeeding and when solids were first introduced) and completed questionnaires assessing symptoms of postnatal depression, at each visit. Mothers also reported their smoking and alcohol consumption and any medications they were taking at each visit. Mothers were observed feeding their infant solid food at 6- and 12-months and reported their infant's eating behaviours at 12-months.

Demographic and Additional Information

Mothers completed a demographic questionnaire at 1-week. It requested age, pre-pregnancy weight, ethnic background, household income, educational level and infant date of birth. It also asked the type of milk the mother intended to feed her baby (breast, formula or a mix of the two). Mothers completed an additional information sheet at each visit, which requested information regarding medications being taken and present smoking and alcohol consumption.

Feeding Information

At each visit, mothers reported whether infants were being breast or formula-fed, and the duration and exclusivity of feeding method. Bottle use among breastfeeding mothers was not measured. At the later time points, mothers were asked if and when they had introduced solid foods.

Edinburgh Postnatal Depression Scale (EPDS(Cox, Holden, & Sagovsky, 1987))

Postnatal depression is associated with maternal-infant interactions (Goodman, 2007) and breastfeeding duration (Henderson, Evans, Straton, Priest, & Hagan, 2003). The EPDS was therefore given to mothers at the 1-, 6- and 12-month visit to establish whether depression needed to be controlled for in the analyses. The EPDS consists of 10 short statements, each of which has four responses to choose from, indicating how the mother has felt during the previous week. Mothers who score 10 or greater are identified as showing symptoms indicative of possible depression.

Child Eating Behaviour Questionnaire (CEBQ; (Wardle, Guthrie, Sanderson, & Rapoport, 2001)

The CEBQ is a reliable and valid parent-rated questionnaire measuring eating styles of children using a five-point rating scale. A modified age-appropriate version of the CEBQ was given at the

12-month visit to assess maternal perception of infants' obesogenic and obesity-protective eating behaviours. Subscales measuring emotional over- and under-eating were deemed not appropriate for infants aged 12-months and so were not included. The original CEBQ consists of 35-items and the current modified version consists of 23-items. The modified version was piloted on 59 mothers of infants with a mean age of 7.5-months. Overall reliability was shown to be good to moderate (.62). The Cronbach's alphas for the six subscales were .83 for enjoyment of food and satiety responsiveness, .74 for slowness in eating, .84 for food fussiness, .85 for responsiveness and .88 for desire to drink.

Mealtime observation

The Feeding Interaction Scale (FIS; (Wolke, Sumner, McDermott, & Skuse, 1992) was used to code positive maternal behaviours and some infant eating behaviours during the feeding observations (Table 1 details subscales used and behaviours assessed). In order to investigate observable warm and sensitive feeding behaviours, maternal vocalisations and appropriateness were chosen in addition to sensitivity. The FIS has clinical validity and has been used to assess maternal-infant feeding interactions and diagnose feeding problems (Farrow & Blissett, 2005; Lindberg, Bohlin, Hagekull, & Palmerus, 1996; Skuse, Wolke, & Reilly, 1992).

Feeding sessions took place at participants' homes and were recorded using a video-camcorder and tripod. Feeding observations of solid food took place at either lunch or dinnertime and did not include milk feeds. Mothers informed the researcher what time the meal would be and decided what to feed their infant. There was no restriction imposed regarding when the child last ate. Videos were watched and scored later by the researcher and research assistant. Intra-class

correlation coefficients were all greater than .76.

Table 1. Subscales and behaviours utilised from the FIS (Wolke et al., 1992).

Subscale	Behaviour	Scoring
Maternal verbal involvement	Proportion of session mother is talking to infant including initiating conversation and spontaneous comments	1 (never talks to infant) to 9 (very much)
Appropriateness of maternal mealtime behaviour	Feeding is appropriate if it is pleasurable for mother and infant.	1 (very inappropriate) to 5 (very appropriate)
Maternal sensitivity	Infant in sensible position including freedom of arm movement and eye contact with mother, close proximity to mother, feedback on infant's behaviour, variation of stimulation	1 (highly insensitive) to 9 (highly sensitive)
Frequency of offers	Offers (mother-to-infant or infant-to-self) semi-solid or solid food. An offer is defined as food which reaches within 5 inches of the infant's mouth	
Frequency of acceptances	Food is counted as accepted when it is kept in the mouth for longer than 5 seconds	

Anthropometric Measures

Infants were weighed naked with Seca electronic baby scales by the researcher at each home visit. Infant weight was then converted to a standard deviation score (SDS), which adjusts measurements for age and sex (Freeman et al., 1995). Mothers were weighed at each home visit

wearing light indoor clothing, without shoes, using electronic scales; maternal height was measured at 1-week postpartum using a portable stadiometer.

Data analysis

Kolmogorov-Smirnov tests and histograms indicated that breastfeeding duration, demographic factors and postnatal depression were not normally distributed. Two-tailed non-parametric Spearman's rho correlations were therefore used to assess whether these variables were associated with breastfeeding duration.

One-tailed partial correlations (controlling for: household income category, maternal age, education, BMI and quantity of cigarettes smoked during pregnancy, infant birth weight SDS and age at which introduced to solids) were used to assess the relationship between: (1) breastfeeding duration and infant weight SDS at 1-week, 1-, 6-, and 12-months, weight gain SDS from 1- to 6- and 1- to 12-months; (2) breastfeeding duration and infant eating behaviours at 12-months. One-tailed partial correlations (controlling for: household income category, maternal age, education and quantity of cigarettes smoked during pregnancy, and infant age introduced to solids) were used to assess the relationships between breastfeeding duration and observed positive maternal feeding behaviours. Post hoc analyses included partial correlations to assess whether controlling for maternal sensitivity, in addition to aforementioned covariates, affected the relationship between breastfeeding duration, infant weight and eating behaviours.

Results

Descriptive statistics

Eighty-one mother-infant dyads were initially recruited; at the 12-month visit 12 had withdrawn, resulting in a dropout rate of 15%. Mothers who withdrew reported leaving the study due to moving away or having other demands on their time (e.g. caring for other children, returning to work [data not shown]). Table 2 shows the number of mother-infant dyads seen at each home visit, the mean age of infants (weeks) and the percentage of infants being breastfed at each time point (includes exclusive and any breastfeeding). Of the 73% breastfeeding at 1-week, 75% of these were exclusively breastfeeding. Of the 65% breastfeeding at 1-month, 76% of these were exclusively breastfeeding. Of the 52% breastfeeding at 6-months and 32% at 12-months, 71% and 64% had not introduced formula or cow's milk respectively. There was no difference in breastfeeding duration between male ($M = 24.34$, $SE = 3.58$) and female ($M = 30.58$, $SE = 4.03$) infants $t(67) = -1.16$, $p = .25$.

Group comparisons between 'exclusive' 'partial' and 'no' breastfeeding were not conducted due to the small group sizes. Infants partially fed breast milk may have received formula twice per week or multiple times per day and so it was deemed inappropriate to group such infants together in one category. Furthermore, five infants were introduced to solid food before 12-weeks and an additional 59 were introduced to solids before 24-weeks. Timing of introduction of solid food added to the complexity of generating 'pure' groups in terms of breastfeeding exclusivity.

Table 2 also shows that no infants were below the 2nd centile for weight at 1-, 6-, or 12-months. These centiles were plotted using the UK-WHO growth charts. There were no significantly underweight infants in the current sample. As meal content can affect interactions during mealtimes, mothers rated infant familiarity and liking of the food presented. Infants were generally given food they liked and were familiar with (Table 3). The mean age infants were introduced to solid food was 20.41 weeks (SD 3.39). There was no difference between male ($M = 20.20, SE = 0.60$) and female ($M = 20.65, SE = 0.51$) infants in the age at which they were introduced to solid food $t(71) = -.57, p = .57$.

1 **Table 2.** Number of infants, mean age (weeks), percentage receiving any breast milk and weight and centile range at each home visit
 2 (according to the UK-WHO growth charts).

3

Visit	N	Mean age (weeks)	Any breastfeeding	Weight range (kg)		Centile range	
				Males	Females	Males	Females
1-week	81	1.32 (SD 0.36)	73%	2.72 – 4.88	2.81 – 4.37	n/a [§]	n/a [§]
1-month	77	4.77 (SD 0.62)	65%	3.43 – 6.00	3.74 – 5.39	2 nd – 98 th	9 th – 91 st
6-months	73	26.67 (SD 0.99)	52%	6.46 – 10.50	6.59 – 9.38	2 nd – 98 th	25 th – 98 th
12-months	69	52.83 (SD1.73)	32%	8.00 – 12.81	7.71 – 11.82	2 nd – 99.6 th	9 th – 98 th

4 [§]UK-WHO growth charts provide centiles for males and females from 2-weeks to 4-years-old

5

6

7

8 **Table 3.** Maternal ratings of infant’s familiarity and liking of food given during feeding sessions

	Familiarity				Liking (Mean and S.D.)
	Never	Once	A few times	Often	
6-months	5.2%	10.3%	32.8%	51.7%	82.4% (S.D. 17.0%)
12-months	3.6%	5.5%	25.5%	65.5%	80.1% (S.D. 17.0%)

9

10 **Covariates**

11 EPDS score was not significantly associated with breastfeeding duration at: 1-month $r=.21$; 6-
12 months $r=.06$; or 12-months $r=.16$, all $p>.05$. Therefore, postnatal depression was not controlled
13 for in any further analyses.

14
15 One-tailed Spearman's rho correlations revealed that maternal age and educational level were
16 significantly associated with breastfeeding duration at each visit (see Table 4). There were
17 positive associations between breastfeeding duration and household income at 1-week, 1-month
18 and 6-months; positive associations between breastfeeding at 6-months and 12-months and the
19 age that infants were introduced to solid food; and negative associations between breastfeeding
20 duration and cigarettes smoked during pregnancy at 1-week, 1- and 6-months. The
21 aforementioned variables were controlled in further analyses. Birth weight was not related to
22 breastfeeding duration at any point.

23

24

25

26

27

28

29

30

31

32

33

34 **Table 4.** Spearman's Rho bivariate correlations (one-tailed) between breastfeeding duration and
 35 covariates at each time point of the study.
 36

Visit		Birth weight SDS	Household income	Maternal age	Maternal education	Cigarettes smoked during pregnancy	Age infant introduced to solids
1-week	<i>r</i>	-.02	.28	.38	.38	-.21	n/a [§]
	<i>p</i>	.44	.01	<.0001	<.0001	.03	n/a [§]
	<i>n</i>	81	81	81	81	81	n/a [§]
1-month	<i>r</i>	-.12	.34	.41	.43	-.26	n/a [§]
	<i>p</i>	.15	.001	<.0001	<.0001	.01	n/a [§]
	<i>n</i>	77	77	77	77	77	n/a [§]
6-months	<i>r</i>	-.06	.26	.42	.37	-.27	.26
	<i>p</i>	.31	.01	<.0001	.001	.01	.01
	<i>n</i>	73	73	73	73	73	73
12-months	<i>r</i>	-.08	.15	.25	.26	-.13	.21
	<i>p</i>	.26	.11	.02	.02	.14	.04
	<i>n</i>	68	68	68	68	68	68

37 n/a[§] No infants had been introduced to solid food at 1-week or 1-month
 38
 39

40 **Breastfeeding and observations**

41 One-tailed partial correlations were conducted to investigate the relationship between
 42 breastfeeding duration and observed positive maternal mealtime behaviours at 12-months. Table
 43 5 shows that after accounting for covariates, breastfeeding duration was associated with more
 44 positive maternal vocalisations and sensitivity during the meal at 12-months and more
 45 appropriateness at 6- and 12-months.

46 **Breastfeeding and infant weight**

47 One-tailed partial correlations were conducted to investigate the relationship between
48 breastfeeding duration and infant weight SDS. Table 6 shows that (concurrent) breastfeeding
49 duration was negatively associated with infant weight SDS at 6- and 12-months and weight gain
50 SDS from 1- to 6- and 1- to 12-months, but not with infant weight SDS at 1-week or 1-month.
51 The effects seen here are substantial. For example, a 6-month-old boy on the 50th centile weighs
52 8.05kg, a reduction of 1 SD at this time results in a weight of 7.74kg, representing a total
53 reduction in weight of 0.32kg.

54
55 Given the relationship between maternal behaviours and breastfeeding duration, post hoc tests
56 were conducted to investigate whether controlling for maternal sensitivity affected the
57 relationship between breastfeeding duration, infant weight and eating behaviours. Maternal
58 behaviours were highly correlated so, for parsimony and to protect power, only sensitivity was
59 controlled. Controlling for maternal sensitivity did not change the pattern of these results.

60

61 **Breastfeeding and infant eating behaviours**

62 One-tailed partial correlations were conducted to investigate the relationship between
63 breastfeeding duration and infant eating behaviours at 12-months. Table 7 shows that after
64 accounting for covariates, breastfeeding duration was positively associated with slowness in
65 eating at 12-months. Breastfeeding duration was also negatively associated with the number of
66 times mothers offered their infants food at 12-months. There were no other relationships between
67 duration of breastfeeding and maternally reported or observed infant eating behaviours.
68 Controlling for maternal sensitivity did not change the pattern of these results.

69

70 **Table 5.** Partial correlations (one-tailed) between breastfeeding duration and observed positive
 71 maternal mealtime behaviours. Covariates include: maternal age, education, number of cigarettes
 72 smoked during pregnancy, household income and age introduced to solids.
 73

		Vocalisations	Appropriateness	Sensitivity
6-months	Breastfeeding Duration	.12	.26	.22
	<i>p</i>	.21	.03	.07
	<i>df</i>	48	48	48
12-months	Breastfeeding Duration	.24	.32	.29
	<i>p</i>	.05	.01	.02
	<i>df</i>	48	48	48

74

75 **Table 6.** Partial correlations (one-tailed) between concurrent breastfeeding duration and infant weight SDS controlling for maternal age,
 76 education, concurrent BMI, number of cigarettes smoked during pregnancy, household income, infant birth weight SDS and age introduced
 77 to solids.
 78

	1-week weight SDS	1-month weight SDS	6-month weight SDS	12-month weight SDS	1- to 6-month SDS weight gain	1- to 12-month SDS weight gain
Breastfeeding Duration	-.001	-.15	-.33	-.39	-.30	-.38
<i>p</i>	.50	.12	.004	.001	.01	.001
<i>df</i>	64	64	63	58	63	58
<i>Breastfeeding Duration (cont. sensitivity)</i>			-.47	-.45	-.44	-.41
<i>p</i>			<.0001	.001	.001	.002
<i>df</i>			45	44	45	44

79
 80
 81
 82
 83
 84
 85
 86
 87
 88
 89
 90
 91
 92
 93

94
95
96
97

Table 7. Partial correlations (one-tailed) between breastfeeding duration and infant eating behaviours. Covariates include: maternal age, education, BMI, number of cigarettes smoked during pregnancy, household income, infant birth weight SDS and age introduced to solids.

		Maternal report of infant eating behaviours (CEBQ)						Observed eating behaviours (FIS)			
		Satiety responsiveness	Food enjoyment	Food responsiveness	Slowness in eating	Food fussiness	Desire to drink	Self offers	Self acceptances	Maternal offers	Maternal acceptances
12-months	Breastfeeding Duration	.11	-.08	-.11	.25	.04	.03	.17	.17	-.25	-.19
	<i>p</i>	.19	.27	.21	.03	.39	.41	.13	.13	.04	.10
	<i>df</i>	59	59	59	59	59	59	46	46	46	46
	<i>Breastfeeding Duration (cont. sensitivity)</i>	.09	-.12	-.07	.28	.01	.05	.16	.16	-.20	-.13
	<i>p</i>	.28	.22	.32	.03	.47	.36	.14	.14	.09	.19
	<i>df</i>	45	45	45	45	45	45	45	45	45	45

98

99 **Discussion**

100 The results of this study supported the hypothesis that a longer duration of
101 breastfeeding is associated with slower weight gain from 1- to 6- and 1- to 12-months
102 and lower weight at 6- and 12-months. Findings are supportive of previous research
103 (Arenz et al., 2004; Harder et al., 2005; Hornell et al., 2013; Kramer, 1981; McCrory
104 & Layte, 2012; Owen et al., 2005) and are consistent with a dose-response
105 relationship between breastfeeding and infant weight during the first 12-months of
106 life, which most clearly manifests in the latter half of the first year.

107

108 Results of the current study also show that infants breastfed for longer are perceived
109 by their mothers to eat solid food more slowly at 12-months. It is possible that
110 breastfeeding influences the development of slower eating styles, which may in turn
111 enhance satiety responsiveness in early childhood. Previous research with adults has
112 found that that slower eating increases self-reported satiety (Andrade & Melanson,
113 2007; Ferriday et al., 2015; Shah et al., 2014). Further research is required to establish
114 whether breastfeeding helps infants develop obesity-protective eating behaviours.

115 This is likely, given that breastfed infants adjust their intake according to the ever-
116 changing fat content of the milk (Tyson et al., 1992) and may therefore better learn
117 their internal cues of hunger and satiety (Birch & Fisher, 1998) than those fed formula
118 milk. Breastfeeding may also encourage a slower rate of eating due to breastfed
119 infants having to work harder for their food than formula-fed infants; introducing a
120 bottle may speed-up feeding rate (Cao et al., 2009).

121

122 Although previous research has found significant relationships between breastfeeding
123 duration and satiety responsiveness (Brown & Lee, 2012), infants were 6- to 12-

124 months younger in the current study. Further research is required to investigate
125 whether relationships between slowness in eating and weight extend to satiety
126 responsiveness after infancy.

127

128 It is also important to consider that such research did not control for the age at which
129 infants were introduced to solid food (Brown & Lee, 2012). Previous research has
130 found that breastfed infants, and those breastfed for longer, tend to be introduced to
131 solid food later than those breastfed for shorter durations or not at all (Huh, Rifas-
132 Shiman, Taveras, Oken, & Gillman, 2011). Furthermore, earlier introduction of solid
133 food has also been related to greater weight gain during the first year of life (Baird et
134 al., 2008; Baker, Michaelsen, Rasmussen, & Sorensen, 2004; Forsyth, Ogston, Clark,
135 Florey, & Howie, 1993; Kramer et al., 1985; Lande et al., 2005). Timing of
136 introduction of solid food was controlled for in the current study as it was related to
137 both breastfeeding duration and infant weight.

138

139 The current research did not find any significant relationships between breastfeeding
140 duration and food enjoyment, food responsiveness, fussiness or desire to drink, as
141 measured by parental report at 1 year. Future studies are required to investigate
142 whether relationships between breastfeeding duration and these eating behaviours
143 emerge after infancy, once children have more control during feeding and are more
144 able to feed themselves. Furthermore, in the current study, measures were taken
145 before the stage at which increases in fussy/picky eating and neophobia are seen
146 (Taylor, Wernimont, Northstone, & Emmett, 2015). Future studies could explore
147 these factors in early childhood as it is possible that some of these infants may go on
148 to develop fussier eating habits with time.

149

150 Although it is possible that longer breastfeeding may promote slower weight gain
151 resulting in lower weight at 12-months, it is important to consider that causality
152 cannot be assumed. Less hungry infants, or those with smaller appetites, may be less
153 demanding and easier to breastfeed, and so are breastfed for longer. Indeed, it has
154 been reported that one of the main reasons why mothers stop breastfeeding is because
155 they perceive their infant was no longer satisfied by breast milk alone (Li, Fein, Chen,
156 & Grummer-Strawn, 2008). In addition, mothers who feed to comfort and soothe their
157 infant may also breastfeed for shorter durations (Paul et al., 2011). It is possible that
158 these infants may gain weight more slowly and may develop slower eating styles due
159 to their smaller appetite rather than it being due to breastfeeding alone. However,
160 weight at birth, 1-week and 1-month was not related to breastfeeding duration in the
161 current study, which suggests that it was not only the smaller infants who were
162 breastfed for longer in this sample. Neither did breastfeeding duration relate to
163 observations of infant eating behaviour such as the frequency of infant self-offering or
164 accepting of food, suggesting that breastfeeding was not related to this index of infant
165 appetite.

166

167 Whilst considering the results of this study it is important to take into account that
168 infant feeding cues are influenced by environmental, physical and psychological
169 factors; perception of these cues is affected by both maternal and infant characteristics
170 (McNally et al., 2016). Mothers perceive hunger signals more easily than satiety
171 signals, but interpretation of feeding cues does get easier as children age (McNally et
172 al., 2016). Future research into early weight gain and eating should therefore move
173 forward by investigating the impact of observed infant characteristics on feeding

174 behaviours. An improved understanding of the factors affecting the interpretation of,
175 and response to, infant feeding cues will aid the development of interventions to
176 promote sensitive and responsive feeding.

177

178 Results of the current study show that increased maternal positive vocalisations and
179 greater observed maternal appropriateness and sensitivity during a mealtime at 12-
180 months are significantly associated with a longer duration of breastfeeding. This
181 supports previous research that found mothers who demonstrate greater maternal
182 sensitivity during infancy and higher quality interactions at 12-months, breastfeed for
183 longer (Britton, Britton, & Gronwaldt, 2006; Gutman, Brown, & Akerman, 2009;
184 Tharner et al., 2012). Results are also supportive of research that has found
185 breastfeeding mothers provide a more ideal feeding environment and feed more
186 responsively than those who bottle feed or feed solids (Shloim et al., 2015). Although
187 causality cannot be inferred from these analyses, it is possible that breastfeeding
188 increases positive maternal behaviours. However, it is also true that more sensitive
189 mothers choose to breastfeed (Tharner et al., 2012). Breastfeeding for at least 6-
190 months may therefore be a practice that more sensitive mothers undertake. It is
191 interesting to highlight here that post hoc analyses demonstrated controlling for
192 maternal sensitivity did not remove the significance of the relationship between
193 breastfeeding duration and weight or slowness in eating. Whilst maternal sensitivity is
194 an important correlate of breastfeeding duration, it does not entirely explain the
195 relationship between breastfeeding and weight/eating behaviour.

196

197 Interestingly, previous research has found that the majority of mothers observed to be
198 responsive to their child during a mealtime had children who were highly responsive

199 to their mothers in return (Hodges et al., 2013). Future research should therefore
200 evaluate parenting sensitivity and responsiveness from a bidirectional perspective,
201 using longitudinal observational methods (Bergmeier, Skouteris, & Hetherington,
202 2015).

203

204 One limitation of the current study is that the sample size is small for the number of
205 variables that were controlled. In addition to this, it is uncertain whether all relevant
206 confounders have been included. However, the current study did assess and control
207 for a large number of important confounders, unlike many previous studies, and has
208 still found some evidence for the relationship between breastfeeding and weight in
209 infancy.

210

211 Although participants were from a variety of demographic, socioeconomic and
212 cultural backgrounds, the educational level achieved by mothers in the current study
213 was significantly higher than the national average (Statistics, 2011). In 2011, 27% of
214 the UK adult population had a Level 4 qualification or above (degree, higher degree
215 or professional qualification), compared to 63% of the mothers in the current study.
216 Furthermore, although fewer women in the current study initiated breastfeeding
217 compared to the national average (75% versus 81%), a higher proportion of women in
218 the current study were breastfeeding at 6-months compared to the UK average (52%
219 versus 34%) (McAndrew, 2010). It is therefore possible that selection bias may have
220 affected whether or not participants continued with the study.

221

222 In addition to this, the current study did not measure the feeding of expressed breast
223 milk via bottles. It is therefore assumed that breastfed infants were fed directly from

224 the breast. Given the increase of feeding expressed milk in recent years (Labiner-
225 Wolfe, Fein, Shealy, & Wang, 2008), when investigating health outcomes in infancy
226 and childhood, future studies should assess the mode by which breast milk is fed as
227 well as the exclusivity and duration of breastfeeding.

228

229 **Conclusions**

230 The current study contributes to the published literature suggesting that breastfeeding
231 is significantly associated with slower weight gain and lower weight and BMI
232 throughout the first year of life. Furthermore, breastfeeding may also encourage the
233 development of obesity-protective eating behaviours through the development of
234 slower eating styles. Slower eating styles may help infants and mothers in the
235 attention, communication and perception of internal signals of hunger and satiety.
236 Future research into breastfeeding and weight gain should move forward by
237 investigating whether relationships between slowness in eating and weight extend to
238 satiety responsiveness after infancy.

239

240

241

242

243

244

245

246

247

248

249 **Acknowledgements**

250 This study was supported by the Economic and Social Research Council Studentship
251 Award ES/G017786/1. The authors have no conflicts of interest or financial
252 relationships relevant to this article to disclose.

253 Samantha L Rogers: Dr. Rogers jointly conceptualised and designed the study,
254 collected the data, contributed to its analysis and interpretation, drafted the initial
255 manuscript, and approved the final manuscript as submitted.

256 Jackie Blissett: Professor Blissett conceptualised and designed the study, supervised
257 data collection, contributed to analysis and interpretation of data, critically reviewed
258 the manuscript, and approved the final manuscript as submitted.

259 Both authors had full access to all of the data in the study and take responsibility for
260 the integrity of the data and the accuracy of the data analysis. Both authors approved
261 the final manuscript as submitted and agree to be accountable for all aspects of the
262 work.

263

264

265

266

267

268

269

270

271

272

273

274

275

276

277

278 **References**

- 279 Andrade, A. M., & Melanson, K. J. (2007). Is eating slowly a potential strategy to
280 maximize satiety? *FASEB Journal*, *21*(5), A319.
- 281 Arenz, S., Ruckerl, R., Koletzko, B., & Von Kries, R. (2004). Breast-feeding and
282 childhood obesity - A systematic review. *International Journal of Obesity and*
283 *Related Metabolic Disorders*, *28*(10), 1247-1256.
284 doi:doi:10.1038/sj.ijo.0802758
- 285 Baird, J., Poole, J., Robinson, S., Marriott, L., Godfrey, K., Cooper, C., . . .
286 Southampton Women's Survey Study, G. (2008). Milk feeding and dietary
287 patterns predict weight and fat gains in infancy. *Paediatric and Perinatal*
288 *Epidemiology*, *22*(6), 575-586. doi:10.1111/j.1365-3016.2008.00963.x
- 289 Baker, J. L., Michaelsen, K. F., Rasmussen, K. M., & Sorensen, T. I. A. (2004).
290 Maternal prepregnant body mass index, duration of breastfeeding, and timing
291 of complementary food introduction are associated with infant weight gain.
292 *American Journal of Clinical Nutrition*, *80*(6), 1579-1588.
- 293 Bergmeier, H., Skouteris, H., Haycraft, E., Haines, J., & Hooley, M. (2015). Reported
294 and observed controlling feeding practices predict child eating behavior after
295 12 months. *The Journal of nutrition*, *145*(6), 1311-1316.
296 doi:10.3945/jn.114.206268
- 297 Bergmeier, H., Skouteris, H., & Hetherington, M. (2015). Systematic research review
298 of observational approaches used to evaluate mother-child mealtime
299 interactions during preschool years. *The American journal of clinical*
300 *nutrition*, *101*(1), 7-15. doi:10.3945/ajcn.114.092114
- 301 Birch, L. L., & Fisher, J. O. (1998). Development of eating behaviors among children
302 and adolescents. *Pediatrics*, *101*(3 II SUPPL.), 539-549.
- 303 Britton, J. R., Britton, H. L., & Gronwaldt, V. (2006). Breastfeeding, sensitivity, and
304 attachment. *Pediatrics*, *118*(5), e1436-1443. doi:10.1542/peds.2005-2916
- 305 Brown, A., & Lee, M. (2012). Breastfeeding during the first year promotes satiety
306 responsiveness in children aged 18-24 months. *Pediatric Obesity*, *7*(5), 382-
307 390.
- 308 Cao, Y., Rao, S. D., Phillips, T. M., Umbach, D. M., Bernbaum, J. C., Archer, J. I., &
309 Rogan, W. J. (2009). Are Breast-fed Infants More Resilient? Feeding Method
310 and Cortisol in Infants. *Journal of Pediatrics*, *154*(3), 452-454.
- 311 Cox, J. L., Holden, J. M., & Sagovsky, R. (1987). Detection of Postnatal Depression:
312 Development of the 10-item Edinburgh Postnatal Depression scale. *British*
313 *Journal of Psychiatry*, *150*(JUNE), 782-786.
- 314 Davis, J. N., Weigensberg, M. J., Shaibi, G. Q., Crespo, N. C., Kelly, L. A., Lane, C.
315 J., & Goran, M. I. (2007). Influence of breastfeeding on obesity and type 2
316 diabetes risk factors in latino youth with a family history of type 2 diabetes.
317 *Diabetes Care*, *30*(4), 784-789.
- 318 Farrow, C. V., & Blissett, J. M. (2005). Is maternal psychopathology related to
319 obesigenic feeding practices at 1 year? *Obesity Research*, *13*(11), 1999-2005.
320 doi:10.1038/oby.2005.245
- 321 Ferriday, D., Bosworth, M. L., Lai, S., Godinot, N., Martin, N., Martin, A. A., . . .
322 Brunstrom, J. M. (2015). Effects of eating rate on satiety: A role for episodic
323 memory? *Physiol Behav*, *152*(Pt B), 389-396.
324 doi:10.1016/j.physbeh.2015.06.038
- 325 Forsyth, J. S., Ogston, S. A., Clark, A., Florey, C. D. V., & Howie, P. W. (1993).
326 Relation between early introduction of solid food to infants and their weight

327 and illnesses during the first two years of life. *British Medical Journal*,
328 306(6892), 1572-1576.

329 Freeman, J. V., Cole, T. J., Chinn, S., Jones, P. R., White, E. M., & Preece, M. A.
330 (1995). Cross sectional stature and weight reference curves for the UK, 1990.
331 *Arch Dis Child*, 73(1), 17-24.

332 Gillman, M. W., Rifas-Shiman, S. L., Berkey, C. S., Frazier, A. L., Rockett, H. R. H.,
333 Camargo, C. A., Jr., . . . Colditz, G. A. (2006). Breast-feeding and overweight
334 in adolescence: within-family analysis corrected. *Epidemiology (Cambridge,*
335 *Mass.)*, 17(1), 112-114. doi:10.1097/01.ede.0000181629.59452.95

336 Gillman, M. W., Rifas-Shiman, S. L., Camargo, C. A., Jr., Berkey, C. S., Frazier, A.
337 L., Rockett, H. R. H., . . . Colditz, G. A. (2001). Risk of overweight among
338 adolescents who were breastfed as infants. *JAMA (Journal of the American*
339 *Medical Association)*, 285(19), 2461-2467. doi:10.1001/jama.285.19.2461

340 Goodman, S. H. (2007). Depression in mothers. *Annual review of clinical psychology*,
341 3, 107-135. doi:10.1146/annurev.clinpsy.3.022806.091401

342 Gutman, L. M., Brown, J., & Akerman, R. (2009). Nurturing parenting capability:
343 The early years. Centre for Research on the Wider Benefits of Learning.
344 Retrieved from
345 <http://www.learningbenefits.net/Publications/ResReps/ResRep30.pdf>

346 Harder, T., Bergmann, R., Kallischnigg, G., & Plagemann, A. (2005). Duration of
347 breastfeeding and risk of overweight: a meta-analysis. *Am J Epidemiol*,
348 162(5), 397-403. doi:10.1093/aje/kwi222

349 Heinig, M. J., Nommsen, L. A., Peerson, J. M., Lonnerdal, B., & Dewey, K. G.
350 (1993). Energy and protein intakes of breast-fed and formula-fed infants
351 during the first year of life and their association with growth velocity: The
352 darling study. *American Journal of Clinical Nutrition*, 58(2), 152-161.

353 Henderson, J. J., Evans, S. F., Straton, J. A. Y., Priest, S. R., & Hagan, R. (2003).
354 Impact of postnatal depression on breastfeeding duration. *Birth (Berkeley,*
355 *Calif.)*, 30(3), 175-180. doi:10.1046/j.1523-536X.2003.00242.x

356 Hodges, E. A., Johnson, S. L., Hughes, S. O., Hopkinson, J. M., Butte, N. F., &
357 Fisher, J. O. (2013). Development of the responsiveness to child feeding cues
358 scale. *Appetite*, 65, 210-219. doi:10.1016/j.appet.2013.02.010

359 Hornell, A., Lagstrom, H., Lande, B., & Thorsdottir, I. (2013). Breastfeeding,
360 introduction of other foods and effects on health: a systematic literature review
361 for the 5th Nordic Nutrition Recommendations. *Food Nutr Res*, 57.
362 doi:10.3402/fnr.v57i0.20823

363 Huh, S. Y., Rifas-Shiman, S. L., Taveras, E. M., Oken, E., & Gillman, M. W. (2011).
364 Timing of solid food introduction and risk of obesity in preschool-aged
365 children. *Pediatrics*, 127(3), e544-551. doi:10.1542/peds.2010-0740

366 Jenness, R. (1979). The composition of human milk. *Semin Perinatol*, 3(3), 225-239.

367 Jiang, M., & Foster, E. M. (2013). Duration of breastfeeding and childhood obesity: a
368 generalized propensity score approach. *Health Serv Res*, 48(2 Pt 1), 628-651.
369 doi:10.1111/j.1475-6773.2012.01456.x

370 Kramer, M. S. (1981). Do breast-feeding and delayed introduction of solid foods
371 protect against subsequent obesity? *The Journal of Pediatrics*, 98, 883-887.

372 Kramer, M. S., Barr, R. G., Leduc, D. G., Boisjoly, C., McVey-White, L., & Pless, I.
373 B. (1985). DETERMINANTS OF WEIGHT AND ADIPOSITY IN THE 1ST
374 YEAR OF LIFE. *Journal of Pediatrics*, 106(1), 10-14. doi:10.1016/s0022-
375 3476(85)80456-x

376 Kudlová, E., & Schneidrová, D. (2012). Dietary patterns and their changes in early
377 childhood. *Central European Journal of Public Health*, 20(2), 126-134.

378 Labiner-Wolfe, J., Fein, S. B., Shealy, K. R., & Wang, C. (2008). Prevalence of breast
379 milk expression and associated factors. *Pediatrics*, 122(Suppl. S), S63-S68.
380 doi:10.1542/peds.2008-1315h

381 Lande, B., Andersen, L. F., Henriksen, T., Baerug, A., Johansson, L., Trygg, K. U., . .
382 . Veierod, M. B. (2005). Relations between high ponderal index at birth,
383 feeding practices and body mass index in infancy. *Eur J Clin Nutr*, 59(11),
384 1241-1249. doi:10.1038/sj.ejcn.1602235

385 Li, R., Fein, S. B., Chen, J., & Grummer-Strawn, L. M. (2008). Why mothers stop
386 breastfeeding: Mothers' self-reported reasons for stopping during the first year.
387 *Pediatrics*, 122(Suppl. S), S69-S76. doi:10.1542/peds.2008-1315i

388 Lindberg, L., Bohlin, G., Hagekull, B., & Palmerus, K. (1996). Interactions between
389 mothers and infants showing food refusal. *Infant Mental Health Journal*,
390 17(4), 334-347.

391 Martin, R. M., Patel, R., Kramer, M. S., Guthrie, L., Vilchuck, K., Bogdanovich, N., .
392 . . Oken, E. (2013). Effects of promoting longer-term and exclusive
393 breastfeeding on adiposity and insulin-like growth factor-I at age 11.5 years: A
394 randomized trial. *JAMA - Journal of the American Medical Association*,
395 309(10), 1005-1013.

396 McAndrew, F., Thompson, J., Fellows, L., Large, A., Speed, M., & Renfrew, M.J.
397 (2010). Infant Feeding Survey 2010: Summary. NHS Information Centre for
398 Health and Social Care [PDF document]. Retrieved from
399 [https://catalogue.ic.nhs.uk/publications/public-health/surveys/infant-feed-](https://catalogue.ic.nhs.uk/publications/public-health/surveys/infant-feed-surv-2010/ifs-uk-2010-sum.pdf)
400 [surv-2010/ifs-uk-2010-sum.pdf](https://catalogue.ic.nhs.uk/publications/public-health/surveys/infant-feed-surv-2010/ifs-uk-2010-sum.pdf)

401 McCrory, C., & Layte, R. (2012). Breastfeeding and risk of overweight and obesity at
402 nine-years of age. *Soc Sci Med*, 75(2), 323-330.
403 doi:10.1016/j.socscimed.2012.02.048

404 McNally, J., Hugh-Jones, S., Caton, S., Vereijken, C., Weenen, H., & Hetherington,
405 M. (2016). Communicating hunger and satiation in the first 2years of life: a
406 systematic review. *Matern Child Nutr*, 12(2), 205-228.
407 doi:10.1111/mcn.12230

408 Mennella, J. A., Jagnow, C. P., & Beauchamp, G. K. (2001). Prenatal and Postnatal
409 Flavor Learning by Human Infants. *Pediatrics*, 107(6), e88-e88.
410 doi:10.1542/peds.107.6.e88

411 Michels, K. B., Willett, W. C., Graubard, B. I., Vaidya, R. L., Cantwell, M. M.,
412 Sansbury, L. B., & Forman, M. R. (2007). A longitudinal study of infant
413 feeding and obesity throughout life course. *Int J Obes (Lond)*, 31(7), 1078-
414 1085. doi:10.1038/sj.ijo.0803622

415 Nommsen, L. A., Lovelady, C. A., Heinig, M. J., Lönnerdal, B., & Dewey, K. G.
416 (1991). Determinants of energy, protein, lipid, and lactose concentrations in
417 human milk during the first 12 mo of lactation: The DARLING Study.
418 *American Journal of Clinical Nutrition*, 53(2), 457-465.

419 Novaes, J. F., Lamounier, J. A., Colosimo, E. A., Franceschini, S. C., & Priore, S. E.
420 (2012). Breastfeeding and obesity in Brazilian children. *Eur J Public Health*,
421 22(3), 383-389. doi:10.1093/eurpub/ckr067

422 Oddy, W. H., Sherriff, J. L., De Klerk, N. H., Kendall, G. E., Sly, P. D., Beilin, L. J., .
423 . . Stanley, F. J. (2004). The relation of breastfeeding and body mass index to
424 asthma and atopy in children: A prospective cohort study to age 6 years. *Am J*
425 *Public Health*, 94(9), 1531-1537.

426 Owen, C. G., Martin, R. M., Whincup, P. H., Smith, G. D., & Cook, D. G. (2005).
427 Effect of infant feeding on the risk of obesity across the life course: a
428 quantitative review of published evidence. *Pediatrics*, *115*(5), 1367-1377.
429 doi:10.1542/peds.2004-1176

430 Paul, I. M., Savage, J. S., Anzman, S. L., Beiler, J. S., Marini, M. E., Stokes, J. L., &
431 Birch, L. L. (2011). Preventing obesity during infancy: a pilot study. *Obesity*
432 (*Silver Spring, Md.*), *19*(2), 353-361. doi:10.1038/oby.2010.182

433 Reynolds, D., Hennessy, E., & Polek, E. (2014). Is breastfeeding in infancy predictive
434 of child mental well-being and protective against obesity at 9 years of age?
435 *Child Care Health Dev*, *40*(6), 882-890. doi:10.1111/cch.12126

436 Sacco, M. R., de Castro, N. P., Euclides, V. L. V., Souza, J. M., & Rondo, P. H. C.
437 (2013). Birth weight, rapid weight gain in infancy and markers of overweight
438 and obesity in childhood. *Eur J Clin Nutr*, *67*(11), 1147-1153.
439 doi:10.1038/ejcn.2013.183

440 Shah, M., Copeland, J., Dart, L., Adams-Huet, B., James, A., & Rhea, D. (2014).
441 Slower eating speed lowers energy intake in normal-weight but not
442 overweight/obese subjects. *Journal of the Academy of Nutrition and Dietetics*,
443 *114*(3), 393-402. doi:10.1016/j.jand.2013.11.002

444 Shloim, N., Rudolf, M. C. J., Feltbower, R. G., Mohebati, L., & Hetherington, M.
445 (2015). Breast is best: Positive mealtime interactions in breastfeeding mothers
446 from Israel and the United Kingdom. *Health Psychology Open*, *2*(1),
447 2055102915579605.

448 Skledar, M. T., & Milosevic, M. (2015). Breastfeeding and time of complementary
449 food introduction as predictors of obesity in children. *Central European*
450 *Journal of Public Health*, *23*(1), 26-31.

451 Skuse, D., Wolke, D., & Reilly, S. (1992). *FAILURE-TO-THRIVE - CLINICAL AND*
452 *DEVELOPMENTAL ASPECTS* (Vol. 2). Toronto: Hogrefe & Huber
453 Publishers.

454 Smithers, L. G., Kramer, M. S., & Lynch, J. W. (2015). Effects of Breastfeeding on
455 Obesity and Intelligence: Causal Insights From Different Study Designs.
456 *JAMA pediatrics*, *169*(8), 707-708. doi:10.1001/jamapediatrics.2015.0175

457 Statistics, O. f. N. (2011). 2011 Census: KS501EW Qualifications and students, local
458 authorities in England and Wales

459 Stettler, N., Zemel, B. S., Kumanyika, S., & Stallings, V. A. (2002). Infant Weight
460 Gain and Childhood Overweight Status in a Multicenter, Cohort Study.
461 *Pediatrics*, *109*(2), 194-199. doi:10.1542/peds.109.2.194

462 Taveras, E. M., Rifas-Shiman, S. L., Scanlon, K. S., Grummer-Strawn, L. M., Sherry,
463 B., & Gillman, M. W. (2006). To what extent is the protective effect of
464 breastfeeding on future overweight explained by decreased maternal feeding
465 restriction? *Pediatrics*, *118*(6), 2341-2348. doi:10.1542/peds.2006-1814

466 Taylor, C. M., Wernimont, S. M., Northstone, K., & Emmett, P. M. (2015).
467 Picky/fussy eating in children: Review of definitions, assessment, prevalence
468 and dietary intakes. *Appetite*, *95*, 349-359. doi:10.1016/j.appet.2015.07.026

469 Tharner, A., Luijk, M. P. C. M., Raat, H., Ijzendoorn, M. H., Bakermans-Kranenburg,
470 M. J., Moll, H. A., . . . Tiemeier, H. (2012). Breastfeeding and its relation to
471 maternal sensitivity and infant attachment. *Journal of developmental and*
472 *behavioral pediatrics : JDBP*, *33*(5), 396-404.
473 doi:10.1097/DBP.0b013e318257fac3

- 474 Tyson, J., Burchfield, J., Sentance, F., Mize, C., Uauy, R., & Eastburn, J. (1992).
475 Adaptation of feeding to a low fat yield in breast milk. *Pediatrics*, 89(2), 215-
476 220.
- 477 Umer, A., Hamilton, C., Britton, C. M., Mullett, M. D., John, C., Neal, W., & Lilly,
478 C. L. (2015). Association between Breastfeeding and Childhood Obesity:
479 Analysis of a Linked Longitudinal Study of Rural Appalachian Fifth-Grade
480 Children. *Childhood obesity (Print)*, 11(4), 449-455.
481 doi:10.1089/chi.2015.0026
- 482 Wardle, J., Guthrie, C. A., Sanderson, S., & Rapoport, L. (2001). Development of the
483 children's eating behaviour questionnaire. *Journal of Child Psychology and*
484 *Psychiatry and Allied Disciplines*, 42(7), 963-970.
- 485 Webber, L., Hill, C., Saxton, J., Van Jaarsveld, C. H. M., & Wardle, J. (2009). Eating
486 behaviour and weight in children. *Int J Obes (Lond)*, 33(1), 21-28.
- 487 Wolke, D., Sumner, M., McDermott, Y., & Skuse, D. (1992). The feeding interaction
488 scale. In H. Remschmidt & M. Schmidt (Eds.), *Child and youth psychiatry:*
489 *European perspectives, Volume II* (Vol. 2, pp. 46-71). Stuttgart: Hans Huber.
- 490 Yan, J., Liu, L., Zhu, Y., Huang, G., & Wang, P. P. (2014). The association between
491 breastfeeding and childhood obesity: a meta-analysis. *BMC Public Health*, 14,
492 1267. doi:10.1186/1471-2458-14-1267
- 493 Zhou, J., Dang, S., Zeng, L., Gao, W., Wang, D., Li, Q., . . . Yan, H. (2016). Rapid
494 Infancy Weight Gain and 7- to 9-year Childhood Obesity Risk: A Prospective
495 Cohort Study in Rural Western China. *Medicine*, 95(16), e3425.
496 doi:10.1097/md.0000000000003425
497