## An investigation into the effects of E-cigarette aerosols using a physiologically relevant *in-vitro* model

Pranav Vasanthi Bathrinarayanan<sup>1</sup>, Lindsay J. Marshall<sup>2</sup>, James E.P. Brown<sup>2</sup>, Laura J. Leslie<sup>1</sup> 1-School of Engineering and Applied Science, Aston University, Birmingham, B4 7ET, UK 2-School of Life and Health Sciences, Aston University, Birmingham, B4 7ET, UK

Although widely popular as smoking cessation tools, limited information is available on the health effects of E-cigarette (EC) use. An urgent requirement of EC research is a standard testing method to investigate possible adverse effects. The present study aims to assess EC cytotoxicity using an *in-vitro* multicellular human airways model.

Human bronchial epithelial cells (CALU3) and pulmonary fibroblasts (MRC5) were co-cultured on permeable membranes for 11–14 days at air–liquid interface. A bespoke smoking machine was used to deliver air, whole cigarette smoke (WCS) or EC vapour (ECV) to the airways model under standard ISO:3308 conditions for 7 m. Considering the prolonged vaping habits of EC users compared to cigarette smoking, ECV exposure was additionally investigated at 1 h, 2 h, 3 h, 4.5 h and 6 h time points.

24 h post exposure, XTT cell viability analysis showed that while WCS had the expected detrimental impact on cell viability, air exposure had no effect at any time point. Interestingly, a steady decrease in the viability of ECV exposed cells was observed at times greater than 2 h. Viability was  $61.31 \pm 5.75\%$  control,  $51.11 \pm 5.56\%$  control and  $42.10 \pm 2.69\%$  control after 3 h, 4.5 h and 6 h respectively. Furthermore, ELISA analysis of supernatants revealed an increase in IL-6/IL-8 pro-inflammatory cytokines at 3 h post ECV exposure, despite the increased cell death.

Results indicate that extended EC exposure ( $\geq$ 3 h) under these conditions has a detrimental impact on cell viability and leads to exaggerated cytokine production in the airways model.