

Impact of mechanical stretch on the cell behaviors of bone and surrounding tissues

Journal of Tissue Engineering
Volume 7: 1–24
© The Author(s) 2016
Reprints and permissions:
sagepub.co.uk/journalsPermissions.nav
DOI: 10.1177/2041731415618342
tej.sagepub.com



Hye-Sun Yu^{1,2,3}, Jung-Ju Kim^{2,3}, Hae-Won Kim^{2,3,4},
Mark P Lewis⁵ and Ivan Wall^{1,2}

Abstract

Mechanical loading is recognized to play an important role in regulating the behaviors of cells in bone and surrounding tissues *in vivo*. Many *in vitro* studies have been conducted to determine the effects of mechanical loading on individual cell types of the tissues. In this review, we focus specifically on the use of the Flexercell system as a tool for studying cellular responses to mechanical stretch. We assess the literature describing the impact of mechanical stretch on different cell types from bone, muscle, tendon, ligament, and cartilage, describing individual cell phenotype responses. In addition, we review evidence regarding the mechanotransduction pathways that are activated to potentiate these phenotype responses in different cell populations.

Keywords

Flexercell, mechanical strain, bone, cartilage, muscle, tendon, ligament

Received: 10 September 2015; accepted: 15 October 2015

Introduction

The bone and surrounding tissues experience many different types of mechanical loading. For example, compressive forces are experienced and tolerated well by articular cartilage. Shear stress resulting from interstitial fluid flow is tolerated in bone tissue and in tendons and ligaments, tensile stretch is tolerated in order to maintain skeletal anatomy. The effect of different forms of mechanical loading of cells has been extensively studied *in vitro* and in animal models and human subjects, where it is commonly reported to have stimulatory effects on those cells. In fact, fluid flow-induced shear stress actively enhances cellular events that potentiate bone formation and remodeling,^{1,2} and compressive loading of articular chondrocytes stimulates their proliferation capacity. In this review, we will focus specifically on the impact of mechanical stretch on bone and surrounding cell responses and in particular, the use of the Flexercell system as a means of applying that stretch.

In 1985, the first reported use of flexible-bottomed plates for studying the effect of mechanical stimulation on cell cultures was made.³ The plates used a computer-controlled vacuum device to apply negative pressure to the underside of the membrane. When the vacuum was applied,

the membrane on which firmly adhered cells were attached was stretched downward and consequently, a strain was imparted on the cells adhered to that membrane. A number of parameters under the control of the operator included the

¹Department of Biochemical Engineering, University College London, London, UK

²Department of Nanobiomedical Science and BK21 Plus NBM Global Research Center for Regenerative Medicine, Dankook University Graduate School, Cheonan, South Korea

³Institute of Tissue Regeneration Engineering (ITREN), Dankook University, Cheonan, South Korea

⁴Department of Biomaterials Science, School of Dentistry, Dankook University, Cheonan, South Korea

⁵Musculo-Skeletal Biology Research Group, School of Sport, Exercise and Health Sciences, Loughborough University, Loughborough, UK

Corresponding authors:

Ivan Wall, Department of Biochemical Engineering, University College London, Torrington Place, London WC1E 7JE, UK.
Email: i.wall@ucl.ac.uk

Hae-Won Kim, Institute of Tissue Regeneration Engineering (ITREN), Dankook University, Cheonan 330-714, South Korea.
Email: kimhw@dku.edu






Figure 1. Experimental setup of the Flexercell system for applying strain to cell populations. (a) Schematic representation of the computer-controlled Flexercell system that applies strain to cell monolayers in custom-made six-well plates. (b) Cross-sectional diagram of one well of a six-well Flexercell plate at rest (top) and upon application of the vacuum (bottom). (c) Strain can be applied in a variety of ways including uniaxial and equiaxial.

magnitude of pressure, waveform, frequency, duration, and incorporation of rest period over the time course of the experiment. As the system is easy to set up and these various parameters are easy to control, the Flexercell system has been a very good platform with which to study the effect of mechanostimulation in many different contexts (Figure 1).

One huge benefit of the widespread use of this system is that even though the many studies of bone and surrounding cell responses use different strain regimes (magnitude, frequency, duration, etc.), the fact that the base technology applies strain using a common method of application means that some comparability between different studies can be made. Strain is applied either uniaxially or biaxially, generally in two-dimensional (2D) culture (although there are some exceptions to this, where a three-dimensional (3D) culture module is used to characterize compressive force) on silicone membranes coated with type I collagen. Dishes can also be coated with other extracellular matrix (ECM) molecules too making it possible to characterize the effect of mechanical stretch on cells anchored to different ECM components, providing a 2D model of *in vivo* mechanical stimulation of tissues.

In the bone and surrounding tissues, mechanical signals are continually generated and experienced by the numerous cell types that make up the component tissues. Understanding the role of mechanotransduction and its effects at the cellular level can greatly enhance our understanding of the role of mechanical signals in maintaining normal healthy musculoskeletal physiology and also provide some insight to mechanisms of pathogenesis, for

instance, during chronic overuse injury. In the following sections, we review the ever-growing body of the literature that describes bone and surrounding cell responses to mechanical strain applied using the Flexercell system.

Effect of mechanical strain on bone cell phenotypes

During bone loading *in vivo*, macroscale forces are transmitted to osteocytes through the canalicular network in the form of microscale shear stresses caused by rapid displacement of interstitial fluid. Prolonged cyclic loading of bone leads to increased mineralized tissue density and bone mass. It was hypothesized some time ago that mechanical force was a positive regulator of bone formation creating a shift in bone homeostasis toward the anabolic state, characterized by enhanced bone-forming activity by osteoblasts and a reduction in bone resorption by osteoclasts.⁴ In addition to shear stress caused by fluid flow (see above; reviewed by Brindley et al.¹ and Yeatts and Fisher²), tensile stretch applied to cells via physical deformation of culture surfaces^{5–7} can enhance key cellular and molecular events that lead to bone formation. A summary of notable studies is presented in Table 1.

Impact on cell survival, proliferation, and growth

Tensile strain applied using the Flexercell increases osteoblast proliferation^{5,9,20,28} although the magnitude of strain appears to have a bearing on the degree of proliferation

Table 1. Summary of Flexercell studies and key findings using osteoblastic cells.

Cell type	Stage of differentiation	Stretch device and regime	Key findings	Reference
Murine osteocytic (MLO-Y ₄)	Osteocytes	Flexercell 4000, 5% elongation, 10 min, 0.05–1 Hz	Estrogen receptor involved in strain-mediated pro-survival signaling via ERK	Aguirre et al. ⁸
Human osteoblasts	24-h culture in differentiation medium (early differentiation)	Flexercell, ^a 3%–9% strain, 8 h, 1 Hz	High strain increased proliferation, migration, VEGF and bFGF production Low strain increased OCN and OPN	Bhatt et al. ⁹
Rat osteoblast (ROS 17/2.8) and mouse osteoblast (MC3T3-E1)	Early osteoblasts	Flexercell 3000, 1% elongation, 10 min, 0.25 Hz	Strain-induced rapid phosphorylation of ERK2 and FAK and activation of Ras/Raf/MEK pathway	Boutahar et al. ¹⁰
Human osteosarcoma (TE-85)	Osteosarcoma	Flexercell, ^a intermittent strain of 3 cycles/min consisting of 10 s strain, 10 s rest, 15–60 min or 1–3 h	Strain-induced reorganized distribution of integrins and increased $\beta 1$ but not αv mRNA levels	Carvalho et al. ¹¹
Human osteosarcoma (SaOS-2)	Osteosarcoma	Flexercell, ^a 5%–12% elongation, 8–24 h, 0.05 Hz	Increased expression of TGF β 1, IGFI, bFGF, IL6, but no change in IL1	Cillo et al. ¹²
Mouse osteoblast (CIMC-4)	Pre-osteoblast	Flexercell, ^a biaxial strain at 0.5%–2%, 24 h, 0.16 Hz	Strain caused a reduction in RANKL, but upregulated OSX and RUNX2 via ERK1/2	Fan et al. ¹³
Rat osteoblast (ROS17/2.8) and human osteoblasts	Single cells; 24-h post-plating	Flexercell 3000, 1% elongation, 1.5–150 min, 0.05–5 Hz	Expression of soluble VEGF isoforms (121, 165) under low frequency Expression of matrix-bound VEGF isoforms (206, 189, 165, 145) under high frequency	Faure et al. ¹⁴
Rat osteoblast (ROS17/2.8)	Mature osteoblasts (3 weeks in differentiation medium)	Flexercell 3000, 1% elongation, 10 min, 0.05 Hz	Stretch-induced activation of Egr-1 and nuclear translocation of NF- κ B	Granet et al. ¹⁵
Rat osteoblast (ROS 17/2.8) and mouse osteoblast (MC3T3-E1)	Early osteoblast	Flexercell 3000, 1% elongation, 15 cycles/min (2-s deformation period followed by a 2-s neutral position)	Induced phosphorylation of FAK, PYK2, paxillin and HIC5	Guignandon et al. ¹⁶
Murine calvarial osteoblasts	Early osteoblast (7–10 days post-plating)	Flexercell 4000, 2.5%–3% elongation, 3–24 h, 0.3 Hz	Stretch resulted in the activation of canonical Wnt signaling	Hens et al. ¹⁷
Human fetal osteoblast (SV-HFO)	Differentiation medium	Bioflex loading stations, 0.4%, 5–60 min or 7–21 days, 0.05 Hz	Stretch-induced phosphorylation of ERK1/2 pathways dependent on differentiation stage	Jansen et al. ¹⁸
Mouse osteoblast (MC3T3-E1)	24 h in differentiation medium	Flexercell 4000, 0.25%–2.5%, 1 Hz, 15 min/day for 7 days	Overall increased RANKL due to increased membrane-bound RANKL but decreased soluble RANKL	Kim et al. ¹⁹
Rat osteoblast and mouse osteoblast (MC3T3-E1)	Osteoblasts	Flexercell, ^a 10%–12%, 0.1 Hz either one single maximal strain load every 6 h or continuous strain for 24 h	Stretch increased DNA synthesis ALP decreased in response to strain and combined strain/TGF β treatment	Knoll et al. ²⁰
Human osteoblast (HOB)	Osteoblasts	Flexercell 4000, 2.5% elongation, 0.1 Hz, intermittent strain 1 h, strain 3 h, and rest for 48 h	Increased DNA synthesis and IL6 production No effect on ALP, COL I or OPG production	Liegibel et al. ⁵
Human osteosarcoma (SaOS-2)	7 days post-plating	Flexercell, ^a 5%–12.5% strain, 24 h, 0.5 Hz	Low strain led to increased COL I and COL III expression High strain led to decreased COL III expression	Liu et al. ²¹

(Continued)

Table 1. (Continued)

Cell type	Stage of differentiation	Stretch device and regime	Key findings	Reference
Mouse osteoblast (MC3T3-E1)	Osteoblasts	Flexercell 2000, 7%–24% elongation, 3 cycles/min, 1–24 h	Increased expression of VEGF and M-CSF	Motokawa et al. ²²
Mouse osteoblast (MC3T3-E1)	Osteoblasts	Flexercell 4000, 0%–9% elongation, 3–24 h, 0.3 Hz	Stretch promoted COX2 but not COX1 expression	Narutomi et al. ²³
Murine osteocytic (MLO-Y ₄)	Osteocytes	Flexercell 4000, 2%–5%, 1–20 min	Activation of ERK via integrin/cytoskeleton/Src-mediated signaling	Plotkin et al. ²⁴
Mouse osteoblast (MC3T3-E1)	Osteoblasts	Flexercell 3000, 3400 microstrain, 2 Hz, 5 h	Increased Wnt/B-catenin target genes (Wnt10B, SFRP1, cyclin D1, FZD2, WISP2, and connexin 43)	Robinson et al. ²⁵
Mouse osteoblast (MC3T3-E1)	Osteoblasts	Flexercell 3000, 6%–18% elongation, 24 h, 0.1 Hz	Increased OPG synthesis and decreased RANKL expression	Tang et al. ⁶
Human osteosarcoma (MG63)	Osteoblasts	Flexercell 3000, 10% elongation, 14 h, 0.5 Hz	Strain inhibited the promoter activation by vitamin D	Toyoshita et al. ²⁶
Rat osteoblast	Differentiation medium	Flexercell, ^a –2 kPa, 1–15 days, 0.05 Hz	Nodule formation was enhanced, depending on the timing of initiation and magnitude of the deformation regimen	Visconti et al. ²⁷
SV40 human osteoblasts	Differentiation medium	Flexercell 4000, 0.8%–3.2% elongation, 48 h, 1 Hz	High-magnitude strain led to increased expression of OC, COL I and Cbfa1/Runx2 Low-magnitude strain led to increased ALP activity	Zhu et al. ⁷

ERK: extracellular-regulated protein kinase; VEGF: vascular endothelial growth factor; bFGF: basic fibroblast growth factor; OCN: osteocalcin; OPN: osteopontin; FAK: focal adhesion kinase; NF- κ B: nuclear factor- κ B; TGF β 1: transforming growth factor β 1; COL: collagen; IGF1: insulin-like growth factor 1; IL: interleukin; PYK2: proline-rich tyrosine kinase 2; COX: cyclooxygenase; M-CSF: macrophage-colony stimulating factor; ALP: alkaline phosphatase; OPG: osteoprotegerin; MEK: mitogen-activated protein kinase (MAPK)-extracellular signal-regulated kinase (ERK) kinase; RANKL: receptor activator of NF- κ B ligand; OSX: osterix; RUNX2: Runt-related transcription factor 2; Egr-1: Early growth response protein 1; HIC5: hydrogen peroxide-inducible clone-5.

^aFlexercell model not specified.

that is permissible.⁹ Application of a high strain level of 10%–12% elongation at 0.1 Hz for 24 h stimulated proliferation in primary rat osteoblasts but the even greater mitogenic effect that transforming growth factor β (TGF β) has in the absence of strain was suppressed by strain itself,²⁰ suggesting that mechanotransduction competes with TGF β signaling to dampen its effects. Application of mechanical strain to osteocytes and very mature osteoblasts failed to stimulate proliferation,²⁹ and it was hence concluded that these cells respond differently to proliferative osteoblasts, instead regulating metabolic responses to strain reflective of their mechanosensing role in vivo.

Lower levels of strain (2.5% elongation at 0.1 Hz) were stimulatory to proliferation of human primary osteoblasts over 48 h, without promoting alkaline phosphatase (ALP) activity or secretion of type I collagen or osteoprotegerin (OPG), suggesting that the cells were maintained in a premature state. However, pre-treating cells with testosterone prior to application of strain promoted osteogenic gene expression at the expense of proliferation,⁵ highlighting that the effects of mechanotransduction are subject to a mechanistic switch regulated by androgens.

A study assaying the effects of strain over a range of 0%–9% elongation for 8 h at 1 Hz revealed that proliferation

of normal human osteoblasts was magnitude dependent, with the highest strain producing the greatest level of proliferation.⁹ Conditioned medium obtained from cultures stretched to 9% elongation stimulated cell proliferation in static culture, indicating the release of a soluble factor.

In summary, the collated evidence indicates that stretch signals have a positive effect on osteoblast proliferation and they do so in a magnitude-dependent fashion. However, there will inevitably be a critical upper threshold at which stretch signals are no longer stimulatory to proliferation and are indeed detrimental to cell viability. Furthermore, biological molecules such as growth factors and hormones may alter the basal cell response to stretch, and this is particularly significant in vivo, where mechanical signals are just one of many facets of the cell microenvironment.

Matrix production

When a range of stretch magnitudes from 0% to 9% was applied to human primary osteoblasts, expression of the type I collagen mRNA, COL1 α II, was highest at 9% elongation, whereas in the case of mature osteoblast markers osteopontin (OPN) and osteocalcin (OCN), highest expression was seen at 3% strain, declining thereafter,⁹ indicating

a biphasic response to tensile strain. This parallels the authors' observations for proliferation, which was superior at the highest magnitude of 9%. Furthermore, COL1 expression by human primary osteoblasts was only upregulated at a low magnitude of 2.5% strain in the presence of testosterone,⁵ further evidence that the effect of mechanical stimulation is subject to hormonal influence. For type III collagen expression, where a "high impact" strain regime of between 5% and 12.5% strain at 0.5 Hz for 24 h on human osteosarcoma cells was used, Liu et al.²¹ found that COL3 mRNA was increased at 5% but then underwent a decline in the expression after 5%, whereas COL1 transcripts were stably increased at all strain levels compared to static control cultures. As type III collagen is normally deposited as an early wound matrix component, it is surprising that it was not increased at the higher strain level.

In summary, early matrix production events, that is, deposition of type I collagenous material, appear to preferentially take place at higher strain levels in parallel with the enhanced proliferation effect. This may reflect the wound healing environment *in vivo* whereby early events of osteoblast proliferation and matrix deposition take place in a mechanically unstable environment prior to the formation of rigid bone, placing large amounts of strain on cells.

Impact on cytokine/growth factor production

Flexercell studies have revealed that numerous growth factors are upregulated in osteoblasts in response to stretch signals. Over a range of stretch from 0% to 9%, vascular endothelial growth factor (VEGF) expression by primary human osteoblasts was highest at 9% stretch.⁹ It also seems that the frequency with which stretch is applied influences the form of VEGF synthesized, as 1% stretch applied to both human and rat osteoblasts at various frequencies from 0.05 to 5 Hz revealed that high-frequency stretch promoted the expression of VEGF isoforms that are matrix bound and low frequencies stimulated production of soluble VEGF isoforms.¹⁴ Upregulation of VEGF, along with macrophage-colony stimulating factor (M-CSF) have been reported to be mechanistically important for the recruitment of osteoclasts in the remodeling of bone,²² and so stretch-induced VEGF expression is pivotal for the regulation of bone remodeling events *in vivo* as well as in revascularization of regenerating tissue.

TGF β expression in human osteosarcoma cells was upregulated in response to mechanical stretch applied at 12% elongation.¹² TGF β is a potent mitogen and can also promote collagen synthesis, both of which are important during the early stages of bone healing. However, binding of TGF β to the TGF β receptor-III is also enhanced by mechanical stretch,²⁰ and this interaction has been shown to inhibit the ability of TGF β to activate osteoblasts.^{30–32} Therefore, mechanical strain seems to finely regulate TGF β signaling and promote similar mitogenic

and differentiation responses to TGF β but via a mechanotransduction pathway that competitively inhibits TGF β signaling.

Mechanical stretch also enhanced osteoblastic expression of insulin-like growth factor (IGF) and basic fibroblast growth factor (bFGF) at 12% maximal stretch at 3 Hz, over a 24-h period.¹² Both factors stimulate early osteoblast healing responses such as osteoblast proliferation and matrix synthesis^{33,34} and the early wound healing events induced in osteoblasts by high-magnitude stretch, such as proliferation and type I collagen expression (see above), may be a consequence of increased expression of these factors. Indeed, osteocytes, which act as mechanosensing cell *in vivo*, produce increasing levels of IGF in response to mechanical strain^{35–37} which may be one of the primary initiation events for the recruitment of osteoblast precursors during bone remodeling and regeneration.

Increased expression of several bone morphogenetic proteins (BMPs), in particular BMP2, in response to low strain of 1% stretch was also reported in mouse osteoblastic MC3T3-E1 cells,³⁸ yet in human primary osteoblasts subjected to stretch over a range of 0%–9% elongation, increased gene expression was only seen at 6% and above.⁹ However, the overall positive effect of mechanical stretch on osteoblastic expression of BMP2 is further evidence of the stimulatory effect of mechanical stretch signals on osteogenesis.

In terms of the bone remodeling balance, mechanical stretch also seems to have a role in regulation of bone resorption by osteoclasts. Interleukin 6 (IL6) has a well-known role in osteoclast production during normal bone homeostasis.³⁹ Increased expression of IL6 was observed in human osteosarcoma cells stretched at 12% elongation at 0.05 Hz¹² and in human primary osteoblasts stretched at 2.5% elongation at 0.1 Hz.⁵ In the latter study, OPG, an inhibitor of osteoclastogenesis, was produced when stretch was applied in combination with testosterone, but not in its absence, demonstrating the coordinated action of mechanotransduction and hormones. The effects of stretch on IL6 are magnitude dependent, as studies of mouse MC3T3-E1 osteoblastic cells exposed to a regime of 0%–9% elongation revealed that the greatest level of IL6 production was seen at 9% stretch and was dependent on prostaglandin E2 production by cyclooxygenase 2.²³

A key downstream consequence of IL6 activity in osteoblasts that leads to osteoclast activation is production of RANKL.³⁹ There is some debate in the literature about the effect of mechanical strain on RANKL expression. Some researchers have reported a magnitude-dependent decline in RANKL expression,^{6,13} and this was accompanied by an increase in the expression of the transcription factors Osterix and Runx2¹³ and OPG.⁶ However, Kim et al.¹⁹ reported an increase in RANKL expression after exposure to mechanical strain. Nevertheless, in this study, a potential mechanism for preventing osteoclast activity in order

to maintain an osteoanabolic process was discovered using MC3T3-E1 cells transfected with RANKL cDNA. The expression of RANKL increased in transfected cells, but when low-level mechanical strain was applied daily for 1 week, most of the protein was in membrane-bound form with a large reduction in soluble RANKL, thus reducing its bioavailability.¹⁹ In any case, RAW264.7 cells that are capable of osteoclastic differentiation in the presence of RANKL failed to do so effectively in the presence of mechanical strain of 10% elongation at 0.5 Hz over 1 week.⁴⁰ Together with the strong hormonal influence as demonstrated for testosterone,⁵ osteoclast activity is subject to further regulation upon exposure to strain.

Overall, mechanical stretch signals induce the expression of factors that invoke early wound healing events such as mesenchymal cell proliferation and angiogenesis. Furthermore, stretch-induced upregulation of factors such as VEGF and IL6 results in osteoclast recruitment and differentiation, which is in turn carefully regulated by androgens. Studies that have assayed growth factor production over a range of different elongations generally concur that growth factor production is magnitude dependent. This likely reflects the early wound healing response where the fracture site is highly unstable and growth factors are in abundance.

Cellular maturation and differentiation

Osteoblastic differentiation, maturation, and ultimately mineralized matrix deposition can be directly enhanced by mechanical strain applied using Flexercell. For example, stretch applied to immortalized mouse calvarial osteoblasts overnight at 2% elongation enhanced the expression of osteogenic transcription factors, osterix and runx2.¹³ Over a range of elongations from 0% to 3.2%, a subsequent study found that runx2 was significantly upregulated only at 3.2% stretch but not at 2.4% or lower, and type I collagen expression was upregulated in proportion to magnitude of stretch.⁷ In this study, OCN was only upregulated at 2.4%–3.2% elongation, whereas ALP (expressed earlier in osteogenic differentiation) was upregulated at 0.8%–1.6% elongation but not above.

The strain magnitude-dependent expression of OCN by human primary osteoblasts was further confirmed using an elongation range of 0%–9%. OCN expression was upregulated at 3%, 6%, and 9% elongation compared to 0% control cultures,⁹ whereas in the case of osteonectin and OPN, greatest gene expression was seen at 3% and then underwent a gradual decline in basal levels by 9%.⁹ At stretch magnitude as low as 1% elongation, bone nodule formation was induced in monolayers of mouse MC3T3-E1 and rat primary osteoblasts.^{27,41} Particularly intriguing is the fact that some researchers reported that early osteogenic events such as proliferation and collagen production were best at a high strain of 9%, whereas the

expression of differentiation markers OPN and OCN was most favorable at low strain of 3% elongation,⁹ and bone nodule formation was reported as low as 1% elongation.^{27,41} Other researchers reported that the osteoanabolic phenotype was enhanced in osteoblasts with increasing magnitude of strain up to 18% elongation, characterized by enhanced expression of OPG,⁶ which is a potent inhibitor of osteoclast formation and activity.⁴² However, it is not just the magnitude of strain that impacts the specific response of osteoblasts but the timing of application and duration of strain that determine the exact nature of cell responses. According to the work of Visconti et al.,²⁷ applying strain after a prolonged static culture period (either 3 or 10 days) produced a greater number of bone nodules than applying strain after just 1 day of static culture. This suggests that a regime that promotes mineralized matrix formation should be applied after early events such as collagen matrix deposition have been given time to occur.

To summarize the impact of mechanical stretch on osteogenic differentiation and maturation, it is clear that lower magnitude strain in the region of <5% elongation is superior for directing these events, a contrast to early events such as proliferation and type I collagen production, which are optimally enhanced at greater magnitudes of stretch (around 9% elongation).

Effect of mechanical strain on skeletal muscle cells

Mechanical strain is transmitted through tendons and ligaments during locomotion, when muscle contraction occurs. Skeletal muscle itself generates contractile forces that are propagated through the muscle fiber network. The nature of force generation in skeletal muscle means that as well as contraction, cells experience stretch stimuli.

In order to fully critically appraise the information in this area, a brief description of how skeletal muscle differentiates and regenerates is appropriate. The tissue itself is characterized by multinucleate muscle fibers encapsulated by connective tissue, mainly collagen, fixed at both ends (junctions).

In terms of mechano-responsiveness, it could be suggested that skeletal muscle is the most relevant tissue in the human body. It is composed of elongated multinucleate muscle fibers that develop from the “end-to-end” fusing of mononuclear muscle precursor cells (MPCs) in a process known as differentiation. In development, these mononuclear cells are derived from the somites and, depending upon the anatomical location, the MPCs either migrate to differentiate at their final destination, for example, limb muscle or differentiate at or near their final destination, for example, head and neck muscles. However and astoundingly, skeletal muscle is not a solely post-mitotic tissue. MPCs located in the periphery of post-natal muscle fibers




Figure 2. A series of processes from MPCs to differentiated myotubes.

can become activated by a variety of stimuli including mechanical. This activation triggers a series of events that lead to the asymmetric division of these cells. This process can be recapitulated with *in vitro* cell culture and is illustrated in Figure 2.

It is therefore very important to understand at which part of the process the cells are being stimulated using the Flexercell apparatus where a large number of studies have demonstrated that mechanical strain activates muscle cells at different stages of maturity from skeletal myoblasts to satellite cells,^{43–47} and a summary of muscle cell studies is presented in Table 2.

Impact on cell survival, proliferation, and growth

In the vast majority of studies, it is striking that cyclic stretching of single MPCs upregulates both proliferation and expression of genes and proteins that are required for further differentiation: MYOD1, myogenin (MYOG), MEF2A, p21 (CDKN1A), and IGF1.⁴⁷ Furthermore, several genes encoding muscle structural proteins are upregulated, including myosin heavy chain (MyHC) isoforms and α -tropomyosin (TPM1), and this ultimately leads to increased myotube production.⁴⁶ Intriguingly, differences in stretch-induced satellite cell responses have been reported from muscles isolated from distinct anatomical regions that contain primarily fast-twitch or slow-twitch fibers.⁵⁰ Further evidence of stretch favorably enhancing cell growth, tissue building, and regeneration includes an increase in total protein levels in response to mechanical stretch,⁶³ and the whole process of stretch-induced cell growth is dependent on mammalian target of rapamycin (mTOR) activation.⁷² However, an antagonist, REDD2, is also expressed in skeletal muscle and is capable of inhibiting stretch-activated mTOR activity.⁴⁹ Nitric oxide synthase (NOS) becomes upregulated⁴⁶ and bFGF expression is increased, also in a strain-dependent fashion.⁶⁰ NOS drives the synthesis and release of nitric oxide (NO) from mature muscle myotubes.⁵⁶ Together with FGF, NO has well-known mitogenic potential and initiates a “wound healing” response. It does this, at least in part, by activation

of satellite cells via matrix metalloproteinase 2 (MMP2)-mediated release of hepatocyte growth factor (HGF) from ECM⁵¹ and subsequently leads to HGF signaling.^{43–45,73,74} It is particularly noteworthy that single MPCs are very tolerant of a wide range of strain with values ranging from 2% to 25% eliciting similar (sometimes with a dose response) effect.

Impact on the differentiation process and differentiated cells (myotubes)

As MPCs are “preparing” to fuse into multinucleate myotubes, their sensitivity to strain appears to increase⁷⁵ although the global responses remain fairly similar—including increased proliferation rates and increased indicators of fusion (Table 2).

In stark contrast to the data seen with single cells, MPCs fused into myotubes respond in quite different ways. One of the most striking observations is that “tolerance” to varying strain rates is lost. Rates nearing in excess of 20% elicited a variety of deleterious effects such as increased creatine kinase (CK) and lactate dehydrogenase (LDH) activity,^{60,68,70,71} membrane wounding and cell “injury,”^{60,71,76} inflammatory cytokine and wound repair growth factor release,^{60,76} sarcolemmal disruption,⁶² increases in molecular chaperonins,⁶³ and increased acetylcholinesterase production.⁶⁶ Overall, stretching myotubes appeared to have no real effect on parameters related to cell proliferation and differentiation; however, there is some evidence of changes to cellular metabolism such as increased glucose uptake⁶⁷ and phosphorylation of a number of anabolic targets such as ribosomal S6 kinase P70S6K^{59,72}, focal adhesion kinase (FAK), protein kinase B (Akt), 4E binding protein 1 (4EBP1), eukaryotic elongation factor 2 (eEF2), extracellular-regulated protein kinase 1/2 (ERK1/2), eukaryotic initiation factor 2 α (eIF2 α), and eukaryotic initiation factor 4E (eIF4).⁵⁹ Notably, however, myofibrillar protein synthesis was actually suppressed.⁵⁹

In conclusion, there is a good deal of data that have been collected, and the system has, for skeletal muscle, some utility for single MPCs work concentrating on some

Table 2. Summary of Flexercell studies and key findings using muscle cells.

Cell type	Stage of differentiation	Stretch device and regime	Key findings	Reference
<i>Proliferating MPCs</i>				
Murine myoblast (C2C12)	Confluent/early fusion	Flexercell 4000, cyclical stretch at 3% at 0.05 Hz for 24–72 h	Increased MyoD, MyoG, Mef2, MHC Stretch abrogated the reductions in the expression of the above by TNF α treatment	Bruce et al. ⁴⁸
Murine myoblast (C2C12)	Single cells; 24-h post-plating	Flexercell, ^a 15% strain, 0.5 Hz, cyclical for up to 48 h	Increased proliferation Increased IGFI and caspase mRNA expression in the initial 24 h of stretch	Iwanuma et al. ⁴⁷
Murine myoblast (C2C12)	MDC	Flexercell 4000, 15% strain for 10 min	Overexpression of REDD2 inhibits response of mTOR to mechanical stimulation	Miyazaki and Esser ⁴⁹
Rat MDCs	>95% desmin-positive single MDCs	Flexercell 2000, 12–36 h post-plating at 25% strain for 12–36 h with 12-s intervals	Increase in BrdU+ cells after 12-h stretch Increase in HGF in CM at 12 h. Anti-HGF antibody prevents increase in BrdU+ cells. Stretch releases pre-existing HGF from ECM	Tatsumi et al. ⁴³
Rat MDCs	>95% Desmin positive single MDCs	Flexercell 2000, 12–36 h post-plating at 25% strain for 24 h with 12 s intervals	Increase in LH limb BrdU+ cells compared to BK and UH cultures following stretch. Increase in HGF in CM of LH cultures stretch compared to BK and UH	Tatsumi et al. ⁵⁰
Rat MDCs	>95% desmin-positive single MDCs	Flexercell 2000, 12–36 h post-plating at 25% strain for 24 h with 12-s intervals	Increase in BrdU+ cells at an optimal pH (7.2) Increase in BrdU+ cells in a HGF- and NO-dependent mechanism	Tatsumi et al. ⁴⁴
Rat MDCs	Single MDCs	Flexercell 2000, 12–36 h post-plating at 25% strain for 24 h with 12-s intervals	Stretch activation of BrdU+ cells is NO dependent	Tatsumi et al. ⁴⁵
Rat MDCs	Single MDCs	Flexercell 2000, 25% strain, 12-s intervals for 2 h	Stretch induced an increase in the active form of MMP2 in a NO-dependent mechanism	Yamada et al. ⁵¹
<i>Aligning MPCs</i>				
Murine myoblast (C2C12) and murine MDCs	Single cells induced to differentiate; 24-h post-seeding (C2C12) or immediately post-seeding (MDCs)	Flexercell 4000, 48 h uniaxial ramp stretch, followed by intermittent strain at 2%–6% strain for 4 days. Cyclic stretch at 1 Hz	2D: reductions in MRF-4 and MYH-4 mRNA in both cell types. Increase in MYH-1 mRNA (C2C12 only) 3D: reductions in MRFs, MyoD, myogenin (both cell types) MRF-4 (C2C12) and MLP (MDCs). Increases in actin and α -actinin mRNA at day 4 (C2C12), reduction in MYH mRNAs (both cell types)	Boonen et al. ⁵²
Murine myoblast (C2C12)	Single cells induced to differentiate	Flexercell 4000, 17% cyclical stretch, 1 Hz for 1 h every 24 h for 5 days	Stretch increased proliferation and inhibited differentiation	Kumar et al. ⁵³
Murine myoblast (C2C12)	Single cells induced to differentiate	Flexercell, ^a 15% cyclical strain for up to 5 days at 0.1, 0.5, and 0.9 Hz	Stretch improved proliferation	Kurokawa et al. ⁵⁴

Table 2. (Continued)

Cell type	Stage of differentiation	Stretch device and regime	Key findings	Reference
			Stretch led to increased MHC-perinatal expression Fast stretch led to early expression, slow led to later expression	
Rat myoblasts (L6)	Single cells induced to differentiate	Flexercell 4000, 20% cyclical strain at 0.5 Hz for up to 24 h	Stretch caused caspase-induced apoptosis during differentiation	Liu et al. ⁵⁵
Murine MDCs isolated from WT and <i>mdx</i> tissue	24 h in differentiation medium (early differentiation)	Flexercell, ^a 10% strain for 2 s, followed by 20–60 s in the non-stretched position	Increase in real time measurement of NO production during stretch (WT). No observed increase in NO in <i>mdx</i> MDCs	Wozniak and Anderson ⁵⁶
PI–P2 rat MDCs	Confluent single cells	Flexercell, ^a 25% strain at 6 cycles/min, each cycle consisting of 3-s strain, 3-s rest. Continuous for 24 h	Stretch induced an increase in Na ⁺ pump activity, increased expression of α -2 subunit of Na ⁺ -K ⁺ -ATPase in the membrane fractions. Mediated through PI3-K	Yuan et al. ⁵⁷
Murine myoblast (C2C12)	Single cells induced to differentiate	Flexercell, ^a 10% strain at 0.5 Hz for 1 h on, 5 h off-0-4 days	Increase in actin fiber formation, myotube maturation (increase in α -actinin and striations) and myotube diameter with stretch. β _{ID} and FAK proposed mediated mechanism	Zhang et al. ⁵⁸
<i>Myotubes</i>				
Rat myoblasts (L6)	Myotubes (7–9 days in culture)	Flexercell 4000, 1 Hz at 15% strain at 2, 15, 30, 60, 90, 120, and 150 min	Sarcoplasmic protein synthesis was unaltered; however, myofibrillar protein synthesis was decreased. Paradoxically, anabolic signaling (phosphorylation of key proteins) was enhanced	Atherton et al. ⁵⁹
Human MDCs	Stretched during differentiation or 10-min stretch after 7-day differentiation	Flexercell 2000, cyclical strain during differentiation 10%–20% cyclical strain	Stretching during differentiation increase CK activity Acute stretching caused membrane wounding and FGF release	Clarke and Feedback ⁶⁰
Mouse diaphragm MDCs	Myotubes (5 days in DM)	Flexercell, ^a 4 h, 10% stretch, 1.5 Hz	No effect of stretch on proinflammatory marker gene expression	Demoule et al. ⁶¹
Rat myoblasts (L6)	Myotubes (7 days in DM)	Flexercell 3000, 10% stretch, 4 h, 1.5 Hz	Mechanical stretch plus oxidative stress (induced by SIN-1) causes sarcolemmal injury to a greater extent than either component alone	Ebihara et al. ⁶²
Rat myoblasts (L6)	Early myotubes	Flexercell 3000, 96 h of cyclical strain (18% stretch, 0.16 Hz) +/- heat stress	Total protein, HSP72 and 90 protein concentrations were increased by stretch and heat in a cumulative manner	Goto et al. ⁶³
Murine myoblast (C2C12)	Myotubes (5–6 days post-induction of differentiation)	Flexercell 3000, 15% stretch, 1 Hz, 10 min, cyclical	Increased phosphorylation of p70s6k following multiaxial versus uniaxial	Hornberger et al. ⁶⁴
Murine myoblast (C2C12)	Myotubes (5–6 days post-induction of differentiation)	Flexercell 3000, 15% stretch, 1 Hz, 10 min, cyclical	Stretch caused phosphorylation of p70s6k	Hornberger ⁶⁵

(Continued)

Table 2. (Continued)

Cell type	Stage of differentiation	Stretch device and regime	Key findings	Reference
Rat MDCs	Early myotubes (8–10 days post initial plating)	Flexercell 2000, up to 24 h cyclical stretch, 0.25 Hz, 24% strain	Addition of locally acting growth factors to myotubes did not have the same effect 24 h stretching increased Ach esterase activity	Hubatsch and Jasmin ⁶⁶
Rat myoblasts (L6)	Myotubes; 7–8 days post-plating and in DM	Flexercell 100C, 25% cyclical stretch at 0.5 Hz for up to 48 h	Stretch caused increased glucose uptake in myotubes, not myoblasts independent of GLUT1 and 4 receptor content	Mitsumoto and Klip ⁶⁷
Murine myoblast (C2C12)	Early myotubes (24-h DM)	Flexercell, ^a 1 Hz, 20% strain consisting of 20 s on, 20 s off. Continuous for 24 h	Increase LDH into CM. Neutrophil cytotoxicity of muscle cell membrane mediated by MPO	Nguyen et al. ⁶⁸
Human MDCs	Myotubes (5 days in DM)	Flexercell 4000, 0.25 Hz at 1.5, 4.9, 9.5, and 15% strains, for 30 min, 2 s on, 2 s off	Increased injury index and neutrophil chemotaxis with increased strain. Increases in IL8 and MCP-1 in conditioned media	Peterson and Pizza ⁶⁹
Rat myoblasts (L6) and rat MDC-sarcoglycan (SG) deficient	Myotubes	Flexercell, ^a 20% strain at 6 cycles/min for 1 h. Bioflex plates also used for microscopy	Increase in creatine phosphokinase in CM following stretch. Stretch-induced damage in SG-deficient cells caused by alteration in Ca ²⁺ dynamics	Sampaolesi et al. ⁷⁰
Human MDCs	Myotubes (5 days in DM)	Flexercell 4000, 0.25 Hz for 2 h at 5%, 10%, 20%, or 30% strain	Lower myotube injury index at lower strains compared to higher strains and scrape injury (based on LDH in CM). Evidence for membrane rupture and an increase in lanthanum-rimmed membrane blebs (sign of injury) at 30% strain. Increased chemotaxis index of neutrophils using CM at higher strains	Tsivitse et al. ⁷¹

Mef2: mouse embryonic fibroblasts 2; MHC: myosin heavy chain; TNF α : tumor necrosis factor α ; IGF1: insulin-like growth factor 1; MDC: myeloid-derived suppressor cell; mTOR: mammalian target of rapamycin; BrdU: bromodeoxyuridine; HGF: hepatocyte growth factor; CM: conditioned media; ECM: extracellular matrix; NO: nitric oxide; MMP2: matrix metalloproteinase 2; 2D: two-dimensional; MRF: myogenic regulatory factor; 3D: three-dimensional; PI3-K: phosphoinositide 3-kinase; FAK: focal adhesion kinase; CK: creatine kinase; FGF: fibroblast growth factor; GLUT1 and 4: glucose transporters 1 and 4; LDH: lactate dehydrogenase; IL8: interleukin 8; MCP-1: monocyte chemoattractant protein-1; MyoG: myogenin; REDD2: regulated in development and DNA damage responses 2; MYH: myosin, heavy chain; MLP: muscle LIM protein; WT: normal C57BL/6 wild-type; SIN-1: 3-morpholinonydnimine-N-ethylcarbamide; HSP72: heat shock proteins; LH: lower hind; BK: back; UH: upper hind; Ach: acetylcholinesterase; MPO: myeloperoxidase.

^aFlexercell model not specified.

aspects of cellular signaling. There are clearly wider considerations when it comes to interpretation of the myotube data and also around the biomimetic nature of the systems and the interplay with other cell types.⁷⁷ The most powerful use is for comparing the same conditions across varying cell types, as described in this review.

Effect of mechanical strain on tendon and ligament cells

Tendon and ligament are important connective tissues, essential for healthy musculoskeletal physiology. Ligaments bind skeletal components together, maintaining their correct anatomical arrangement. Tendons connect muscles to bone and facilitate the transmission of contractile force from those muscles to control movement. They are therefore

able to withstand high mechanical strain during normal physiological conditions. However, chronic overuse or pathologically high magnitudes of strain result in damage, causing considerable pain and impaired mobility. Using the Flexercell system, the effect of mechanical strain on cellular constituents of tendons and ligaments can be studied, facilitating our understanding of how these tissues respond to, and propagate, mechanical signals (Table 3). This in turn may help us to understand the kind of mechanical stimulation appropriate for engineering replacement tissues.

Impact on cell survival, proliferation, and growth

One of the earliest studies using Flexercell demonstrated that applied cyclic stretch alone was not capable of

Table 3. Summary of Flexercell studies and key findings using ligament cells.

Cell type	Stage of differentiation	Stretch device and regime	Key findings	Reference
Human PDL	Confluent single cells	Flexercell, ^a TENS-L (3%, 6%, 8%), TENS-H (15%), 30–120 min	Suppressed IL1 β -induced COX2 expression TENS-H upregulated COX2 expression and PGE ₂ synthesis	Agarwal et al. ⁷⁸
Human PDL	Single cells	Flexercell, ^a 20 kPa, 12 h	Increased expression of MMP1, MMP2, TIMP1, TIMP2, α 6 and β 1	Bolcato-Bellemin et al. ⁷⁹
Human PDL	Single cells	Flexercell, ^a 15% 30 cycles/min consisting of 1 s strain, 1 s rest	Decreased ALP activity	Chiba and Mitani ⁸⁰
Canine ACL and MCL	Single cells	Flexercell 2000, 5% (0.05 strain) strain at 6 cycles/min for 2 or 22 h daily for 3 days	Increased α 5 and β 1 expression Increased β 3 expression when cells grown on laminin	Hannafin et al. ⁸¹
Canine ACL	Confluent single cells	Flexercell 2000, 5%–15% strain at 6 cycles/min for 2 h followed by 22-h rest for 5 consecutive days	Increased α 5 and β 1 expression	Henshaw et al. ⁸²
Human PDL	Confluent single cells	Flexercell 2000, 9% strain at 6 cycles/min consisting of 5 s strain, 5 s for 6 days	Increased UNCL expression	Kim et al. ⁸³
Human PDL	Single cells	Flexercell, ^a 5% elongation, 3 cycles/min for 24 h on 7 days	Increased TGF β 1 levels Decreased M-CSF expression	Kimoto et al. ⁸⁴
Canine ACL and MCL	Confluent single cells	Flexercell, ^a 60 cycles/min with 0.05–0.075 strain for 0.5–24 h	ACL: Increased COL I MCL: Increased COL III	Hsieh et al. ⁸⁵
Rat MCL	Confluent single cells	Flexercell, ^a 3.5% elongation, 2 h, 1 Hz, microgrooved substrate	Increased sensitivity to load and ability to propagate a calcium wave	Jones et al. ⁸⁶
Human PDL	Osteoblast-like characteristics	Flexercell, ^a 6%–15%, 0.005 Hz, 24 h	Low-magnitude strain inhibited IL1 β -induced synthesis of IL1 I, IL6, and IL8 and induced IL10 synthesis	Long et al. ⁸⁷
Human PDL	Confluent single cells	Flexercell 2000, 9%–18% strain for 6 cycles/min consisting of 5 s strain, 5 s rest, 1–5 days	Increased ALP and OCN expression EGF/EGF-R system acts as a negative regulator of differentiation	Matsuda et al. ⁸⁸
Human LFC	Confluent	Flexercell 2000, 10%–20% strain, 0.16 Hz, 24–48 h	Increased COL I, III, and V via TGF β 1	Nakatani et al. ⁸⁹
Bovine PDL	Confluent	Bioflex loading stations, 0.2%–18% stretch for 6 cycles/min consisting of 5 s strain, 5 s rest, 48 h	High-magnitude strain increased COL I and decorin expression but decreased ALP	Ozaki et al. ⁹⁰
Human PDL	Confluent	Flexercell, ^a 18% stretch for 6 cycles/min consisting of 5 s strain, 5 s rest, 1–5 days	Increased activity of plasminogen activator	Ozawa et al. ⁹¹
Rat PDL (6-week young rats and 60-week old rats)	Confluent	Flexercell, ^a 9%–18% stretch for 6 cycles/min consisting of 5 s strain, 5 s rest, 1–5 days	Increased PGE ₂ and IL1 β production by old versus young in response to strain	Shimizu et al. ⁹²
Human PDL	Confluent	Flexercell, ^a 15% stretch for 30 cycles/min consisting of 1 s strain, 1 s rest, 30–90 min and 6 h	Rapid, transient increase in C-FOS expression No change in expression of osteogenic genes	Yamaguchi et al. ⁹³
Human PDL	Confluent	Flexercell 4000, 12% for 6 s every 90 s, 6, 12, and 24 h	Increased expression of BMP2, BMP6, ALP, SOX9, MSX1, and VEGFA Decreased expression of BMP4 and EGF	Wescott et al. ⁹⁴
Human PDL	Confluent	Flexercell, ^a 7%–21% stretch for 12 cycles/min consisting 2.5 s strain, 2.5 s rest, 24 h	Increased VEGF expression and secretion	Yoshino et al. ⁹⁵

PDL: periodontal ligament; IL: interleukin; COX2: cyclooxygenase 2; PGE₂: prostaglandin E2; MMP: matrix metalloproteinase; TIMP: tissue inhibitors of metalloproteinases; ALP: alkaline phosphatase; ACL: anterior cruciate ligament; MCL: medial collateral ligament; UNCL: ulnar collateral ligament; TGF β 1: transforming growth factor β 1; M-CSF: macrophage-colony stimulating factor; OCN: osteocalcin; EGF: epidermal growth factor; EGF-R: epidermal growth factor receptor; BMP: bone morphogenetic protein; VEGFA: vascular endothelial growth factor A; TENS-L: tensile strain of low magnitude; TENS-H: tensile strain of high magnitude; LFC: Ligamentum flavum cells; C-FOS: FBJ murine osteosarcoma viral oncogene; SOX9: SRY (sex determining region Y)-box 9; MSX1: Msh homeobox 1.

^aFlexercell model not specified.

increasing flexor tendon proliferation, but when applied in combination with platelet-derived growth factor (PDGF), or a cocktail containing both PDGF and IGF, synthesis of DNA was significantly enhanced.⁹⁶ This could be prevented by blocking gap junction activity with the gap junction inhibitor octanol, indicating the importance of cell–cell communication for cell proliferation.⁹⁷ Therefore, in terms of proliferation, fibroblastic cells appear not to respond as favorably to mechanical stretch signals as osteoblasts or MPCs. For tissue engineering purposes, it will be necessary to screen the appropriate combination of mechanical stretch and growth factors to create functional tendon tissue with appropriate cell number or seed the scaffold with the desired number of cells required to form mature tissue at the outset.

Matrix production

Mechanical stretch has been widely reported to upregulate type I collagen synthesis by tendon cells^{97,98} and ligament cells.^{85,89,90,99} Curiously in the case of the knee joint, type I collagen is upregulated by fibroblasts of the anterior cruciate ligament (ACL)^{85,99} but not those from the medial collateral ligament (MCL).⁸⁵ Conversely, type III collagen is upregulated upon application of strain in MCL cells, but not ACL.^{85,99} These differences in collagen expression profiles might explain differences in healing of ACL versus MCL evident *in vivo*, with lack of healing in ACL due to lack of type III collagen production.⁸⁵ They also deliver a resounding reminder that seemingly comparable cell populations from different topographic sites can behave rather differently to each other, as demonstrated with topographically distinct skin fibroblast populations that have distinct phenotype and gene expression profiles.^{100–102}

In fibroblasts from ligamentum flavum, mechanical stretch induces the expression of collagen types I, III, and V via a TGF β -dependent mechanism.⁸⁹ In static culture, estrogen increases the expression of collagen types I and III,^{99,103} but combined with mechanical stretch signals, actually reduces their expression, along with biglycan.^{99,103} The compound effect of stretch and estrogen might account for the up to eightfold increase in risk of ACL injury in female athletes.^{104–106}

Aside from synthesis of matrix components, it is important to have a controlled balance between proteases that degrade ECM proteins and their inhibitors during matrix turnover and remodeling, and a number of different proteases are upregulated in periodontal ligament (PDL) fibroblasts upon mechanical stimulation. For example, plasminogen activator (PA) is upregulated in mechanically stretched PDL cells.^{91,107} PA is a wide-spectrum serine protease that activates plasminogen to produce plasmin, which in turn activates MMPs to degrade ECM components and cell adhesion molecules,¹⁰⁸ important events during matrix remodeling, cell mobilization, and differentiation. Furthermore, transcript levels of MMPs themselves

(MMP1 and MMP2), along with their inhibitors (tissue inhibitors of metalloproteinase 1 and 2 (TIMP1 and TIMP2)) are upregulated relative to unstretched controls,⁷⁹ again indicating that mechanical stimulation enhances matrix remodeling, which is essential for maintaining structural integrity and strength in ligament tissue *in vivo*. It has also been reported that cell aging leads to increased PA activity,¹⁰⁹ which may reflect the impaired healing capacity that is typically seen in aged individuals.

Exposing tendon cells to IL1 decreased their elastic modulus, thereby increasing their survival after exposure to high-magnitude stretch in 3D collagen gels (see Table 4).¹¹⁵ Coupled with this, IL1 enhanced elastin expression but suppressed type I collagen expression by tendon cells cultured within the constructs,¹¹⁴ which will have a bearing on the mechanical properties of bioengineered constructs that are produced this way. Therefore, a potential means of tuning the mechanical properties of engineered tendon tissues for regenerative therapy may be achievable via regulation not just of the biomaterial substrate but also of the tendon cells themselves.

Impact on cytokine/growth factor production

Growth factors and cytokines can have both autocrine and paracrine effects in the local microenvironment where they are produced. Therefore, the effect of stretch signals applied using Flexercell can give some indication regarding the potential influence of mechanical stretch signals *in vivo* and also potentially enable “priming” of cells to upregulate particular factors for therapeutic applications.

Stretching PDL cells for 24 h (5% elongation at 0.05 Hz) enhanced TGF β production after 7 days⁸⁴ but only in PDL cells from adult teeth and not deciduous teeth. This perhaps reflects the intrinsic difference in longevity between the two as TGF β has a prominent role in matrix formation, maturation, and scar tissue formation that is not needed by deciduous teeth that are shed. On the other hand, stretch signals led to a decrease in M-CSF levels, even after pre-stimulation with vitD3,⁸⁴ which normally stimulates M-CSF production. VEGF production by PDL cells was also upregulated after stretching,^{94,95} as were osteogenic factors BMP2 and BMP6.⁹⁴ VEGF production by tendon fibroblasts was similarly increased and together with the simultaneous upregulation of PGE₂ expression was suggested as a mediator of inflammation and fibrosis, as typically responsible for carpal tunnel syndrome.¹¹³ This suggests that stretch-mediated upregulation of growth factors and cytokines is not necessarily a positive event for fibroblastic cells, and risk of scar tissue formation due to excessive stretch *in vivo* can perhaps be modeled *in vitro* using Flexercell systems.

The magnitude of strain seems to be an important factor in determining cell responses. Low-magnitude tensile strain (from 2% to 8%) was found to have anti-inflammatory effects on PDL cells.^{78,119,120} For example, IL1-mediated

Table 4. Summary of Flexercell studies and key findings using tendon cells.

Cell type	Stage of differentiation	Stretch device and regime	Key findings	Reference
Rabbit Achilles tendon	Tenocytes	Flexercell 3000, 5% elongation, 0.33 Hz for 6 h with 18 h of rest	Synergistic effect of strain and IL1 β to upregulate the production of stromelysin pro-enzyme Strain alone induced no effect	Archambault et al. ¹¹⁰
Canine patellar tendon	Tenocytes	Flexercell 3000, 1%–9% strain, 0.5–3.0 Hz, 15–120 min	Transient JNK activation peaking at 30 min and returning to basal levels by 2 h	Arnoczky et al. ¹¹¹
Avian flexor tendon	Epitenon and tendon internal fibroblasts	Flexercell 2000, 5% elongation, 1 Hz for 8 h	Strain stimulates proliferation in the presence of PDGF-BB and IGF1 and promotes phosphorylation in multiple proteins	Banes et al. ⁹⁶
Avian flexor tendon	Tenocytes	Flexercell, ^a 0.65%, 1 Hz, 8 h/day, 16-h rest	Induced expression of numerous novel genes	Banes et al. ¹¹²
Porcine posterior tibial tendon	Tenocytes	Flexercell, ^a 5% strain, 0.5 Hz for 24 h	Increased COL I and decorin gene expression	Chen et al. ⁹⁸
Human tenosynovial fibroblasts	Tenosynovial fibroblasts	Flexercell, ^a 20%, 1 Hz, 36 h	Increased expression of PGE ₂ and VEGF	Hirata et al. ¹¹³
Human tendon	Tendon internal fibroblasts forming bioartificial tendons	Flexercell 4000, 30% strain for 10 s	IL1 β low dose upregulated elastin expression and high dose suppressed COL I expression to increase elasticity and prevent rupture due to strain	Qi et al. ¹¹⁴
Human tendon	Tendon internal fibroblasts forming bioartificial tendons	Flexercell 4000, 3.5% elongation, 1 Hz, for 1 h/day	IL1 β reduced Young's modulus in bioartificial tendons, increasing their tolerance to mechanical strain	Qi et al. ¹¹⁵
Avian flexor tendons	Tenocytes	Flexercell, ^a 75 millistrain at 1 Hz in a regime of 8 h on, 16 h resting, for 96 h	Increased junctional components n-cadherin and vinculin, but no change in actin levels	Ralphs et al. ¹¹⁶
Human flexor tendons	Tenocytes	Flexercell, ^a 3.5% strain at 1 Hz for 5–30 min and 1–2 h	Stimulation of ecto-ATP activity	Tsuzaki et al. ¹¹⁷
Human flexor tendons	Tenocytes	Flexercell, ^a 3.5% strain at 1 Hz for 2-h and 18-h rest	Induced IL1 β , COX2 and MMP3, but not MMP1 ATP modulated stretch-induced gene expression	Tsuzaki et al. ¹¹⁷
Human flexor tendons	Tendon internal fibroblasts	Flexercell 3000, 3%–5% strain at 1 Hz for 1–6 h	Connexin 43 colocalization with actin increased with strain	Wall et al. ¹¹⁸

IL1 β : interleukin 1 β ; PDGF-BB: platelet-derived growth factor-BB; IGF1: insulin-like growth factor 1; PGE₂: prostaglandin E₂; VEGF: vascular endothelial growth factor; COX2: cyclooxygenase 2; MMP: matrix metalloproteinase; JNK: c-Jun N-terminal kinase.

^aFlexercell model not specified.

transcription of IL1, IL6, and IL8 was blocked, whereas production of IL10, which in turn can block synthesis of IL1, IL6, and IL8,¹²¹ was enhanced.¹¹⁹ In this low range, strain was also found to significantly inhibit IL1-mediated activation of COX2 gene expression and PGE₂ synthesis in a dose-dependent fashion,⁷⁸ reinforcing the argument that low-magnitude strain is anti-inflammatory. In contrast, high strain levels (10%–18.5% strain) induce pro-inflammatory responses such as nuclear factor- κ B (NF- κ B)-mediated COX2 gene expression.⁷⁸ Moreover, high strain levels (9% and 18%) significantly enhanced production of IL1 and PGE₂ and the effects were more prominent in PDL cells

from aged tissue versus young.⁹² Cellular aging was also found to impact significantly COX2 gene expression, with aged (late passage) PDL cells producing higher COX2 mRNA levels than young cells after strain.¹²²

High strain levels (18% elongation applied for 12 h at 0.1 Hz) were also found to significantly enhance NO production,¹²³ which has many physiologic functions that include inflammation, which is likely in this case given the other reported inflammatory mediators upregulated by high-magnitude strain. In rabbit Achilles tendon fibroblasts, the application of strain in the presence of IL1 also led to a 17-fold increase in pro-MMP3 production, but no

difference in COX2 expression was observed compared with IL1 treatment alone.¹²⁴ Strain-induced expression of inflammatory genes could be suppressed by the addition of exogenous ATP to the culture.¹¹⁷

Overall, it seems that a growth factor/cytokine expression profile associated with inflammation is produced in response to mechanical stretch signals that is magnitude dependent. However, low-magnitude stretch is actually anti-inflammatory, and this biphasic response runs parallel with physiologic scale observations where chronic overuse results in injury and inflammation.

Cellular maturation and differentiation

Functional tendons are dependent on adequate cell–cell communication and gap junctions are crucial for this.⁹⁷ Specific gap junction proteins n-cadherin and vinculin are both significantly upregulated in tendon cells after exposure to cyclic stretch¹¹⁶ along with connexin-43, which is localized to gap junctions at the ends of actin filaments.¹¹⁸ Enhanced cell–cell communication is very important for stretch-mediated responses, particularly so in 3D constructs populated with tendon fibroblasts, where application of mechanical strain using a non-Flexercell method resulted in a threefold increase in tensile strength of constructs compared with non-loaded constructs after 7 days of loading.¹²⁵ In addition, supplementing cultures with the anabolic steroid, nandrolone decanoate, synergistically enhances matrix remodeling and ultimately tensile strength of bioengineered tendons.¹²⁶ Therefore, applying mechanical strain to tissue engineered constructs creates a more physiologically relevant tissue. Moreover, when considering information gained from many published Flexercell studies, translating the relevance of this information from 2D to 3D microenvironment needs to be considered.

PDL cells can acquire a bone-like phenotype, and mechanical strain is generally considered to promote and enhance this. Wescott et al.⁹⁴ subjected PDL cells to 12% cyclic strain for up to 24 h and then screened an array of 78 different genes associated with osteogenic differentiation and bone metabolism. Several genes associated with osteogenesis were upregulated, including BMP2, BMP6, ALP, SOX9, MSX1, and vascular endothelial growth factor A (VEGFA), suggesting that a program of osteogenic gene expression was invoked. However, there seems to be a minimum duration of exposure to strain, or culture period following exposure, that is required for osteogenic induction to take place as short-term exposure (6 h) was insufficient to enhance the expression of any key osteogenic genes.^{93,94} Over a longer culture period of 48 h, the expression levels of type I collagen and decorin were elevated after stretch of up to 18% elongation.⁹⁰ Continuous cyclic strain of 9% or 18% elongation at 0.1 Hz applied to PDL cells for up to 6 days resulted in enhanced osteoblastic differentiation characterized by increased ALP activity and

expression of OCN, with paralleled decrease in mitogenic potential and epidermal growth factor (EGF) receptor expression.^{88,127} Differentiation was inhibited by the addition of EGF to the culture medium, which reversed the profile.

In a separate study, ALP synthesis was downregulated in PDL cells subjected to cyclic stretch of 15% elongation at 0.5 Hz and even treatment with vitamin D (an agonist of ALP production) was unable to compensate.⁸⁰ Therefore, the PDL contains a source of multipotent cells that can differentiate toward a bone-forming phenotype, and a careful balance between both biological and mechanical signals is required to maintain the niche and direct cell fate. Furthermore, at high strain levels it seems that a fine balance between osteogenic differentiation and inflammation occurs. Similar to osteoblasts, in PDL cells, cyclic strain also suppresses osteoclastogenesis as although osteoclast genes OPG and RANKL are both upregulated,¹²⁸ the former inhibits the stimulatory effects of the latter.

In summary, mechanical stretch signals applied to tendon and ligament cells using Flexercell have been reported to be largely positive, with matrix synthesis and remodeling being upregulated when using moderate strain levels. Moderate strain levels also have documented anti-inflammatory properties and are able to abrogate the effects of cytokines on these cell types. Conversely, high strain levels induce inflammatory pathways and this likely reflects the potential tissue damage caused by high strains *in vivo*.

Effect of mechanical strain on cartilage cells

Hyaline cartilage has a crucial protective role in the bone–bone contacts, such as knee, where it protects the ends of opposing bone structures and dissipates loads evenly across the full surface area of the cartilage. Consequently, in addition to the load bearing that is typically associated with cartilage function, the constituent chondrocytes also experience tensile and shear stresses.^{129–131} A summary of studies assessing the impact of mechanical stretch on chondrocytes using Flexercell is presented in Table 5.

Impact on cell survival, proliferation, and growth

Chondrocyte proliferation is upregulated in response to strain.¹⁵¹ It is not just magnitude of strain that can impact cell responses, but also frequency. This was demonstrated when mechanical stretch of 3% elongation was applied to chondrocytes at 2, 30, or 150 cycles/min (corresponding to 0.03, 0.5, or 2.5 Hz), with the higher frequency increasing DNA synthesis.¹⁵² From this observation we can conclude that a delicate interplay between several factors including magnitude and frequency of applied stretch can influence cell behavior.

Table 5. Summary of Flexercell studies and key findings using chondrocytes.

Cell type	Stage of differentiation	Stretch device and regime	Key findings	Reference
Rabbit articular cartilage	Chondrocytes	Flexercell, ^a 2%–18%, 0.05 Hz, in the presence or absence of IL1 β	Low stretch: inhibitor of IL1 β -dependent NF- κ B nuclear translocation High stretch: involvement of NF- κ B nuclear translocation and synthesis	Agarwal et al. ¹³²
Rat articular cartilage	Chondrocytes	Flexercell 2000, 7% elongation, 0.5 Hz, cyclic stretch (1 s on, 1 s off)	Upregulated expression of MMP13 and cathepsin B No effect on the expression of aggrecan and COL II	Doi et al. ¹³³
Rat articular cartilage	Chondrocytes	Flexercell 4000, 3%, 0.25 Hz, in the presence or absence of IL1 β	Inhibitor of IL1 β -dependent NF- κ B nuclear translocation and cytoplasmic degradation of I κ B β and I κ B α	Dossumbekova et al. ¹³⁴
Human chondrosarcoma (HCS-2/8)	Chondrocytes	Flexercell, ^a high frequency: 30 cycles/min Mid frequency: 1 cycle/2 min Low frequency: 1 cycle/4 min	High magnitude and frequency gene expression of IL1 and MMP9 Continuous stress induces the production of IL1 and MMP9	Fujisawa et al. ¹³⁵
Bovine articular cartilage	Chondrocytes	Flexercell, ^a higher stress at 10 cycles/min, 17% Lower stress at 10 cycle/h, 5%	Increased proteoglycan synthesis at low frequency and magnitude Decreased proteoglycan synthesis at high frequency and magnitude	Fukuda et al. ¹³⁶
Rabbit articular cartilage	Chondrocytes	Flexercell, ^a 0.05 Hz, 20% elongation, in the presence or absence of IL1 β	Anti-inflammatory effect by inhibiting iNOS and therefore NO in IL1 β -activated chondrocytes	Gassner et al. ¹³⁷
Rabbit articular cartilage	Chondrocytes (retain their differentiated phenotype)	Flexercell, ^a 3 cycles/min consisting of 10 s strain, 10 s rest	Reverses IL1 β -induced suppression of proteoglycan synthesis	Gassner et al. ¹³⁸
Bovine articular cartilage and human chondrosarcoma (I05KC)	Chondrocytes	Flexercell, ^a 24% maximal (average 10%), cyclic (2 s on, 2 s off), 1–20 h	Primary cells: increased COL II expression with no change in β 1 integrin I05KC: increased α 5 expression with no change in β 1, α 2 or α v	Holmvalld et al. ¹³⁹
Porcine articular cartilage	Chondrocytes	Flexercell 4000, 10% strain, 0.5 Hz, 1–24 h	Increased NO, PGE ₂ and COX2 expression Anabolic response: Increased COL II and aggrecan expression as an early response Catabolic response: Increased TGF β and MMP1 expression at 24 h	Huang et al. ¹⁴⁰
Human articular cartilage	Chondrocytes	Flexercell, ^a –20 kPa, 0.5 Hz, 24 h	Increased proliferation Enhanced expression of COL II and aggrecan Enhanced integrin α 2 but no change in α 5 or β 1	Lahiji et al. ¹⁴¹
Rabbit articular cartilage	Chondrocytes (retaining their differentiated phenotype)	Flexercell, ^a 6%, 0.05 Hz	Abrogated TNF α -induced inhibition of proteoglycan synthesis	Long et al. ⁸⁷

(Continued)

Table 5. (Continued)

Cell type	Stage of differentiation	Stretch device and regime	Key findings	Reference
Rat articular cartilage	Chondrocytes (retaining their differentiated phenotype)	Flexercell 4000, 3%, 0.25 Hz, 4–24 h	Blocking of IL1 β -dependent pro-inflammatory gene expression (iNOS, COX2, MMP9 and MMP13)	Madhavan et al. ¹⁴²
Bovine articular cartilage	Chondrocytes	Flexercell 3000, 7%, the frequency (10 cycles/min, 3 s)	Enhanced NO synthesis that inhibited PG synthesis	Matsukawa et al. ¹⁴³
Rat articular cartilage	Chondrocytes	Flexercell, ^a elongation 7%, 30 cycles/min, 12–24 h	Protective effect of IL4 on matrix synthesis	Shimizu et al. ¹⁴⁴
Rat articular cartilage	Chondrocytes	Flexercell, ^a 5, 17%	Decreased proteoglycan synthesis and PKC activity	Tanaka et al. ¹⁴⁵
Bovine articular cartilage	Chondrocytes	Flexercell 3000, high (10 cycles/min), low (10 cycles/h)	Caused depolymerization of HA and induced ROS generation	Yamazaki et al. ¹⁴⁶
Rabbit fibrochondrocytes from TMJ	Chondrocytes (retain their differentiated phenotype)	Flexercell, ^a 6% stretch, 3 cycles/min, 0.05 Hz	Stretch suppresses IL1 β -dependent induction of COX2 and PGE ₂ synthesis	Agarwal et al. ¹⁴⁷
Human chondrosarcoma (CS-OKB)	Confluent	Flexercell, ^a 25%, 0.05 Hz, 48 h	Induces the expression of a heat shock protein termed stress-induced chondrocytic (SIC) I	Chano et al. ¹⁴⁸
Rat fibrochondrocytes from TMJ	Chondrocytes (retain their differentiated phenotype)	Bioflex loading stations, 20%, 0.05 Hz, 1–20 h	IL1 β -induced expression of several MMPs inhibited	Deschner et al. ¹⁴⁹
Rat fibrochondrocytes from TMJ	Chondrocytes (retain their differentiated phenotype)	Flexercell 4000, 20%, 0.05 Hz, 1–24 h	Inhibits IL1 β -induced expression of TNF α , TNFR2 and iNOS, but not TNFR1	Deschner et al. ¹⁵⁰

IL1 β : interleukin 1 β ; NF- κ B: nuclear factor- κ B; MMP: matrix metalloproteinase; iNOS: inducible nitric oxide synthase; NO: nitric oxide; TNF α : tumor necrosis factor α ; COX2: cyclooxygenase 2; PG: proteoglycan; PKC: protein kinase C; ROS: reactive oxygen species; TMJ: temporomandibular joint; PGE₂: prostaglandin E₂; TNFR: tumor necrosis factor receptor; I κ B: nuclear factor of kappa light polypeptide gene enhancer in B-cells inhibitor; HA: Hyaluronan.

^aFlexercell model not specified.

Matrix production

Multiple studies using hyaline cartilage have concluded that application of mechanical stretch increases production of the major collagen component of cartilage, type II collagen,^{139,140,152,153} and the integrin subunit α 2,^{139,153} which binds to collagen. Similarly, the proteoglycan aggrecan is also upregulated^{140,143,152,153} as are several important cartilage glycosaminoglycans (GAGs) including chondroitin-6-sulfate, hyaluronic acid, and dermatan sulfate.¹⁵⁴

However, not all studies agree as stretch was found to downregulate transcript levels of aggrecan and type II collagen, but this could be reversed by the addition of IL4.¹⁴⁴ Hyaluronic acid was also found to undergo depolymerization in the presence of strain due to generation of reactive oxygen species.¹⁴⁶ These conflicting observations can perhaps be explained by the recurring theme of magnitude-dependent effects. In the case of matrix production, particularly proteoglycans that along with their component GAGs are important for producing a highly hydrated matrix, the effects of stretch are magnitude dependent.

Low-magnitude strain enhances proteoglycan production and matrix synthesis, whereas high-magnitude strain diminishes it resulting in ECM degradation and onset of inflammation.^{132,133,135,136,155} The process is dependent on protein kinase C activity.¹³⁶ Moreover, application of high-magnitude cyclic strain to cultured chondrocytes was found to upregulate the expression of MMPs that degrade cartilage ECM.^{133,135,155} Cyclic strain is able to prevent suppression of proteoglycan synthesis caused by IL1^{138,142} and tumor necrosis factor α (TNF α)⁸⁷ but again may be magnitude dependent.

Impact on cytokine/growth factor production

In terms of inflammation, the same biphasic response as seen for tendon/ligament cells can be observed. Exposure of chondrocytes to low magnitudes of strain (<8% elongation) is reported to invoke anti-inflammatory effects, whereas high strain levels (up to 18% elongation) augment the inflammatory response.¹³² This indicates some conservation in responses to different cell populations

from throughout the musculoskeletal system with respect to mechanical signal responsiveness. In particular, low-level strain has generally been observed to block the expression of inflammation-related genes that augment cartilage destruction.^{87,132,134,137,142,147,149,150} For example, using high frequency and low strain (3% elongation at 25 MHz), IL1 pro-inflammatory activity was inhibited, due to block of IL1-mediated NF- κ B nuclear translocation,¹³⁴ in a transforming growth factor- β -activated kinase-1 (TAK1)-dependent fashion.¹⁵⁶ This prevented subsequent transcription of inflammatory gene targets, which normally result from NF- κ B activation. Similarly, TNF α -mediated inflammatory responses were blocked by strain *in vitro*.⁸⁷

There are some anomalies to this general trend however, with high-magnitude strain (20% elongation) applied at low frequency (0.05 Hz) reported to produce anti-inflammatory effects by downregulating IL1-dependent inducible NOS production.¹³⁷ Additionally, cyclic strain of 7% elongation at 0.5 Hz was found to induce IL1 β expression and that of MMP13 and cathepsin B,¹³³ both proteolytic enzymes associated with cartilage pathogenesis.

Interestingly, one study reported that in the presence of a single strain regime of 10% elongation at 0.5 Hz, both catabolic/pro-inflammatory and anabolic responses were observed.¹⁴⁰ While production of NO, PGE₂, and COX2 (characteristic of a pro-inflammatory response) were increased, so too were expression levels of type II collagen and the cartilage-specific proteoglycan aggrecan, as mentioned above. This early response was then replaced by elevations of expression of TGF β 1 and TGF β 3, along with MMP1, indicating that matrix remodeling ensues. Collectively, the above data echoes at the cellular level the patient response where gentle exercise and joint mobility are reported to be beneficial for inflamed joints.^{157,158}

In summary, Flexercell studies indicate that low to moderate levels of strain generally promote cartilage anabolic responses, terms of chondrocyte proliferation, GAG, and proteoglycan production, while also conferring anti-inflammatory effects. However, high levels of strain have a tendency to drive inflammatory responses and provide a good deal of correlation to the trauma and injury associated with high strain and overuse in patients.

The effect of strain applied using Flexercell on mechanotransduction mediators

Most of the bone-stimulating effects of mechanical strain occur due to the activation of mechanotransduction pathways, and the major cell surface receptors involved in these pathways are the integrins. In human osteosarcoma cells, a strain regime of 20 kPa at 0.05 Hz resulted in the upregulation of β 1 integrin mRNA expression^{11,28} along with

redistribution of β 1 integrin subunits from the cytoplasm to the cell membrane.¹¹ However, α v integrin expression and distribution were unaffected by mechanical strain,¹¹ indicating that functional activation and continued recruitment of integrins in response to strain are specific to certain subunits of integrin. Interestingly, vitronectin and OPN, both ligands for α v-containing α v β 3, induce greater calcium influx into osteoblasts than other matrix molecules,¹⁵⁹ and the lack of change in α v integrin subunit expression and distribution upon mechanical loading using Flexercell suggests that the cell response to this form of mechanical strain is not calcium-dependent. It has also been reported that both α 2 β 1 and α 5 β 1 can transmit the effect of stretching from the extracellular environment to activate ERK downstream.²⁴ However, it seems that the α -subunits are the most potent mediators, not β 1, and their signal transduction capacity is actin dependent.^{24,160} Disruption of the actin cytoskeleton, for example, by treatment with cytochalasin D, an actin-depolymerizing agent, attenuates stress response pathways in osteoblast cells.^{161–163} This indicates the importance of the cytoskeleton in sustaining mechanotransduction from extracellular input signals.

Mechanotransduction downstream of integrins involves the coordinated regulation of several other key molecules. Src kinase interacts with integrin intracellular domains²⁴ and FAK¹⁰ at the focal adhesion site when mechanical strain is applied. In osteoblasts, this universally leads to stretch-induced activation of mitogen-activated protein kinase (MAPK) pathways and in particular ERK, a key MAPK effector molecule that activates mechanoresponsive transcription factors.^{7,8,10,13,15,18,24,26,164} At the focal adhesion site formed upon mechanical activation of integrins, FAK also undergoes enhanced and sustained association with another tyrosine kinase, proline-rich tyrosine kinase 2 (PYK2),^{10,16} which seems to stabilize FAK and remove the inhibitory Hic5 adaptor protein from the focal adhesion complex upon activation.¹⁶ Collectively, these data confirm that integrins and their associated intracellular partners are necessary for signal transduction downstream of the strain stimulus to be executed.

Canonical wnt signaling in Flexercell-stretched osteoblasts is highly activated, with enhanced β -catenin transcriptional activity reported.^{17,25} This leads to upregulation of numerous wnt target genes such as those encoding cell cycle proteins, the gap junction protein connexin-43, and other transcription factors.²⁵ Phosphorylation of AKT can also promote β -catenin transcriptional activity via glycogen synthase kinase 3 (GSK β) inactivation and subsequent translocation of β -catenin to the nucleus, where the wnt target genes *Wisp1* and *Cox2* are then transcribed.¹⁶⁵ Upregulation of the wnt target gene encoding connexin-43 seems to have a role in enhancing cell–cell communication among mechanically stimulated osteoblasts by enhancing gap junction communication due to increased levels of phosphorylated connexin-43 protein,^{166,167} and in tendon

cells its colocalization with actin presumably facilitates cell communication in the direction of force application.

The role of integrins in mechanotransduction is explicit as they anchor cells to the ECM substrate in which they reside.^{168,169} Less obvious candidates for mechanotransduction are cell surface receptors that do not have an active role in physical anchorage of cells to ECM. It is well known that androgens, in combination with mechanical cues, can control bone metabolism and invoke a shift from high turnover to osteoanabolic mode. Typically, mechanical signaling via the estrogen receptors (ERs) enhances osteoblast proliferation.¹⁷⁰ ERs have a critical role in mechanotransduction and ERK activation, as osteoblasts from ER $\alpha\beta^{-/-}$ mice lacking both α and β forms of ER were unable to activate ERK when stimulated with 5% elongation at 0.05 Hz.⁸ This phenotype could be partially rescued by transfecting ER $\alpha\beta^{-/-}$ osteoblasts with cDNA for either receptor or fully rescued by transfection with cDNA for both receptors. Additionally, the active estrogen compound estradiol acts in synergy with mechanical strain to promote ER-dependent ALP activity.¹⁷¹ U2OS osteosarcoma cells that do not express ERs and have intrinsically low ALP activity were compared with transfected U2OS cells expressing either ER α or ER β for COL1 expression and ALP activity after being subjected to 1.5% cyclic strain at 0.05 Hz. Mechanical strain increased ALP activity and decreased COL1 expression in ER α and ER β mutants, with estrogen having a synergistic effect on ALP activity. However, estrogen itself is dispensable to the process of ER-mediated mechanotransduction as transfecting ER $\alpha\beta^{-/-}$ osteoblasts with mutant ERs incapable of binding to estrogen did not impair ERK activation, demonstrating that the process was not dependent on estrogen. Even so, it cannot be discounted that differences in the cell lines or the strain regimes applied to cells may have a bearing on the outcome.

Integrins that mediate cell adhesion initiate mechanotransduction signals and have shown significant impact on ligament cells upon stretch stimulation. In PDL fibroblasts, assessment of integrin gene expression showed that both $\alpha 6$ and $\beta 1$ were upregulated by stretch.⁷⁹ Cells from the ACL or MCL plated onto type I collagen, fibronectin, or laminin express elevated levels of both $\alpha 5$ and $\beta 1$ when stretched at 5% elongation for 2 h,⁸¹ and in avian tendon cells, src kinase phosphorylation is also increased after mechanical stimulation.⁹⁶ In 3D collagen gels loaded with ACL fibroblasts, strain enhances both $\alpha 5$ and $\beta 1$, but not in $\alpha 1$ expression.⁸² However, the expression of the $\alpha 5$ integrin subunit in PDL cell monolayers subjected to stretch was actually found to be downregulated upon application of 20 kPa negative pressure to each well for 12 h.⁷⁹

As well as integrin-mediated mechanotransduction, calcium signaling has an important role in cell responsiveness to mechanical forces. Calcium signaling between MCL cells is enhanced by mechanical stretch, as determined by

their ability to maintain a calcium wave between cells both aligned longitudinally in microgrooves and in non-aligned cells.⁸⁶ In the case of tendon cells, activation of JNK, a “stress-activated protein kinase,” occurs in response to mechanical strain, and the level of JNK activation is dependent on magnitude of strain applied.¹¹¹ It also seems to take place downstream of calcium signaling as increased intracellular calcium levels also enhance JNK activation independent of strain.¹¹¹ Chronic, sustained activation of JNK in tendon cells could result in cell death and tendon injury.

There was also prominent upregulation of ulnar collateral ligament (UNCL) protein in PDL cells after applied strain.⁸³ UNCL has a central role in mechanotransduction pathways based on evidence that loss-of-function mutations to the homolog in *Drosophila* result in failure of mechanotransduction¹⁷² and so might have important function in ligament physiology.

Summary

Cell behavior is regulated by both biochemical and physical signals. The importance of biophysical cues in determining cell phenotype is becoming more widely understood, as are mechanotransduction pathways that are activated upon mechanical stimulation. Application of mechanical strain, most commonly applied using the Flexercell system, can provide physical stimulation to cells in the bone and surrounding tissues and aid our understanding of mechanical strain impact on health and injury. It may also potentially be used as a means of priming cell components of the tissues prior to seeding them into scaffolds for tissue engineering and in vivo transplantation.

Declaration of conflicting interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This work was supported by the Priority Research Centers Program (Grant No. 2009-0093829), the Global Research Laboratory Program (Grant No. 2015032163) and KRIBB Research Initiative Program, Republic of Korea.

References

1. Brindley D, Moorthy K, Lee JH, et al. Bioprocess forces and their impact on cell behavior: implications for bone regeneration therapy. *J Tissue Eng* 2011; 2011: 620247.
2. Yeatts AB and Fisher JP. Bone tissue engineering bioreactors: dynamic culture and the influence of shear stress. *Bone* 2011; 48(2): 171–181.

3. Banes AJ, Gilbert J, Taylor D, et al. A new vacuum-operated stress-providing instrument that applies static or variable duration cyclic tension or compression to cells in vitro. *J Cell Sci* 1985; 75: 35–42.
4. Burger EH and Klein-Nulend J. Mechanotransduction in bone—role of the lacuno-canalicular network. *FASEB J* 1999; 13(Suppl.): S101–S112.
5. Liegibel UM, Sommer U, Tomakidi P, et al. Concerted action of androgens and mechanical strain shifts bone metabolism from high turnover into an osteoanabolic mode. *J Exp Med* 2002; 196(10): 1387–1392.
6. Tang L, Lin Z and Li YM. Effects of different magnitudes of mechanical strain on Osteoblasts in vitro. *Biochem Biophys Res Commun* 2006; 344(1): 122–128.
7. Zhu J, Zhang X, Wang C, et al. Different magnitudes of tensile strain induce human osteoblasts differentiation associated with the activation of ERK1/2 phosphorylation. *Int J Mol Sci* 2008; 9(12): 2322–2332.
8. Aguirre JI, Plotkin LI, Gortazar AR, et al. A novel ligand-independent function of the estrogen receptor is essential for osteocyte and osteoblast mechanotransduction. *J Biol Chem* 2007; 282(35): 25501–25508.
9. Bhatt KA, Chang EI, Warren SM, et al. Uniaxial mechanical strain: an in vitro correlate to distraction osteogenesis. *J Surg Res* 2007; 143(2): 329–336.
10. Boutahar N, Guignandon A, Vico L, et al. Mechanical strain on osteoblasts activates autophosphorylation of focal adhesion kinase (FAK) and proline-rich tyrosine kinase 2 (PYK2) tyrosine sites involved in ERK activation. *J Biol Chem* 2004; 16: 30588–30599.
11. Carvalho R, Bumann A, Schwarzer C, et al. A molecular mechanism of integrin regulation from bone cells stimulated by orthodontic forces. *Eur J Orthod* 1996; 18(1): 227–235.
12. Cillo JE Jr, Gassner R, Koepsel RR, et al. Growth factor and cytokine gene expression in mechanically strained human osteoblast-like cells: implications for distraction osteogenesis. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2000; 90(2): 147–154.
13. Fan X, Rahnert JA, Murphy TC, et al. Response to mechanical strain in an immortalized pre-osteoblast cell is dependent on ERK1/2. *J Cell Physiol* 2006; 207(2): 454–460.
14. Faure C, Linossier M-T, Malaval L, et al. Mechanical signals modulated vascular endothelial growth factor-A (VEGF-A) alternative splicing in osteoblastic cells through actin polymerisation. *Bone* 2008; 42(6): 1092–1101.
15. Granet C, Boutahar N, Vico L, et al. MAPK and SRC-kinases control EGR-1 and NF- κ B inductions by changes in mechanical environment in osteoblasts. *Biochem Biophys Res Commun* 2001; 284(3): 622–631.
16. Guignandon A, Boutahar N, Rattner A, et al. Cyclic strain promotes shuttling of PYK2/Hic-5 complex from focal contacts in osteoblast-like cells. *Biochem Biophys Res Commun* 2006; 343(2): 407–414.
17. Hens JR, Wilson KM, Dann P, et al. TOPGAL mice show that the canonical Wnt signaling pathway is active during bone development and growth and is activated by mechanical loading in vitro. *J Bone Miner Res* 2005; 20(7): 1103–1113.
18. Jansen J, Weyts F, Westbroek I, et al. Stretch-induced phosphorylation of ERK1/2 depends on differentiation stage of osteoblasts. *J Cell Biochem* 2004; 93(3): 542–551.
19. Kim DW, Lee HJ, Karmin JA, et al. Mechanical loading differentially regulates membrane-bound and soluble RANKL availability in MC3T3-E1 cells. *Ann N Y Acad Sci* 2006; 1068(1): 568–572.
20. Knoll BI, McCarthy TL, Centrella M, et al. Strain-dependent control of transforming growth factor- β function in osteoblasts in an in vitro model: biochemical events associated with distraction osteogenesis. *Plast Reconstr Surg* 2005; 116(1): 224–233.
21. Liu X, Zhang X and Luo Z-P. Strain-related collagen gene expression in human osteoblast-like cells. *Cell Tissue Res* 2005; 322(2): 331–334.
22. Motokawa M, Kaku M, Tohma Y, et al. Effects of cyclic tensile forces on the expression of vascular endothelial growth factor (VEGF) and macrophage-colony-stimulating factor (M-CSF) in murine osteoblastic MC3T3-E1 cells. *J Dent Res* 2005; 84(5): 422–427.
23. Narutomi M, Nishiura T, Sakai T, et al. Cyclic mechanical strain induces interleukin-6 expression via prostaglandin E2 production by cyclooxygenase-2 in MC3T3-E1 osteoblast-like cells. *J Oral Biosci* 2007; 49(1): 65–73.
24. Plotkin LI, Mathov I, Aguirre JI, et al. Mechanical stimulation prevents osteocyte apoptosis: requirement of integrins, Src kinases, and ERKs. *Am J Physiol Cell Physiol* 2005; 289(3): C633–C643.
25. Robinson JA, Chatterjee-Kishore M, Yaworsky PJ, et al. Wnt/ β -catenin signaling is a normal physiological response to mechanical loading in bone. *J Biol Chem* 2006; 281(42): 31720–31728.
26. Toyoshita Y, Iida S, Koshino H, et al. CYP24 promoter activity is affected by mechanical stress and mitogen-activated protein kinase in MG63 osteoblast-like cells. *日本補綴歯科学会雑誌 [Nihon Hotetsu Shika Gakkai Zasshi]* 2008; 52(2): 171–174.
27. Visconti L, Yen E and Johnson R. Effect of strain on bone nodule formation by rat osteogenic cells in vitro. *Arch Oral Biol* 2004; 49(6): 485–492.
28. Carvalho RS, Scott JE and Yen EH. The effects of mechanical stimulation on the distribution of beta 1 integrin and expression of beta 1-integrin mRNA in TE-85 human osteosarcoma cells. *Arch Oral Biol* 1995; 40(3): 257–264.
29. Mikuni-Takagaki Y, Suzuki Y, Kawase T, et al. Distinct responses of different populations of bone cells to mechanical stress. *Endocrinology* 1996; 137(5): 2028–2035.
30. Centrella M, Ji C and McCarthy TL. Control of TGF- β receptor expression in bone. *Front Biosci* 1998; 3: d113–d124.
31. Ji C, Chen Y, McCarthy TL, et al. Cloning the promoter for transforming growth factor-beta type III receptor. Basal and conditional expression in fetal rat osteoblasts. *J Biol Chem* 1999; 274(43): 30487–30494.
32. Centrella M, Casinighino S, Kim J, et al. Independent changes in type I and type II receptors for transforming growth factor beta induced by bone morphogenetic protein 2 parallel expression of the osteoblast phenotype. *Mol Cell Biol* 1995; 15(6): 3273–3281.

33. Baylink DJ, Finkelstein RD and Mohan S. Growth factors to stimulate bone formation. *J Bone Miner Res* 1993; 8(Suppl. 2): S565–S572.
34. Neidlinger-Wilke C, Stalla I, Claes L, et al. Human osteoblasts from younger normal and osteoporotic donors show differences in proliferation and TGF beta-release in response to cyclic strain. *J Biomech* 1995; 28(12): 1411–1418.
35. Kawata A and Mikuni-Takagaki Y. Mechanotransduction in stretched osteocytes—temporal expression of immediate early and other genes. *Biochem Biophys Res Commun* 1998; 246(2): 404–408.
36. Lean JM, Mackay AG, Chow JW, et al. Osteocytic expression of mRNA for c-fos and IGF-I: an immediate early gene response to an osteogenic stimulus. *Am J Physiol* 1996; 270(6 Pt 1): E937–E945.
37. Skerry TM, Bitensky L, Chayen J, et al. Early strain-related changes in enzyme activity in osteocytes following bone loading in vivo. *J Bone Miner Res* 1989; 4(5): 783–788.
38. Siddhivarn C, Banas A, Champagne C, et al. Mechanical loading and delta12prostaglandin J2 induce bone morphogenetic protein-2, peroxisome proliferator-activated receptor gamma-1, and bone nodule formation in an osteoblastic cell line. *J Periodontal Res* 2007; 42(5): 383–392.
39. Kwan Tat S, Padrines M, Theoleyre S, et al. IL-6, RANKL, TNF-alpha/IL-1: interrelations in bone resorption pathophysiology. *Cytokine Growth Factor Rev* 2004; 15(1): 49–60.
40. Suzuki N, Yoshimura Y, Deyama Y, et al. Mechanical stress directly suppresses osteoclast differentiation in RAW264.7 cells. *Int J Mol Med* 2008; 21(3): 291–296.
41. Siddhivarn C, Banas A, Champagne C, et al. Prostaglandin D2 pathway and peroxisome proliferator-activated receptor gamma-1 expression are induced by mechanical loading in an osteoblastic cell line. *J Periodontal Res* 2006; 41(2): 92–100.
42. Hofbauer LC and Heufelder AE. Role of receptor activator of nuclear factor-kappaB ligand and osteoprotegerin in bone cell biology. *J Mol Med* 2001; 79(5–6): 243–253.
43. Tatsumi R, Sheehan SM, Iwasaki H, et al. Mechanical stretch induces activation of skeletal muscle satellite cells in vitro. *Exp Cell Res* 2001; 267(1): 107–114.
44. Tatsumi R, Hattori A, Ikeuchi Y, et al. Release of hepatocyte growth factor from mechanically stretched skeletal muscle satellite cells and role of pH and nitric oxide. *Mol Biol Cell* 2002; 13(8): 2909–2918.
45. Tatsumi R, Hattori A, Allen RE, et al. Mechanical stretch-induced activation of skeletal muscle satellite cells is dependent on nitric oxide production in vitro. *Anim Sci J* 2002; 73(3): 235–239.
46. Chandran R, Knobloch TJ, Anghelina M, et al. Biomechanical signals upregulate myogenic gene induction in the presence or absence of inflammation. *Am J Physiol Cell Physiol* 2007; 293(1): C267–C276.
47. Iwanuma O, Abe S, Hiroki E, et al. Effects of mechanical stretching on caspase and IGF-1 expression during the proliferation process of myoblasts. *Zoolog Sci* 2008; 25(3): 242–247.
48. Bruce SJ, Gardiner BB, Burke LJ, et al. Dynamic transcription programs during ES cell differentiation towards mesoderm in serum versus serum-freeBMP4 culture. *BMC Genomics* 2007; 8(1): 365.
49. Miyazaki M and Esser KA. REDD2 is enriched in skeletal muscle and inhibits mTOR signaling in response to leucine and stretch. *Am J Physiol Cell Physiol* 2009; 296(3): C583–C592.
50. Tatsumi R, Mitsunashi K, Ashida K, et al. Comparative analysis of mechanical stretch-induced activation activity of back and leg muscle satellite cells in vitro. *Anim Sci J* 2004; 75(4): 345–351.
51. Yamada M, Sankoda Y, Tatsumi R, et al. Matrix metalloproteinase-2 mediates stretch-induced activation of skeletal muscle satellite cells in a nitric oxide-dependent manner. *Int J Biochem Cell Biol* 2008; 40(10): 2183–2191.
52. Boonen KJ, Langelaan ML, Polak RB, et al. Effects of a combined mechanical stimulation protocol: value for skeletal muscle tissue engineering. *J Biomech* 2010; 43(8): 1514–1521.
53. Kumar A, Murphy R, Robinson P, et al. Cyclic mechanical strain inhibits skeletal myogenesis through activation of focal adhesion kinase, Rac-1 GTPase, and NF-κB transcription factor. *FASEB J* 2004; 18(13): 1524–1535.
54. Kurokawa K, Abe S, Sakiyama K, et al. Effects of stretching stimulation with different rates on the expression of MyHC mRNA in mouse cultured myoblasts. *Biomed Res* 2007; 28(1): 25–31.
55. Liu J, Liu J, Mao J, et al. Caspase-3-mediated cyclic stretch-induced myoblast apoptosis via a Fas/FasL-independent signaling pathway during myogenesis. *J Cell Biochem* 2009; 107(4): 834–844.
56. Wozniak A and Anderson J. The dynamics of the nitric oxide release-transient from stretched muscle cells. *Int J Biochem Cell Biol* 2009; 41(3): 625–631.
57. Yuan X, Luo S, Lin Z, et al. Cyclic stretch translocates the α2-subunit of the Na pump to plasma membrane in skeletal muscle cells in vitro. *Biochem Biophys Res Commun* 2006; 348(2): 750–757.
58. Zhang SJ, Truskey GA and Kraus WE. Effect of cyclic stretch on β1D-integrin expression and activation of FAK and RhoA. *Am J Physiol Cell Physiol* 2007; 292(6): C2057–C2069.
59. Atherton P, Szewczyk N, Selby A, et al. Cyclic stretch reduces myofibrillar protein synthesis despite increases in FAK and anabolic signalling in L6 cells. *J Physiol* 2009; 587(14): 3719–3727.
60. Clarke M and Feedback DL. Mechanical load induces sarcoplasmic wounding and FGF release in differentiated human skeletal muscle cultures. *FASEB J* 1996; 10(4): 502–509.
61. Demoule A, Decailliot F, Jonson B, et al. Relationship between pressure-volume curve and markers for collagen turn-over in early acute respiratory distress syndrome. *Intensive Care Med* 2006; 32(3): 413–420.
62. Ebihara S, Hussain SN, Danialou G, et al. Mechanical ventilation protects against diaphragm injury in sepsis: interaction of oxidative and mechanical stresses. *Am J Respir Crit Care Med* 2002; 165(2): 221–228.
63. Goto K, Okuyama R, Sugiyama H, et al. Effects of heat stress and mechanical stretch on protein expression in cultured skeletal muscle cells. *Pflugers Arch* 2003; 447(2): 247–253.

64. Hornberger TA, Armstrong DD, Koh TJ, et al. Intracellular signaling specificity in response to uniaxial vs. multi-axial stretch: implications for mechanotransduction. *Am J Physiol Cell Physiol* 2005; 288(1): C185–C194.
65. Hornberger TA. *Regulation of skeletal muscle mass through stretch-induced signaling events*. Chicago, IL: University of Illinois at Chicago, 2004.
66. Hubatsch DA and Jasmin BJ. Mechanical stimulation increases expression of acetylcholinesterase in cultured myotubes. *Am J Physiol Cell Physiol* 1997; 273(6): C2002–C2009.
67. Mitsumoto Y and Klip A. Development regulation of the subcellular distribution and glycosylation of GLUT1 and GLUT4 glucose transporters during myogenesis of L6 muscle cells. *J Biol Chem* 1992; 267(7): 4957–4962.
68. Nguyen HX, Lusic AJ and Tidball JG. Null mutation of myeloperoxidase in mice prevents mechanical activation of neutrophil lysis of muscle cell membranes in vitro and in vivo. *J Physiol* 2005; 565(2): 403–413.
69. Peterson JM and Pizza FX. Cytokines derived from cultured skeletal muscle cells after mechanical strain promote neutrophil chemotaxis in vitro. *J Appl Physiol* 2009; 106(1): 130–137.
70. Sampaolesi M, Yoshida T, Iwata Y, et al. Stretch-induced cell damage in sarcoglycan-deficient myotubes. *Pflugers Arch* 2001; 442(2): 161–170.
71. Tsivitsse SK, Mylona E, Peterson JM, et al. Mechanical loading and injury induce human myotubes to release neutrophil chemoattractants. *Am J Physiol Cell Physiol* 2005; 288(3): C721–C729.
72. Hornberger TA, Stuppard R, Conley KE, et al. Mechanical stimuli regulate rapamycin-sensitive signalling by a phosphoinositide 3-kinase-, protein kinase B- and growth factor-independent mechanism. *Biochem J* 2004; 380(Pt 3): 795–804.
73. Wozniak AC, Pilipowicz O, Yablonka-Reuveni Z, et al. C-Met expression and mechanical activation of satellite cells on cultured muscle fibers. *J Histochem Cytochem* 2003; 51(11): 1437–1445.
74. Wozniak AC and Anderson JE. Nitric oxide-dependence of satellite stem cell activation and quiescence on normal skeletal muscle fibers. *Dev Dyn* 2007; 236(1): 240–250.
75. Kurokawa K, Itoh T, Kuwahara T, et al. Comparative metagenomics revealed commonly enriched gene sets in human gut microbiomes. *DNA Res* 2007; 14(4): 169–181.
76. Fraser C, Galeotti C, Grandi G, et al. *Neisseria meningitidis antigens and compositions*. Patent, 2009, <http://www.google.co.in/patents/EP2261345A3?cl=ko>
77. Dugan JM, Cartmell SH and Gough JE. Uniaxial cyclic strain of human adipose-derived mesenchymal stem cells and C2C12 myoblasts in coculture. *J Tissue Eng* 2014; 5: 2041731414530138.
78. Agarwal S, Long P, Seyedain A, et al. A central role for the nuclear factor- κ B pathway in anti-inflammatory and proinflammatory actions of mechanical strain. *FASEB J* 2003; 17(8): 899–901.
79. Bolcato-Bellemin A, Elkaim R, Abehsera A, et al. Expression of mRNAs encoding for α and β integrin subunits, MMPs, and TIMPs in stretched human periodontal ligament and gingival fibroblasts. *J Dent Res* 2000; 79(9): 1712–1716.
80. Chiba M and Mitani H. Cytoskeletal changes and the system of regulation of alkaline phosphatase activity in human periodontal ligament cells induced by mechanical stress. *Cell Biochem Funct* 2004; 22(4): 249–256.
81. Hannafin JA, Attia EA, Henshaw R, et al. Effect of cyclic strain and plating matrix on cell proliferation and integrin expression by ligament fibroblasts. *J Orthop Res* 2006; 24(2): 149–158.
82. Henshaw DR, Attia E, Bhargava M, et al. Canine ACL fibroblast integrin expression and cell alignment in response to cyclic tensile strain in three-dimensional collagen gels. *J Orthop Res* 2006; 24(3): 481–490.
83. Kim H-J, Choi YS, Jeong M-J, et al. Expression of UNCL during development of periodontal tissue and response of periodontal ligament fibroblasts to mechanical stress in vivo and in vitro. *Cell Tissue Res* 2007; 327(1): 25–31.
84. Kimoto S, Matsuzawa M, Matsubara S, et al. Cytokine secretion of periodontal ligament fibroblasts derived from human deciduous teeth: effect of mechanical stress on the secretion of transforming growth factor- β 1 and macrophage colony stimulating factor. *J Periodontal Res* 1999; 34(5): 235–243.
85. Hsieh AH, Tsai CMH, Ma QJ, et al. Time-dependent increases in type-III collagen gene expression in medial collateral ligament fibroblasts under cyclic strains. *J Orthop Res* 2000; 18(2): 220–227.
86. Jones BF, Wall ME, Carroll RL, et al. Ligament cells stretch-adapted on a microgrooved substrate increase intercellular communication in response to a mechanical stimulus. *J Biomech* 2005; 38(8): 1653–1664.
87. Long P, Gassner R and Agarwal S. Tumor necrosis factor α -dependent proinflammatory gene induction is inhibited by cyclic tensile strain in articular chondrocytes in vitro. *Arthritis Rheum* 2001; 44(10): 2311–2319.
88. Matsuda N, Yokoyama K, Takeshita S, et al. Role of epidermal growth factor and its receptor in mechanical stress-induced differentiation of human periodontal ligament cells in vitro. *Arch Oral Biol* 1998; 43(12): 987–997.
89. Nakatani T, Marui T, Hitora T, et al. Mechanical stretching force promotes collagen synthesis by cultured cells from human ligamentum flavum via transforming growth factor- β 1. *J Orthop Res* 2002; 20(6): 1380–1386.
90. Ozaki S, Kaneko S, Podyma-Inoue K, et al. Modulation of extracellular matrix synthesis and alkaline phosphatase activity of periodontal ligament cells by mechanical stress. *J Periodontal Res* 2005; 40(2): 110–117.
91. Ozawa Y, Shimizu N and Abiko Y. Low-energy diode laser irradiation reduced plasminogen activator activity in human periodontal ligament cells. *Lasers Surg Med* 1997; 21(5): 456–463.
92. Shimizu N, Yamaguchi M, Uesu K, et al. Stimulation of prostaglandin E2 and Interleukin-1 β production from old rat periodontal ligament cells subjected to mechanical stress. *J Gerontol A Biol Sci Med Sci* 2000; 55(10): B489–B495.
93. Yamaguchi N, Chiba M and Mitani H. The induction of c-fos mRNA expression by mechanical stress in human periodontal ligament cells. *Arch Oral Biol* 2002; 47(6): 465–471.

94. Wescott D, Pinkerton M, Gaffey B, et al. Osteogenic gene expression by human periodontal ligament cells under cyclic tension. *J Dent Res* 2007; 86(12): 1212–1216.
95. Yoshino H, Morita I, Murota SI, et al. Mechanical stress induces production of angiogenic regulators in cultured human gingival and periodontal ligament fibroblasts. *J Periodontal Res* 2003; 38(4): 405–410.
96. Banes AJ, Tsuzaki M, Hu P, et al. PDGF-BB, IGF-I and mechanical load stimulate DNA synthesis in avian tendon fibroblasts in vitro. *J Biomech* 1995; 28(12): 1505–1513.
97. Banes AJ, Weinhold P, Yang X, et al. Gap junctions regulate responses of tendon cells ex vivo to mechanical loading. *Clin Orthop Relat Res* 1999; 367(Suppl.): S356–S370.
98. Chen CH, Marymont JV, Huang MH, et al. Mechanical strain promotes fibroblast gene expression in presence of corticosteroid. *Connect Tissue Res* 2007; 48(2): 65–69.
99. Lee C-Y, Liu X, Smith CL, et al. The combined regulation of estrogen and cyclic tension on fibroblast biosynthesis derived from anterior cruciate ligament. *Matrix Biol* 2004; 23(5): 323–329.
100. Chang HY, Chi JT, Dudoit S, et al. Diversity, topographic differentiation, and positional memory in human fibroblasts. *Proc Natl Acad Sci U S A* 2002; 99(20): 12877–12882.
101. Enoch S, Wall I, Peake M, et al. Increased oral fibroblast lifespan is telomerase-independent. *J Dent Res* 2009; 88(10): 916–921.
102. Enoch S, Peake MA, Wall I, et al. “Young” oral fibroblasts are geno/phenotypically distinct. *J Dent Res* 2010; 89(12): 1407–1413.
103. Lee CY, Smith CL, Zhang X, et al. Tensile forces attenuate estrogen-stimulated collagen synthesis in the ACL. *Biochem Biophys Res Commun* 2004; 317(4): 1221–1225.
104. Gray J, Taunton JE, McKenzie DC, et al. A survey of injuries to the anterior cruciate ligament of the knee in female basketball players. *Int J Sports Med* 1985; 6(6): 314–316.
105. Hutchinson MR and Ireland ML. Knee injuries in female athletes. *Sports Med* 1995; 19(4): 288–302.
106. Zelisko JA, Noble HB and Porter M. A comparison of men’s and women’s professional basketball injuries. *Am J Sports Med* 1982; 10(5): 297–299.
107. Yamaguchi M, Shimizu N, Ozawa Y, et al. Effect of tension-force on plasminogen activator activity from human periodontal ligament cells. *J Periodontal Res* 1997; 32(3): 308–314.
108. Werb Z, Mainardi CL, Vater CA, et al. Endogenous activation of latent collagenase by rheumatoid synovial cells. *N Engl J Med* 1977; 296(18): 1017–1023.
109. Miura S, Yamaguchi M, Shimizu N, et al. Mechanical stress enhances expression and production of plasminogen activator in aging human periodontal ligament cells. *Mech Ageing Dev* 2000; 112(3): 217–231.
110. Archambault JM, Elfervig-Wall MK, Tsuzaki M, et al. Rabbit tendon cells produce MMP-3 in response to fluid flow without significant calcium transients. *J Biomech* 2002; 35(3): 303–309.
111. Arnoczky SP, Tian T, Lavagnino M, et al. Activation of stress-activated protein kinases (SAPK) in tendon cells following cyclic strain: the effects of strain frequency, strain magnitude, and cytosolic calcium. *J Orthop Res* 2002; 20(5): 947–952.
112. Banes AJ, Horesovsky G, Larson C, et al. Mechanical load stimulates expression of novel genes in vivo and in vitro in avian flexor tendon cells. *Osteoarthritis Cartilage* 1999; 7(1): 141–153.
113. Hirata H, Nagakura T, Tsujii M, et al. The relationship of VEGF and PGE2 expression to extracellular matrix remodelling of the tenosynovium in the carpal tunnel syndrome. *J Pathol* 2004; 204(5): 605–612.
114. Qi J, Chi L, Maloney M, et al. Interleukin-1beta increases elasticity of human bioartificial tendons. *Tissue Eng* 2006; 12(10): 2913–2925.
115. Qi J, Fox AM, Alexopoulos LG, et al. IL-1beta decreases the elastic modulus of human tenocytes. *J Appl Physiol* 2006; 101(1): 189–195.
116. Ralphs J, Waggett A and Benjamin M. Actin stress fibres and cell-cell adhesion molecules in tendons: organisation in vivo and response to mechanical loading of tendon cells in vitro. *Matrix Biol* 2002; 21(1): 67–74.
117. Tsuzaki M, Bynum D, Almekinders L, et al. ATP modulates load-inducible IL-1 β , COX 2, and MMP-3 gene expression in human tendon cells. *J Cell Biochem* 2003; 89(3): 556–562.
118. Wall ME, Otey C, Qi J, et al. Connexin 43 is localized with actin in tenocytes. *Cell Motil Cytoskeleton* 2007; 64(2): 121–130.
119. Long P, Hu J, Piesco N, et al. Low magnitude of tensile strain inhibits IL-1beta-dependent induction of pro-inflammatory cytokines and induces synthesis of IL-10 in human periodontal ligament cells in vitro. *J Dent Res* 2001; 80(5): 1416–1420.
120. Long P, Liu F, Piesco NP, et al. Signaling by mechanical strain involves transcriptional regulation of proinflammatory genes in human periodontal ligament cells in vitro. *Bone* 2002; 30(4): 547–552.
121. Pertolani M. Interleukin-10: an anti-inflammatory cytokine with therapeutic potential. *Clin Exp Allergy* 1999; 29: 1164–1171.
122. Ohzeki K, Yamaguchi M, Shimizu N, et al. Effect of cellular aging on the induction of cyclooxygenase-2 by mechanical stress in human periodontal ligament cells. *Mech Ageing Dev* 1999; 108(2): 151–163.
123. Kikuiiri T, Hasegawa T, Yoshimura Y, et al. Cyclic tension force activates nitric oxide production in cultured human periodontal ligament cells. *J Periodontol* 2000; 71(4): 533–539.
124. Archambault J, Tsuzaki M, Herzog W, et al. Stretch and interleukin-1beta induce matrix metalloproteinases in rabbit tendon cells in vitro. *J Orthop Res* 2002; 20(1): 36–39.
125. Garvin J, Qi J, Maloney M, et al. Novel system for engineering bioartificial tendons and application of mechanical load. *Tissue Eng* 2003; 9(5): 967–979.
126. Triantafillopoulos IK, Banes AJ, Bowman KF Jr, et al. Nandrolone decanoate and load increase remodeling and strength in human supraspinatus bioartificial tendons. *Am J Sports Med* 2004; 32(4): 934–943.
127. Matsuda N, Morita N, Matsuda K, et al. Proliferation and differentiation of human osteoblastic cells associated with differential activation of MAP kinases in response to epidermal growth factor, hypoxia, and mechanical stress in vitro. *Biochem Biophys Res Commun* 1998; 249(2): 350–354.

128. Kanzaki H, Chiba M, Sato A, et al. Cyclical tensile force on periodontal ligament cells inhibits osteoclastogenesis through OPG induction. *J Dent Res* 2006; 85(5): 457–462.
129. Soltz MA and Ateshian GA. A Conewise Linear Elasticity mixture model for the analysis of tension-compression nonlinearity in articular cartilage. *J Biomech Eng* 2000; 122(6): 576–586.
130. Huang CY, Soltz MA, Kopacz M, et al. Experimental verification of the roles of intrinsic matrix viscoelasticity and tension-compression nonlinearity in the biphasic response of cartilage. *J Biomech Eng* 2003; 125(1): 84–93.
131. Seo S-J, Mahapatra C, Singh RK, et al. Strategies for osteochondral repair: focus on scaffolds. *J Tissue Eng* 2014; 5: 2041731414541850.
132. Agarwal S, Deschner J, Long P, et al. Role of NF- κ B transcription factors in antiinflammatory and proinflammatory actions of mechanical signals. *Arthritis Rheum* 2004; 50(11): 3541–3548.
133. Doi H, Nishida K, Yorimitsu M, et al. Interleukin-4 downregulates the cyclic tensile stress-induced matrix metalloproteinases-13 and cathepsin B expression by rat normal chondrocytes. *Acta Med Okayama* 2008; 62(2): 119–126.
134. Dossumbekova A, Anghelina M, Madhavan S, et al. Biomechanical signals inhibit IKK activity to attenuate NF- κ B transcription activity in inflamed chondrocytes. *Arthritis Rheum* 2007; 56(10): 3284–3296.
135. Fujisawa T, Hattori T, Takahashi K, et al. Cyclic mechanical stress induces extracellular matrix degradation in cultured chondrocytes via gene expression of matrix metalloproteinases and interleukin-1. *J Biochem* 1999; 125(5): 966–975.
136. Fukuda K, Asada S, Kumano F, et al. Cyclic tensile stretch on bovine articular chondrocytes inhibits protein kinase C activity. *J Lab Clin Med* 1997; 130(2): 209–215.
137. Gassner R, Buckley MJ, Georgescu H, et al. Cyclic tensile stress exerts antiinflammatory actions on chondrocytes by inhibiting inducible nitric oxide synthase. *J Immunol* 1999; 163(4): 2187–2192.
138. Gassner RJ, Buckley MJ, Studer RK, et al. Interaction of strain and interleukin-1 in articular cartilage: effects on proteoglycan synthesis in chondrocytes. *Int J Oral Maxillofac Surg* 2000; 29(5): 389–394.
139. Holmvall K, Camper L, Johansson S, et al. Chondrocyte and chondrosarcoma cell integrins with affinity for collagen type II and their response to mechanical stress. *Exp Cell Res* 1995; 221(2): 496–503.
140. Huang J, Ballou L and Hasty K. Cyclic equibiaxial tensile strain induces both anabolic and catabolic responses in articular chondrocytes. *Gene* 2007; 404(1): 101–109.
141. Lahiji K, Polotsky A, Hungerford DS, et al. Cyclic strain stimulates proliferative capacity, α 2 and α 5 integrin, gene marker expression by human articular chondrocytes propagated on flexible silicone membranes. *In Vitro Cell Dev Biol Anim* 2004; 40(5–6): 138–142.
142. Madhavan S, Anghelina M, Rath-Deschner B, et al. Biomechanical signals exert sustained attenuation of proinflammatory gene induction in articular chondrocytes. *Osteoarthritis Cartilage* 2006; 14(10): 1023–1032.
143. Matsukawa M, Fukuda K, Yamasaki K, et al. Enhancement of nitric oxide and proteoglycan synthesis due to cyclic tensile strain loaded on chondrocytes attached to fibronectin. *Inflamm Res* 2004; 53(6): 239–244.
144. Shimizu A, Watanabe S, Iimoto S, et al. Interleukin-4 protects matrix synthesis in chondrocytes under excessive mechanical stress in vitro. *Mod Rheumatol* 2004; 14(4): 296–300.
145. Tanaka S, Hamanishi C, Kikuchi H, et al. Factors related to degradation of articular cartilage in osteoarthritis: a review. *Seminars in Arthritis and Rheumatism* 1998; 27(6): 392–399.
146. Yamazaki K, Fukuda K, Matsukawa M, et al. Cyclic tensile stretch loaded on bovine chondrocytes causes depolymerization of hyaluronan: involvement of reactive oxygen species. *Arthritis Rheum* 2003; 48(11): 3151–3158.
147. Agarwal S, Long P, Gassner R, et al. Cyclic tensile strain suppresses catabolic effects of interleukin-1 β in fibrochondrocytes from the temporomandibular joint. *Arthritis Rheum* 2001; 44(3): 608–617.
148. Chano T, Tanaka M, Hukuda S, et al. Mechanical stress induces the expression of high molecular mass heat shock protein in human chondrocytic cell line CS-OKB. *Osteoarthritis Cartilage* 2000; 8(2): 115–119.
149. Deschner J, Rath-Deschner B and Agarwal S. Regulation of matrix metalloproteinase expression by dynamic tensile strain in rat fibrochondrocytes. *Osteoarthritis Cartilage* 2006; 14(3): 264–272.
150. Deschner J, Rath-Deschner B, Wypasek E, et al. Biomechanical strain regulates TNFR2 but not TNFR1 in TMJ cells. *J Biomech* 2007; 40(7): 1541–1549.
151. Marques MR, Hajjar D, Franchini KG, et al. Mandibular appliance modulates condylar growth through integrins. *J Dent Res* 2008; 87(2): 153–158.
152. Ueki M, Tanaka N, Tanimoto K, et al. The effect of mechanical loading on the metabolism of growth plate chondrocytes. *Ann Biomed Eng* 2008; 36(5): 793–800.
153. Lahiji K, Polotsky A, Hungerford D, et al. Cyclic strain stimulates proliferative capacity, alpha-2 integrin by human articular chondrocytes from osteoarthritic knee joints. *Univ Pennsylvania Orthop J* 2002; 15: 75–81.
154. Carvalho RS, Yen EH and Suga DM. Glycosaminoglycan synthesis in the rat articular disk in response to mechanical stress. *Am J Orthod Dentofacial Orthop* 1995; 107(4): 401–410.
155. Honda K, Ohno S, Tanimoto K, et al. The effects of high magnitude cyclic tensile load on cartilage matrix metabolism in cultured chondrocytes. *Eur J Cell Biol* 2000; 79(9): 601–609.
156. Madhavan S, Anghelina M, Sjoström D, et al. Biomechanical signals suppress TAK1 activation to inhibit NF- κ B transcriptional activation in fibrochondrocytes. *J Immunol* 2007; 179(9): 6246–6254.
157. Salter RB. The physiologic basis of continuous passive motion for articular cartilage healing and regeneration. *Hand Clin* 1994; 10(2): 211–219.
158. Kim HK, Kerr RG, Cruz TF, et al. Effects of continuous passive motion and immobilization on synovitis and cartilage degradation in antigen induced arthritis. *J Rheumatol* 1995; 22(9): 1714–1721.
159. Miyachi A, Gotoh M, Kamioka H, et al. AlphaVbeta3 integrin ligands enhance volume-sensitive calcium influx in mechanically stretched osteocytes. *J Bone Miner Metab* 2006; 24(6): 498–504.

160. Geng WD, Boskovic G, Fultz ME, et al. Regulation of expression and activity of four PKC isozymes in confluent and mechanically stimulated UMR-108 osteoblastic cells. *J Cell Physiol* 2001; 189(2): 216–228.
161. Hara F, Fukuda K, Asada S, et al. Cyclic tensile stretch inhibition of nitric oxide release from osteoblast-like cells is both G protein and actin-dependent. *J Orthop Res* 2001; 19(1): 126–131.
162. Hara F, Fukuda K, Ueno M, et al. Pertussis toxin-sensitive G proteins as mediators of stretch-induced decrease in nitric-oxide release of osteoblast-like cells. *J Orthop Res* 1999; 17(4): 593–597.
163. Yamamoto N, Fukuda K, Matsushita T, et al. Cyclic tensile stretch stimulates the release of reactive oxygen species from osteoblast-like cells. *Calcif Tissue Int* 2005; 76(6): 433–438.
164. Granet C, Vico AG, Alexandre C, et al. MAP and src kinases control the induction of AP-1 members in response to changes in mechanical environment in osteoblastic cells. *Cell Signal* 2002; 14(8): 679–688.
165. Case N, Ma M, Sen B, et al. Beta-catenin levels influence rapid mechanical responses in osteoblasts. *J Biol Chem* 2008; 283(43): 29196–29205.
166. Grimston SK, Screen J, Haskell JH, et al. Role of connexin43 in osteoblast response to physical load. *Ann N Y Acad Sci* 2006; 1068: 214–224.
167. Ziambaras K, Lecanda F, Steinberg TH, et al. Cyclic stretch enhances gap junctional communication between osteoblastic cells. *J Bone Miner Res* 1998; 13(2): 218–228.
168. Ross TD, Coon BG, Yun S, et al. Integrins in mechanotransduction. *Curr Opin Cell Biol* 2013; 25(5): 613–618.
169. McNamara LE, Sjöström T, Seunarine K, et al. Investigation of the limits of nanoscale filopodial interactions. *J Tissue Eng* 2014; 5: 2041731414536177.
170. Damien E, Price JS and Lanyon LE. Mechanical strain stimulates osteoblast proliferation through the estrogen receptor in males as well as females. *J Bone Miner Res* 2000; 15(11): 2169–2177.
171. Lima F, Vico L, Lafage-Proust MH, et al. Interactions between estrogen and mechanical strain effects on U2OS human osteosarcoma cells are not influenced by estrogen receptor type. *Bone* 2004; 35(5): 1127–1135.
172. Kernan M, Cowan D and Zuker C. Genetic dissection of mechanosensory transduction: mechanoreception-defective mutations of *Drosophila*. *Neuron* 1994; 12(6): 1195–1206.