

Omrani, H., A. Alizadeh, A. Emrouznejad and T. Teplova (2021) Data Envelopment Analysis Model with Decision Makers' Preferences: A Robust Credibility Approach, *Annals of Operations Research*, Accepted. <https://10.1007/s10479-021-04262-2>.

Data Envelopment Analysis Model with Decision Makers' Preferences: A Robust Credibility Approach

Hashem Omrani^{1*}, Arash Alizadeh², Ali Emrouznejad³ and Tamara Teplova⁴

¹ Faculty of Industrial Engineering, Urmia University of Technology, Urmia, Iran, h.omrani@uut.ac.ir

² Department of Industrial and Manufacturing Systems Engineering, University of Missouri, Columbia, MO 65211, USA, aadhb@mail.missouri.edu

³ Aston Business School, Aston University, Birmingham, UK, a.emrouznejad@aston.ac.uk

⁴ National Research University Higher School of Economics, Russian Federation, toma@sani-k.ru

Abstract

Data envelopment analysis (DEA) is one of the widely used methods to measure the efficiency scores of decision making units (DMUs). Conventional DEA is unable to consider both uncertainty in data and decision makers' (DMs) judgments in the evaluations. This study, to address the shortcomings of the conventional DEA, proposes a new best worst method (BWM)- robust credibility DEA (BWM-RCDEA) model to estimate the efficiency scores of DMUs considering DMs' preferences and uncertain data, simultaneously. First, to handle uncertainty in input and output variables, fuzzy credibility model has been applied. Additionally, uncertainty in constructing fuzzy sets is modeled using robust optimization with fuzzy perturbation degree. In this paper, two new types of RCDEA models are proposed: RCDEA model with exact perturbation in fuzzy inputs and outputs and RCDEA model with fuzzy perturbation in fuzzy inputs and outputs. In addition, to deal with flexibility of weights and incorporating DMs' judgement into the RCDEA model, a bi-objective BWM-RCDEA model is introduced. Finally, the proposed bi-objective model is solved using min-max approach. To illustrate the usefulness and capability of the proposed model, efficiency scores of 39 distribution companies in Iran is investigated and results are analyzed and discussed. Finally, based on the results, recommendations have been made for policy makers.

* Corresponding Author, Tel: +98-4433554180, Fax: +98-4413554181, Email address: h.omrani@uut.ac.ir

Keywords: DEA, Credibility theory, Robust optimization, BWM, Efficiency

1. Introduction

Electricity industry plays a major role in economic development of the countries. Indeed, electricity industry as the mother industry, is the engine of economic growth in countries (Tavassoli et al., 2020). Electricity industry consists of three important sectors of power generation, transmission, and electricity distribution. Electricity distribution companies deliver final product to end customers and have direct relationship with customers. Therefore, electricity distribution companies' performance evaluation has a great importance among decision makers, managers, and regulators. In other words, in most cases, the quality of the electricity delivered is directly related to the performance of the distribution companies. Hence, high performance of the distribution companies could improve the economic growth and customers satisfaction. For evaluating distribution companies' performance various methods have been applied in the literature. Data envelopment analysis (DEA) is one of the widely used approach to estimate efficiency score of decision making units (DMUs). Data envelopment analysis (DEA) is a non-parametric method to measure the relative efficiency of a set of DMUs with homogenous inputs and outputs (Charnes et al. 1978). DEA is based on a series of optimizations to specify an efficient frontier curve using linear programming technique. While the numerous advantages of DEA in the field of efficiency evaluations have led researchers devote a considerable number of literatures to develop and apply DEA method in recent two decades (Chen et al., 2019; Emrouznejad and Yang, 2018; Hatami-Marbini et al., 2012), conventional DEA model suffers from some shortcomings such as weight flexibility and uncertainty in input and output data which have been addressed in this study.

One of the crucial assumptions in the conventional DEA is that input and output data are deterministic. In fact, DEA uses a deterministic amount of input to produce a deterministic amount of output. However, in real world applications, data are sometimes contaminated with uncertainty and vagueness. Since DEA is a linear programming model, hence, imprecise and uncertain data may lead to unreliable, incomplete and even infeasible results (Ben Tal and Nemirovski, 2000; Alizadeh and Omrani, 2019). Thereby, some researchers have addressed uncertainties in DEA data using various methods, such as fuzzy sets, robust optimization, stochastic programming, interval programming and etc. methods. This paper deals with uncertainty in input and output variables of DEA using mixed fuzzy sets and robust optimization approaches simultaneously. It is notable that in engineering and management practices it is difficult to collect data with knowing the probability distribution function of data which resulted to unavailability of probability models such as stochastic programming (Yin et al., 2018). In such cases, fuzzy sets and robust optimization are the most appropriate

ways to handle uncertainties. So, a review of the literature on fuzzy sets and robust optimization is summarized in the next section.

In this paper, in order to incorporate DMs' preferences and reducing flexibility in input and output weights, a fuzzy robust DEA model is combined to the latest multiple criteria decision making (MCDM) technique, namely best-worst method (BWM). BWM is a multi-criteria decision making (MCDM) method which was first introduced by Rezaei (2015). BWM applies reference comparisons in order to obtain the weights of criteria. The advantages of BWM in generating more consistent and reliable weights and also needing less pairwise comparison rather than other popular techniques such as AHP led to the application of this method in literatures (Vafadarnikjoo et al., 2020). Readers can refer to comprehensive review of application of BWM technique in decision making problems by Mi et al., (2019).

In summary, the main contributions of this study are as follows:

- Considering uncertainty in inputs and outputs using fuzzy credibility theory
- Considering uncertainty in constructing fuzzy sets using robust optimization approach
- Considering perturbation degree of data as a fuzzy number
- Incorporating preferences into robust-credibility DEA (RCDEA) model using BWM

The rest of the paper is as follows: Literature has been reviewed in section 2. Section 3 presents the methodology (definitions, models and proofs) used in this study. In section 4, the proposed model is applied to evaluate efficiency scores of Iranian electricity distribution companies. Section 5 analyzes the results of the model. Recommendations have been made for policy makers in section 6 and finally, conclusion and summary is made in section 7.

2. Literature Review

In this paper, for considering uncertainty in input and output variables of DEA, fuzzy sets have been applied. Additionally, uncertainty in constructing fuzzy sets has been modeled using robust optimization. We proceed one more level in considering uncertainty and used fuzzy value for perturbation level in robust optimization. Finally, to incorporate DMs' judgement and reduce weight flexibility in conventional DEA, BWM has been used. In this section literature of the used models have been reviewed. y

2.1. Fuzzy DEA

The first steps of incorporating fuzzy set theory into DEA (Emrouznejad and Tavana, 2014) models was taken by Sengupta (1992). Sengupta (1992) introduced a DEA model with fuzzy objective and constraints by defining a tolerance level on DEA constraint violations and analyzed the model using Zimmermann (1978) approach. Later, Triantis and Girod (1998) applied a novel three stage DEA model to measure the technical efficiency in a fuzzy environment. In the first stage, for input and output variables a membership function was defined, so that imprecise input and output variables are expressed in terms of their risk-free and impossible bounds. In the second stage, conventional DEA-VRS and DEA-CRS were formulated in terms of the risk-free and impossible bounds and the membership function for each of the fuzzy input and output variables. Finally, technical efficiencies were measured according to the different membership function values. Their proposed three stage DEA model is categorized in the most popular fuzzy α -level based approach. This method enables decision maker (DM) to observe the impact of modifying input and output values between risk-free and impossible bounds on the efficiency scores. In other words, the main purpose of the α -level based approach is to provide a pair of parametric programs which measures the upper and lower bounds of the α -level based membership function of efficiency scores (Emrouznejad et al., 2014). The α -level based approach was later extended and applied in many literatures. For instance, Liu (2008) in order to evaluate the upper and lower bound performance of flexible manufacturing systems (FMS) alternatives when the input and output data are represented as crisp and fuzzy data, transformed a two-level mathematical program into a one-level DEA assurance region model. Soltanzadeh and Omrani (2018) to deal with fluctuations in data which can be represented by fuzzy numbers, extended the dynamic network DEA model in a fuzzy framework. Their proposed method provided more information for management, since data was presented by membership function. Peykani et al. (2019) proposed an adjustable fuzzy DEA approach while Hatami- Marbini and Saati (2018) proposed a common-weight DEA method to evaluate system efficiency and the component process efficiencies in fuzzy environment.

Another approach to deal with fuzziness in DEA model is the fuzzy ranking which was first developed by Guo and Tanaka (2001). In the fuzzy ranking approach, DM defines a possibility level and the model converts to a linear programming model with crisp constraints and ranking occurs using the comparison rule for fuzzy numbers. Like α -level based approach, fuzzy ranking method has been widely applied in literatures. For instance, Guo (2009) proposed a fuzzy DEA for evaluating objects with fuzzy inputs and outputs and applied his method to analyze a case study involving a restaurant location problem in detail.

Finally, Lertworasikul et al. (2003a; 2003b) based on the Zadeh's (1978) fundamental principles of possibility theory for fuzzy sets, proposed "possibility" and "credibility" approaches to overcome ranking problem in DEA-CRS model. Later, Lertworasikul et al. (2003) extended the possibility approach to fuzzy DEA-VRS model. Their model converted the fuzzy DEA model to a linear model which can be solved using linear programming solvers. Although "possibility" and "credibility" approaches are strong tool to deal with fuzzy and uncertain data, "credibility" mathematical programming is complex and difficult to solve (Amini et al., 2019). However, development of computer and appearance of algorithms have facilitated the use of these approaches in optimization problems (Liu and Liu, 2002). In the credibility approach, the fuzzy DEA model converted into a credibility programming DEA model and fuzzy variables were replaced by expected credits, which were obtained by applying credibility measure (Lertworasikul et al., 2003). Unlike possibility approach, in credibility approach, expected credits are replaced with fuzzy data to deal with fuzzy constraints and fuzzy objectives and there is no need to define any parameter to rank fuzzy efficiency by DM. Generally, credibility approach has been applied in a narrow of literatures. For example, Fasanghari et al., (2015) applied fuzzy credibility constrained programming and P-robust approaches simultaneously for analyzing enterprise architecture scenarios. The catered DEA model was linear, flexible, robust and successful in discrimination power improvement. Also, Amini et al., (2019) to estimate the road safety efficiency of provinces in Iran used a credibility DEA based on road safety (DEA-RS) model. In fact, the constraints of DEA-RS model are considered as credibility constraints and a counterpart credibility DEA-RS (CreDEA-RS) model was proposed. For a comprehensive review on literatures about fuzzy DEA readers can refer to Hatami-Marbini et al. (2011).

2.2. Robust Optimization

As mentioned before, fuzzy sets constructed based on the DMs' opinions. In real world applications, due to the dynamic and continuous change in preferences of stakeholders and DMs, considering certain values for DMs' preferences may lead to the unrealistic and reliable evaluations (Omrani et al., 2018). Therefore, considering perturbation and noise, for DMs assigned values, in order to produce more realistic and precise results is inevitable. One of the most effective approaches to incorporate uncertainty and immunizes model against uncertainty in linear programming is robust optimization (RO) which proposed by Ben-Tal and Nemirovski (2000) and Bertsimas and Sim (2004). According to the Bertsimas and Sim (2004), it is almost implausible that all parameters get their worst case values, therefore, Bertsimas and Sim (2004) for each constraint i introduced a new parameter, Γ_i , to make a trade-off

between the degree of conservatism of the solution and the protection level of the constraint i . Robust optimization has been used in diverse management and engineering practices such as supply chain management (Omrani et al. 2017), waste management (Saeidi-Mobarakeh et al., 2020), investment management (Kim et al., 2018), project selection (Lee et al., 2020), and portfolio optimization (Toloo and Mensah, 2019). Also, robust optimization has been applied on the conventional DEA as a well-known linear programming model widely. Robust DEA (RDEA) is first introduced by Sadjadi and Omrani (2008) and then developed in many researches (Sadjadi et al., (2011a; 2011b), Omrani (2013), Salahi et al., (2016), Shabanpour et al., (2017) and Toloo and Mensah (2018)). In this paper, the robust optimization approach is used to handle uncertainties in both input and output data, simultaneously. In fact, since data are considered as fuzzy sets, a small perturbation in DMs' preferences in specifying upper, medium and lower bounds may lead to unreal and imprecise results. In addition, in RO, perturbation degree (percentage of uncertainty) is considered as exact values which intensifies the impreciseness of the results. Therefore, in this study, perturbation level is handled using fuzzy sets.

2.3. Assurance Region

The second issue addressed in the paper is weight flexibility in DEA. Each DMU in order to be on the efficient frontier is free to set its own weight (Omrani, 2013). This implies that a DMU can make a full benefit of the weighting system to maximize its own efficiency. This weights flexibility could be considered as a strength of DEA, because inefficient units interpreting is meaningful. In other words, these units cannot reach to the efficient frontier even in situation they can choose weights, freely. However, weight flexibility is the weakness of DEA models, too. Since, in some circumstances, some input and output variables take zero or extreme values (Liu, 2014). Therefore, it will be difficult to interpret the impact of these variables on the efficiency scores. Generally, in the literature, four approaches including common weights (Roll et al., 1991), weight restriction (Dyson and Thanassoulis, 1988), cone ratio (Charnes, et al., 1990) and assurance region (AR) (Thompson et al., 1986) have been proposed to reduce weight flexibility. Among weight restriction approaches, AR is a well- known and appropriate approach to incorporate value judgment such as preference concepts, expert opinions and prior information in analysis (Liu, 2014). AR impose ratios between weights to be in a certain range of values (Khalili et al., 2010). AR is categorized to two ARI and ARII approaches. ARI determines bounds on using a priori information on marginal rates. ARII specifies bound for ratios of inputs/ outputs

weights and mainly is applicable in profit efficiency analysis. Both ARI and ARII are sensitive to scaling of inputs and outputs (Allen et al., 1997).

Sarica and Or (2007) to evaluate performance of 65 electricity generation plants in Turkey applied DEA/AR model. Wang et al., (2008) presented a DEA model with AR for priority derivation in the analytic hierarchy process (AHP), which is referred as the DEA/AR model. Their proposed model overcame the shortcomings of the DEAHP such as illogical local weights, over insensitivity to some comparisons, information loss, and overestimation of some local weights, and provide better priority estimate and better decision conclusions than the DEAHP. Lai et al. (2015) applied AHP method to incorporate the weightings of input and output variables into DEA to evaluate 24 international airports. The results revealed that the discrimination power of their model has been increased in comparison with conventional DEA. Moreover, Degl'Innocenti et al. (2017) examined the efficiency of 116 banks for nine new EU members in Central and Eastern European (CEE) countries over the period 2004–2015 using weight assurance region in two-stage DEA model. They also extended their work by including a window-based approach to take into account the patterns of efficiency over time. Ebrahimi and Khalili (2018) proposed a mixed integer AR- imprecise DEA approach to find the best DMU and applied their model to solve a supplier selection problem. Recently, Omrani et al. (2019) integrated best-worst method (BWM) and DEA to measure the efficiency score of 39 electricity distribution companies in Iran. Also, they extended their proposed model to find common weights of input and output variables in DEA considering DMs' judgement.

The main motivation of this study was to propose a novel model to handle uncertainty in data and DMs' judgement comprehensively using the state-of-the-art techniques. As mentioned before, due to the complexity, few research studies have addressed uncertainty using credibility approach for evaluating the efficiency of DMUs. So, the gap of applying credibility DEA approach in the efficiency evaluation of DMUs was obvious. Although, mixed robust fuzzy DEA method has been applied in a few research studies, however, to the best of the authors knowledge, no study has investigated the robust credibility DEA approach with uncertain perturbation level and weight flexibility yet.

3. Methodology

In this section, the proposed BWM-RCDEA model has been illustrated. In summary, first a fuzzy credibility DEA model is introduced. Then, considering uncertainty in constructing fuzzy sets, a robust credibility DEA (RCDEA) model is used. Finally, DMs' preferences is incorporated into the RCDEA model using BWM and a novel bi-objective

BWM-CRDEA model is constructed. The proposed bi-objective BWM-RCDEA model is solved using a min-max approach. In the following, methods applied into the proposed model have been defined separately.

3.1. Preliminaries

First, a brief review of fuzzy sets definitions and terms have been explained in the following. For more details, the readers can refer to Dubois and Prade (1978), Zimmermann (2001), Liu and Liu (2002, 2003) and Li and Liu (2006). Moreover, the parameters used in models have been presented in Table (1).

----- [Table 1 about here] -----

Definition 1: Let U be a universe set. A fuzzy set \tilde{A} of U is defined by a membership function $\mu_{\tilde{A}}(x) \rightarrow [0, 1], \forall x \in U$.

Definition 2: The α -cut of fuzzy set \tilde{A} , \tilde{A}_{α} , is the crisp set $\tilde{A}_{\alpha} = \{x | \mu_{\tilde{A}}(x) \geq \alpha\}$.

Definition 3: A fuzzy number L-R type is expressed as $\tilde{A} = (m, \alpha, \beta)_{LR}$ with below membership function:

$$\mu_{\tilde{A}}(r) = \begin{cases} L(\frac{m-r}{\alpha}) & r \leq m \\ R(\frac{r-m}{\beta}) & r \geq m \end{cases} \quad (1)$$

where L and R are the left and right functions, respectively, and α and β are the (non-negative) left and right spreads, respectively.

Definition 4: An L-R fuzzy number, $\tilde{A} = (m, \alpha, \beta)_{LR} = (m, \alpha, \beta)$ is a triangular fuzzy number if

$$L(x) = R(x) = \begin{cases} 1-x & 0 \leq x \leq 1 \\ 0 & otherwise \end{cases} \quad (2)$$

Definition 5: Let $\tilde{A} = (m, \alpha, \beta)_{LR}$ and $\tilde{B} = (\bar{m}, \bar{\alpha}, \bar{\beta})_{LR}$ be two positive triangular fuzzy numbers. The addition, subtraction and multiplication of \tilde{A} and \tilde{B} are as follows:

Addition: $\tilde{A} + \tilde{B} = (m, \alpha, \beta)_{LR} + (\bar{m}, \bar{\alpha}, \bar{\beta})_{LR} = (m + \bar{m}, \alpha + \bar{\alpha}, \beta + \bar{\beta})_{LR}$

Subtraction: $\tilde{A} - \tilde{B} = (m, \alpha, \beta)_{LR} - (\bar{m}, \bar{\alpha}, \bar{\beta})_{LR} = (m - \bar{m}, \alpha + \bar{\beta}, \beta + \bar{\alpha})_{LR}$

Multiplication (approximation):

$$\tilde{A} \otimes \tilde{B} = (m, \alpha, \beta)_{LR} \otimes (\bar{m}, \bar{\alpha}, \bar{\beta})_{LR} = (m\bar{m}, m\bar{\alpha} + \bar{m}\alpha - \alpha\bar{\alpha}, m\bar{\beta} + \bar{m}\beta + \beta\bar{\beta})_{LR}$$

Definition 6: A possibility space is defined as $(\Theta, P(\Theta), Pos)$ where Θ is nonempty set, $P(\Theta)$ is the power set of Θ and Pos is the possibility measure. The possibility measure satisfies the below axioms:

1. $Pos(\emptyset) = 0, Pos(X) = 1$;
2. $\forall A, B \in P(\Theta), \text{ if } A \subseteq B \rightarrow Pos(A) \leq Pos(B)$;
3. $Pos(A_1 \cup A_2 \dots \cup A_k) = Sup Pos_j(A_j)$

where X is the universe set.

Definition 7: The necessity measure is defined as $Nec(A) = 1 - Pos(A^c)$ where A^c is the complementary set of A set. The necessity measure satisfies the below axioms:

1. $Nec(\emptyset) = 0, Nec(X) = 1$;
2. $\forall A, B \in P(\Theta), \text{ if } A \subseteq B \rightarrow Nec(A) \leq Nec(B)$;
3. $Nec(A_1 \cap A_2 \dots \cap A_k) = Inf Nec_j(A_j)$

Definition 8: The credibility measure is defined as $Cre(A) = \frac{1}{2} \{Pos(A) + Nec(A)\}$. The credibility measure satisfies the below axioms:

1. $Cre(\emptyset) = 0, Cre(X) = 1$;
2. $\forall A, B \in P(\Theta), \text{ if } A \subseteq B \rightarrow Cre(A) \leq Cre(B)$;
3. $Cre(A) + Cre(A^c) = 1, \forall A \subseteq P(X)$

Definition 9: Let λ be a fuzzy variables. The possibility, necessity and credibility of the fuzzy event $(\lambda \geq r)$ are defined as:

$$Pos(\lambda \geq r) = Sup_{t \geq r} \mu_\lambda(t) \tag{3}$$

$$Nec(\lambda \geq r) = 1 - Pos(\lambda < r) = 1 - Sup_{\lambda < r} \mu_\lambda(t) \tag{4}$$

$$Cre(\lambda \geq r) = \frac{1}{2} \{Pos(\lambda \geq r) + Nec(\lambda \geq r)\} \quad (5)$$

3.2. Data Envelopment Analysis (DEA)

DEA is a nonparametric method that uses linear programming to measure the efficiency of DMUs with multiple inputs and multiple outputs. In DEA, efficiency is defined as a ratio of weighted sum of outputs to a weighted sum of inputs. The data form a frontier, DMUs which are on the frontier are evaluated as efficient. Output-oriented DEA models maximize output for a given quantity of input factors, contrariwise input-oriented models minimize input factors required for a given level of output. The input oriented DEA-CCR model is as follows (Charnes et al., 1978):

$$\begin{aligned} \theta_o^{CCR} &= \max \sum_{i=t+1}^{t+s} w_i y_{io} \\ st : \\ \sum_{i=t+1}^{t+s} w_i y_{ij} - \sum_{i=1}^t w_i x_{ij} &\leq 0, \quad j = 1, \dots, n \\ \sum_{i=1}^t w_i x_{io} &= 1 \\ w_i &\geq 0, \quad i = 1, \dots, t + s \end{aligned} \quad (6)$$

In model (6), j th DMU uses t inputs x_{1j}, \dots, x_{tj} for producing s outputs $y_{(t+1)j}, \dots, y_{(t+s)j}$. Also, the efficiency score of DMU under evaluation is θ_o^{CCR} .

Multiplier form of DEA model (6) contains an equality constraint $\sum_{i=1}^t w_i x_{io} = 1$. Since, this paper assumes uncertainty in both inputs and outputs and the equality constraints have not the robust counterpart, hence, to handle uncertainty in inputs using robust optimization approach, the alternative DEA model (7) introduced by Zohrehbandian et al., (2010) is used instead of model (6).

$$\begin{aligned}
& \max z_o \\
& s.t : \\
& \sum_{i=t+1}^{t+s} w_i y_{io} - \theta_o^{CCR} \sum_{i=1}^t w_i x_{io} \geq z_o ; \\
& \sum_{i=t+1}^{t+s} w_i y_{ij} - \theta_j^{CCR} \sum_{i=1}^t w_i x_{ij} \leq 0, j = 1, \dots, n \\
& \sum_{i=1}^{t+s} w_i = 1 \\
& w_i \geq 0, i = 1, \dots, t+s \\
& z_o \text{ free}
\end{aligned} \tag{7}$$

where θ_j^{CCR} is the efficiency score estimated for j th DMU in model (7). In model (7), the normalization constraint

$\sum_{i=1}^t w_i x_{io} = 1$ has been replaced by $\sum_{i=1}^{t+s} w_i = 1$. Zohrehbandian et al. (2010) proved that the optimal solution of the

models (6) and (7) are the same. In other words, the weights generated by the models (6) and (7) are same.

3.3. Credibility DEA (CDEA) model

In this section, the credibility DEA (CDEA) model is described. For developing DEA using fuzzy credibility theory, first following lemma is proven.

Lemma: Let $\lambda_1 = (m_1, \alpha_1, \beta_1)_{LR}$ and $\lambda_2 = (m_2, \alpha_2, \beta_2)_{LR}$ be two L-R fuzzy numbers with continuous membership functions. For a given confidence level $\gamma \in [0, 1]$ it is proven that (Tavana et al., 2012):

$$I) \text{ If } \gamma \leq 0.5, \text{ then } Cr(\tilde{\lambda}_1 \geq \tilde{\lambda}_2) \geq \gamma \Leftrightarrow m_1 + \beta_1 R^{-1}(2\gamma) \geq m_2 - \alpha_2 R^{-1}(2\gamma)$$

$$II) \text{ If } \gamma > 0.5, \text{ then } Cr(\tilde{\lambda}_1 \geq \tilde{\lambda}_2) \geq \gamma \Leftrightarrow m_1 - \alpha_1 L^{-1}(2(1-\gamma)) \geq m_2 + \beta_2 L^{-1}(2(1-\gamma))$$

Proof. Suppose that

$$\lambda = \lambda_1 - \lambda_2 = (m_1, \alpha_1, \beta_1)_{LR} \oplus (-m_2, \beta_2, \alpha_2)_{LR} = (m_1 - m_2, \alpha_1 + \beta_2, \alpha_2 + \beta_1)_{LR} = (\bar{m}, \bar{\alpha}, \bar{\beta})_{LR} \tag{8}$$

According to definition 9, we have:

$$\begin{aligned}
Cr(\lambda_1 \geq \lambda_2) &= Cr(\lambda_1 - \lambda_2 \geq 0) = Cr(\lambda \geq 0) = \frac{1}{2} \left[P_0, (\lambda \geq 0) + Nec(\lambda \geq 0) \right] \\
&= \frac{1}{2} \left[P_0, (\lambda \geq 0) + 1 - P_0, (\lambda \geq 0) \right] = \frac{1}{2} \left[\sup_{t \geq 0} \mu(t) + 1 - \sup_{t < 0} \mu(t) \right]
\end{aligned} \tag{9}$$

It is clear that the equation (9) can be expressed as follows:

$$Cr(\lambda \geq 0) = \begin{cases} 1 & 0 \leq \bar{m} - \bar{\alpha} \\ \frac{1}{2} \left[1 + 1 - L\left(\frac{\bar{m}}{\bar{\alpha}}\right) \right] = 1 - \frac{1}{2} L\left(\frac{\bar{m}}{\bar{\alpha}}\right) & \bar{m} - \bar{\alpha} \leq 0 \leq \bar{m} \\ \frac{1}{2} \left[R\left(-\frac{\bar{m}}{\bar{\beta}}\right) + 1 - 1 \right] = \frac{1}{2} R\left(\frac{\bar{m}}{\bar{\beta}}\right) & \bar{m} \leq 0 \leq \bar{m} + \bar{\beta} \\ 0 & \bar{m} + \bar{\beta} < 0 \end{cases} \tag{10}$$

If $\gamma \leq 0.5$, then

$$\begin{aligned}
C_r(\lambda \geq 0) \geq \gamma &\Leftrightarrow \frac{1}{2} R\left(\frac{-\bar{m}}{\bar{\beta}}\right) \geq \gamma \Leftrightarrow R\left(\frac{-\bar{m}}{\bar{\beta}}\right) \geq 2\gamma \Leftrightarrow \frac{-\bar{m}}{\bar{\beta}} \leq R^{-1}(2\gamma) \Leftrightarrow -\frac{m_1 - m_2}{\alpha_2 + \beta_1} \leq R^{-1}(2\gamma) \\
&\Leftrightarrow m_1 - m_2 \leq (\alpha_2 + \beta_1)R^{-1}(2\gamma) \Leftrightarrow m_1 + \beta_1 R^{-1}(2\gamma) \geq m_2 - \alpha_2 R^{-1}(2\gamma)
\end{aligned}$$

If $\gamma > 0.5$, then

$$\begin{aligned}
C_r(\lambda \geq 0) \geq \gamma &\Leftrightarrow 1 - \frac{1}{2} L\left(\frac{\bar{m}}{\bar{\alpha}}\right) \geq \gamma \Leftrightarrow \frac{1}{2} L\left(\frac{\bar{m}}{\bar{\alpha}}\right) \leq 1 - \gamma \Leftrightarrow L\left(\frac{\bar{m}}{\bar{\alpha}}\right) \leq 2(1 - \gamma) \\
&\Leftrightarrow \frac{\bar{m}}{\bar{\alpha}} \geq L^{-1}(2(1 - \gamma)) \Leftrightarrow \frac{m_1 - m_2}{\alpha_1 + \beta_2} \geq L^{-1}(2(1 - \gamma)) \Leftrightarrow m_1 - m_2 \geq (\alpha_1 + \beta_2)L^{-1}(2(1 - \gamma)) \\
&\Leftrightarrow m_1 - \alpha_1 L^{-1}(2(1 - \gamma)) \geq m_2 + \beta_2 L^{-1}(2(1 - \gamma))
\end{aligned}$$

The credibility counterpart of dual of model (7) can be expressed as follows:

$$\begin{aligned}
& \max z_o \\
& s.t : \\
& Cr\left(\sum_{i=t+1}^{t+s} w_i \tilde{y}_{io} \geq \theta_o^{CCR} \sum_{i=1}^t w_i \tilde{x}_{io} + z_o\right) \geq \gamma_o \\
& Cr\left(\sum_{i=t+1}^{t+s} w_i \tilde{y}_{ij} \leq \theta_j^{CCR} \sum_{i=1}^t w_i \tilde{x}_{ij}\right) \geq \gamma_j, j = 1, \dots, n \\
& \sum_{i=1}^{t+s} w_i = 1 \\
& w_i \geq 0, i = 1, \dots, t+s \\
& z_o \text{ free}
\end{aligned} \tag{11}$$

where $\tilde{x}_{ij} = (x_{ij}^m, x_{ij}^\alpha, x_{ij}^\beta)_{LR}, i = 1, \dots, t$ and $\tilde{y}_{ij} = (y_{ij}^m, y_{ij}^\alpha, y_{ij}^\beta)_{LR}, i = t+1, \dots, t+s$ are the L - R fuzzy numbers.

The parameter $\gamma(\gamma_o, \gamma_i)$ indicates the minimum level of confidence in feasibility of the constraints. Since the data and perturbations are considered as the L - R fuzzy numbers, hence, to protect the constraints against violation, the parameter γ is used. As the parameter γ increase, we lose more objective function to protect the constraints. The higher the value of the parameter γ , the higher the constraint confidence level. In other words, with increasing confidence level, the value of the objective function of the model (11) decreases. According to definition 3, the membership functions of inputs and outputs $j=1, \dots, n$ can be expressed as follows, respectively:

$$\mu_{\tilde{x}_{ij}}(r) = \begin{cases} L\left(\frac{x_{ij}^m - r}{x_{ij}^\alpha}\right) & r \leq x_{ij}^m, i = 1, \dots, t \\ R\left(\frac{r - x_{ij}^m}{x_{ij}^\beta}\right) & r \geq x_{ij}^m, i = 1, \dots, t \end{cases} \tag{12}$$

$$\mu_{\tilde{y}_{ij}}(r) = \begin{cases} L\left(\frac{y_{ij}^m - r}{y_{ij}^\alpha}\right) & r \leq y_{ij}^m, i = t+1, \dots, t+s \\ R\left(\frac{r - y_{ij}^m}{y_{ij}^\beta}\right) & r \geq y_{ij}^m, i = t+1, \dots, t+s \end{cases} \tag{13}$$

According to Zadeh extension principle, the membership functions of the constraints of model (11) can be expressed as follows:

$$\mu_{\sum_{i=1}^t w_i \tilde{x}_{ij}}(r) = \begin{cases} L\left(\frac{\sum_{i=1}^t w_i x_{ij}^m - r}{\sum_{i=1}^t w_i x_{ij}^\alpha}\right) & r \leq \sum_{i=1}^t w_i x_{ij}^m, j = 1, \dots, n \\ R\left(\frac{r - \sum_{i=1}^t w_i x_{ij}^m}{\sum_{i=1}^t w_i x_{ij}^\beta}\right) & r \geq \sum_{i=1}^t w_i x_{ij}^m, j = 1, \dots, n \end{cases} \quad (14)$$

$$\mu_{\sum_{i=t+1}^{t+s} w_i \tilde{y}_{ij}}(r) = \begin{cases} L\left(\frac{\sum_{i=t+1}^{t+s} w_i y_{ij}^m - r}{\sum_{i=t+1}^{t+s} w_i y_{ij}^\alpha}\right) & r \leq \sum_{i=t+1}^{t+s} w_i y_{ij}^m, j = 1, \dots, n \\ R\left(\frac{r - \sum_{i=t+1}^{t+s} w_i y_{ij}^m}{\sum_{i=t+1}^{t+s} w_i y_{ij}^\beta}\right) & r \geq \sum_{i=t+1}^{t+s} w_i y_{ij}^m, j = 1, \dots, n \end{cases} \quad (15)$$

Based on the membership functions (14) to (15), the fuzzy numbers $\sum_{i=t+1}^{t+s} w_i \tilde{y}_{io}$, $\sum_{i=t+1}^{t+s} w_i \tilde{y}_{ij}$, $\theta_j^{CCR} \sum_{i=1}^t w_i \tilde{x}_{ij}$ and

$\theta_o^{CCR} \sum_{i=1}^t w_i \tilde{x}_{io} + z_o$ are shown as below L - R fuzzy numbers:

$$\sum_{i=t+1}^{t+s} w_i y_{io} = \left(\sum_{i=t+1}^{t+s} w_i y_{io}^m, \sum_{i=t+1}^{t+s} w_i y_{io}^\alpha, \sum_{i=t+1}^{t+s} w_i y_{io}^\beta \right)_{LR} \quad (16)$$

$$\sum_{i=t+1}^{t+s} w_i y_{ij} = \left(\sum_{i=t+1}^{t+s} w_i y_{ij}^m, \sum_{i=t+1}^{t+s} w_i y_{ij}^\alpha, \sum_{i=t+1}^{t+s} w_i y_{ij}^\beta \right)_{LR}, j = 1, \dots, n \quad (17)$$

$$\theta_j^{CCR} \sum_{i=1}^t w_i \tilde{x}_{ij} = \left(\theta_j^{CCR} \sum_{i=1}^t w_i x_{ij}^m, \theta_j^{CCR} \sum_{i=1}^t w_i x_{ij}^\alpha, \theta_j^{CCR} \sum_{i=1}^t w_i x_{ij}^\beta \right)_{LR}, j = 1, \dots, n \quad (18)$$

$$\theta_o^{CCR} \sum_{i=1}^t w_i \tilde{x}_{io} + z_o = \left(\theta_o^{CCR} \sum_{i=1}^t w_i x_{io}^m + z_o, \theta_o^{CCR} \sum_{i=1}^t w_i x_{io}^\alpha, \theta_o^{CCR} \sum_{i=1}^t w_i x_{io}^\beta \right)_{LR} \quad (19)$$

In this study, the data are considered as triangular fuzzy numbers. Hence, according to *definition 4*, we have:

$$L(x) = R(x) = L^{-1}(x) = R^{-1}(x) = 1 - x \quad (20)$$

According to above lemma, for $\gamma_o \leq 0.5$, the first constraint of model (11) is expressed as follows:

$$\begin{aligned}
& \sum_{i=t+1}^{t+s} w_i y_{io}^m + R^{-1}(2\gamma_o) \sum_{i=t+1}^{t+s} w_i y_{io}^\beta \geq \theta_o^{CCR} \sum_{i=1}^t w_i x_{io}^m + z_o - R^{-1}(2\gamma_o) \theta_o^{CCR} \sum_{i=1}^t w_i x_{io}^\alpha \\
& \Rightarrow \sum_{i=t+1}^{t+s} w_i (y_{io}^m + R^{-1}(2\gamma_o) y_{io}^\beta) \geq \theta_o^{CCR} \sum_{i=1}^t w_i (x_{io}^m - R^{-1}(2\gamma_o) x_{io}^\alpha) + z_o \\
& \Rightarrow \sum_{i=t+1}^{t+s} w_i (y_{io}^m + (1-2\gamma_o) y_{io}^\beta) \geq \theta_o^{CCR} \sum_{i=1}^t w_i (x_{io}^m - (1-2\gamma_o) x_{io}^\alpha) + z_o \\
& \Rightarrow \sum_{i=t+1}^{t+s} w_i (y_{io}^m + (1-2\gamma_o) y_{io}^\beta) - \theta_o^{CCR} \sum_{i=1}^t w_i (x_{io}^m - (1-2\gamma_o) x_{io}^\alpha) \geq z_o
\end{aligned} \tag{21}$$

In addition, the second constraint of model (11) is converted to a linear constraint as follows:

$$\begin{aligned}
& \sum_{i=t+1}^{t+s} w_i y_{ij}^m - R^{-1}(2\gamma_j) \sum_{i=t+1}^{t+s} w_i y_{ij}^\alpha \leq \theta_j^{CCR} \sum_{i=1}^t w_i x_{ij}^m + R^{-1}(2\gamma_j) \theta_j^{CCR} \sum_{i=1}^t w_i x_{ij}^\beta \\
& \Rightarrow \sum_{i=t+1}^{t+s} w_i (y_{ij}^m - R^{-1}(2\gamma_j) y_{ij}^\alpha) \leq \theta_j^{CCR} \sum_{i=1}^t w_i (x_{ij}^m + R^{-1}(2\gamma_j) x_{ij}^\beta) \\
& \Rightarrow \sum_{i=t+1}^{t+s} w_i (y_{ij}^m - (1-2\gamma_j) y_{ij}^\alpha) \leq \theta_j^{CCR} \sum_{i=1}^t w_i (x_{ij}^m + (1-2\gamma_j) x_{ij}^\beta) \\
& \Rightarrow \sum_{i=t+1}^{t+s} w_i (y_{ij}^m - (1-2\gamma_j) y_{ij}^\alpha) - \theta_j^{CCR} \sum_{i=1}^t w_i (x_{ij}^m + (1-2\gamma_j) x_{ij}^\beta) \leq 0
\end{aligned} \tag{22}$$

By considering the constraints (21) and (22), the final CDEA model for $\gamma_o, \gamma_j \leq 0.5$ is expressed follows:

$$\begin{aligned}
& \max z_o \\
& s.t : \\
& \sum_{i=t+1}^{t+s} w_i (y_{io}^m + (1-2\gamma_o) y_{io}^\beta) - \theta_o^{CCR} \sum_{i=1}^t w_i (x_{io}^m - (1-2\gamma_o) x_{io}^\alpha) \geq z_o \\
& \sum_{i=t+1}^{t+s} w_i (y_{ij}^m - (1-2\gamma_j) y_{ij}^\alpha) - \theta_j^{CCR} \sum_{i=1}^t w_i (x_{ij}^m + (1-2\gamma_j) x_{ij}^\beta) \leq 0, j = 1, \dots, n \\
& \sum_{i=1}^{t+s} w_i = 1 \\
& w_i \geq 0, i = 1, \dots, t+s \\
& z_o \text{ free}
\end{aligned} \tag{23}$$

Similar to above way and according to above lemma, the first constraint of model (11) for $\gamma_o > 0.5$ is expressed as follows:

$$\begin{aligned}
& \sum_{i=t+1}^{t+s} w_i y_{io}^m - L^{-1}(2(1-\gamma_o)) \sum_{i=t+1}^{t+s} w_i y_{io}^\alpha \geq \theta_o^{CCR} \sum_{i=1}^t w_i x_{io}^m + z_o + L^{-1}(2(1-\gamma_o)) \theta_o^{CCR} \sum_{i=1}^t w_i x_{io}^\beta \\
& \Rightarrow \sum_{i=t+1}^{t+s} w_i (y_{io}^m - L^{-1}(2(1-\gamma_o)) y_{io}^\alpha) \geq \theta_o^{CCR} \sum_{i=1}^t w_i (x_{io}^m + L^{-1}(2(1-\gamma_o)) x_{io}^\beta) + z_o \\
& \Rightarrow \sum_{i=t+1}^{t+s} w_i (y_{io}^m - (2\gamma_o - 1) y_{io}^\alpha) \geq \theta_o^{CCR} \sum_{i=1}^t w_i (x_{io}^m + (2\gamma_o - 1) x_{io}^\beta) + z_o \\
& \Rightarrow \sum_{i=t+1}^{t+s} w_i (y_{io}^m - (2\gamma_o - 1) y_{io}^\alpha) - \theta_o^{CCR} \sum_{i=1}^t w_i (x_{io}^m + (2\gamma_o - 1) x_{io}^\beta) \geq z_o
\end{aligned} \tag{24}$$

Also, the second constraint of model (11) is written as follows:

$$\begin{aligned}
& \sum_{i=t+1}^{t+s} w_i y_{ij}^m + L^{-1}(2(1-\gamma_j)) \sum_{i=t+1}^{t+s} w_i y_{ij}^\beta \leq \theta_j^{CCR} \sum_{i=1}^t w_i x_{ij}^m - L^{-1}(2(1-\gamma_j)) \theta_j^{CCR} \sum_{i=1}^t w_i x_{ij}^\alpha \\
& \Rightarrow \sum_{i=t+1}^{t+s} w_i (y_{ij}^m + L^{-1}(2(1-\gamma_j)) y_{ij}^\beta) \leq \theta_j^{CCR} \sum_{i=1}^t w_i (x_{ij}^m - L^{-1}(2(1-\gamma_j)) x_{ij}^\alpha) \\
& \Rightarrow \sum_{i=t+1}^{t+s} w_i (y_{ij}^m + (2\gamma_j - 1) y_{ij}^\beta) \leq \theta_j^{CCR} \sum_{i=1}^t w_i (x_{ij}^m - (2\gamma_j - 1) x_{ij}^\alpha) \\
& \Rightarrow \sum_{i=t+1}^{t+s} w_i (y_{ij}^m + (2\gamma_j - 1) y_{ij}^\beta) - \theta_j^{CCR} \sum_{i=1}^t w_i (x_{ij}^m - (2\gamma_j - 1) x_{ij}^\alpha) \leq 0
\end{aligned} \tag{25}$$

Finally, the CDEA model for $\gamma_o, \gamma_j > 0.5$ is shown as model (26).

$$\begin{aligned}
& \max z_o \\
& s.t.: \\
& \sum_{i=t+1}^{t+s} w_i (y_{io}^m - (2\gamma_o - 1) y_{io}^\alpha) - \theta_o^{CCR} \sum_{i=1}^t w_i (x_{io}^m + (2\gamma_o - 1) x_{io}^\beta) \geq z_o \\
& \sum_{i=t+1}^{t+s} w_i (y_{ij}^m + (2\gamma_j - 1) y_{ij}^\beta) - \theta_j^{CCR} \sum_{i=1}^t w_i (x_{ij}^m - (2\gamma_j - 1) x_{ij}^\alpha) \leq 0, j = 1, \dots, n \\
& \sum_{i=1}^{t+s} w_i = 1 \\
& w_i \geq 0, i = 1, \dots, t+s \\
& z_o \text{ free}
\end{aligned} \tag{26}$$

3.4. Robust Optimization

Robust optimization is a preminent approach in dealing with uncertainty of data (Alem and Morabito, 2012) since robust optimization assumes no probability distribution for uncertain data as well as the capability in modeling

problems with large number of uncertain parameters. In this study we have investigated the robust optimization model with data uncertainty of U (Bental and Nemirovski, 2000).

Although first steps of robust optimization with data uncertainty of U was taken by Soyester (1973) and then developed by Ben-Tal and Nemirovski (2000) and Bertsimas and Sim (2004), however, the advantages of Bertsimas and Sim (2004) in converting models into simple linear models and generating less conservatism solutions led to wide application of this approach in real world problems.

To illustrate the robust optimization model (Bertsimas and Sim, 2004), consider the following optimization model:

$$\begin{aligned}
 & \max c'x \\
 & \text{s.t. :} \\
 & \sum_j a'_{ij}x_j \leq b_i, \forall i \\
 & l_j \leq x_j \leq u_j, \forall j
 \end{aligned} \tag{27}$$

In the optimization model (27), J_i is a set of uncertain coefficients in particular row i of matrix A . Each independent uncertain parameter \tilde{a}_{ij} symmetrically distributed in a bounded interval $[a_{ij} - \hat{a}_{ij}, a_{ij} + \hat{a}_{ij}]$ which centered at point a_{ij} . Also $\hat{a}_{ij} = e_{ij} \times a_{ij}$ where e_{ij} indicates percentage of perturbation in the nominal value a_{ij} . Moreover, Bertsimas and Sim (2004) introduced a scaled deviation $z_{ij} = \frac{(\hat{a}_{ij} - a_{ij})}{\hat{a}_{ij}}$ which take values in interval $[-1, 1]$. Bertsimas and Sim (2004) believed that in real world problems, it is almost implausible that, all parameters take their worst case values simultaneously, which leads to the conservatism solution. Therefore, for each constraint i , Bertsimas and Sim (2004) introduced a new parameter of Γ_i , as a budget of uncertainty, which takes value in interval $[0, |J_i|]$. Indeed Γ_i values make a trade-off between protection level of constraint and conservatism level of solutions. When $\Gamma_i = 0$, the constraints are vulnerable in dealing with uncertainty while in $\Gamma_i = |J_i|$ constraints are fully protected and solutions are too conservatism. Considering all aforementioned details, Bertsimas and Sim (2004) introduced the robust linear counterpart of model (27) as follows: (For more details, readers can refer to Bertsimas and sim (2004) and Alem and Morabito (2012))

$$\begin{aligned}
& \max c'x \\
& \text{s.t:} \\
& \sum_j a'_{ij}x + \Gamma_i P_i + \sum_{j \in J_i} q_{rj} \leq b_i, \quad \forall i \\
& P_j + q_{rj} \geq \hat{a}_{ij} y_j, \quad \forall j \\
& -y_j \leq x_j \leq y_j, \quad \forall j \\
& I_j \leq x_j \leq u_j, \quad \forall j \\
& q_{ij} \geq 0, \quad \forall i, j \in J_i \\
& y_j \geq 0, \quad \forall j \\
& p_i \geq 0, \quad \forall i
\end{aligned} \tag{28}$$

3.5. Best Worst Method (BWM)

Best worst method was first introduced by Rezaei (2015) to weight criteria using pairwise comparison. Indeed, like AHP, BWM assign the weights based on the preferences of Decision makers. However, BWM has advantages in compared with AHP such as less pairwise comparison and higher consistency ratio (Zhao et al.,2019). For example, for a set of n criteria, AHP needs $\frac{n * (n - 1)}{2}$ pairwise comparison while BWM applies $(2 * n) - 3$, which is less than AHP technique.

The final BWM model is as model (29). It is important to note that the steps of the BWM are completely explained in Rezaei (2015 and 2016) and readers for more details can refer to them.

$$\begin{aligned}
& \min \xi_r \\
& \text{s.t:} \\
& \{ | w_i - a_{iW} w_W | \}_r \leq \xi_r, \quad i = 1, \dots, n \\
& \{ | w_B - a_{Bi} w_i | \}_r \leq \xi_r, \quad i = 1, \dots, n \\
& \sum_{i=1}^n w_i = 1 \\
& w_i \geq 0, \quad j = 1, \dots, n
\end{aligned} \tag{29}$$

Where the preference of the best criterion over i th criteria for r th DM is a_{Bi} and preference of i th criteria on worst criterion for r th DM is a_{iW} . In this paper, linear BWM (29) is applied to estimate the indicators' weights. Also

readers can refer to comprehensive review of application of BWM technique in decision making problems by Mi et al., (2019).

3.6. Robust Credibility DEA (RCDEA)

In this section, the counterpart robust CDEA (RCDEA) models are proposed. The proposed RCDEA models are:

- RCDEA model with exact perturbation in fuzzy inputs and fuzzy outputs
- RCDEA model with fuzzy perturbation in fuzzy inputs and fuzzy outputs

Let $\tilde{x}_{ij} = (x_{ij}^m, x_{ij}^\alpha, x_{ij}^\beta)_{LR}, i = 1, \dots, t$ and $\tilde{y}_{ij} = (y_{ij}^m, y_{ij}^\alpha, y_{ij}^\beta)_{LR}, i = t + 1, \dots, t + s$ be the L - R fuzzy numbers. Let

x_{ij}^m and y_{ij}^m are symmetrically distributed in intervals $[x_{ij}^m - \hat{x}_{ij}^m, x_{ij}^m + \hat{x}_{ij}^m]$ and $[y_{ij}^m - \hat{y}_{ij}^m, y_{ij}^m + \hat{y}_{ij}^m]$ where

$\hat{x}_{ij}^m = e_{ij} \times x_{ij}^m$ and $\hat{y}_{ij}^m = e_{ij} \times y_{ij}^m$. Since x_{ij}^m and y_{ij}^m are contaminated with perturbation, hence, $x_{ij}^\alpha, x_{ij}^\beta, y_{ij}^\alpha$ and y_{ij}^β

are symmetrically distributed in determined intervals, too.

3.6.1. RCDEA model with exact perturbation in fuzzy inputs and fuzzy outputs

To show the RCDEA model with exact perturbation in fuzzy inputs and fuzzy outputs, credibility DEA models (23) and (26) are converted to the counterpart robust CDEA models. According to approach proposed by Bertsimas and Sim (2004), the robust counterpart of CDEA models for $\gamma \leq 0.5$ and $\gamma > 0.5$ are as (30) and (31) respectively:

$$\theta_o^{RCDEA} = \max_{\gamma_o, \gamma_i \leq 0.5} \sum_{i=t+1}^{t+s} w_i (y_{io}^m + (1-2\gamma_o)y_{io}^\beta) - \theta_o^{CCR} \sum_{i=1}^t w_i (x_{io}^m - (1-2\gamma_o)x_{io}^\alpha) - \Gamma_o p_o - \sum_{i=t+1}^{t+s} q_{io} - \sum_{i=1}^t q_{io}$$

s.t.:

$$\sum_{i=t+1}^{t+s} w_i (y_{ij}^m - (1-2\gamma_j)y_{ij}^\alpha) - \theta_j^{CCR} \sum_{i=1}^t w_i (x_{ij}^m + (1-2\gamma_j)x_{ij}^\beta) + \Gamma_j p_j + \sum_{i=t+1}^{t+s} q_{ij} + \sum_{i=1}^t q_{ij} \leq 0, j=1, \dots, n$$

$$\sum_{i=1}^{m+s} w_i = 1$$

$$p_o + q_{io} \geq e_{io} (y_{io}^m + (1-2\gamma_o)y_{io}^\beta) z_i, \quad \forall i = t+1, \dots, t+s \quad (30)$$

$$p_o + q_{io} \geq e_{io} (x_{io}^m - (1-2\gamma_o)x_{io}^\alpha) z_i, \quad \forall i = 1, \dots, t$$

$$p_j + q_{ij} \geq e_{ij} (y_{ij}^m - (1-2\gamma_j)y_{ij}^\alpha) z_i, \quad \forall i = t+1, \dots, t+s, \forall j = 1, \dots, n$$

$$p_j + q_{ij} \geq e_{ij} (x_{ij}^m + (1-2\gamma_j)x_{ij}^\beta) z_i, \quad \forall i = 1, \dots, t, \forall j = 1, \dots, n$$

$$-z_i \leq w_i \leq z_i, \quad \forall i = 1, 2, \dots, t+s$$

$$p_j, q_{ij} \geq 0, \quad \forall i, j$$

$$w_i \geq 0, i = 1, \dots, t+s$$

$$\theta_o^{RCDEA} = \max_{\gamma_o, \gamma_i > 0.5} \sum_{i=t+1}^{t+s} w_i (y_{io}^m - (2\gamma_o - 1)y_{io}^\alpha) - \theta_o^{CCR} \sum_{i=1}^t w_i (x_{io}^m + (2\gamma_o - 1)x_{io}^\beta) - \Gamma_o p_o - \sum_{i=t+1}^{t+s} q_{io} - \sum_{i=1}^t q_{io}$$

s.t.:

$$\sum_{i=t+1}^{t+s} w_i (y_{ij}^m + (2\gamma_j - 1)y_{ij}^\beta) - \theta_j^{CCR} \sum_{i=1}^t w_i (x_{ij}^m - (2\gamma_j - 1)x_{ij}^\alpha) + \Gamma_j p_j + \sum_{i=t+1}^{t+s} q_{ij} - \sum_{i=1}^t q_{ij} \leq 0, j=1, \dots, n$$

$$\sum_{i=1}^{m+s} w_i = 1$$

$$p_o + q_{io} \geq e_{io} (y_{io}^m - (2\gamma_o - 1)y_{io}^\alpha) z_i, \quad \forall i = t+1, \dots, t+s \quad (31)$$

$$p_o + q_{io} \geq e_{io} (x_{io}^m + (2\gamma_o - 1)x_{io}^\beta) z_i, \quad \forall i = 1, \dots, t$$

$$p_j + q_{ij} \geq e_{ij} (y_{ij}^m + (2\gamma_j - 1)y_{ij}^\beta) z_i, \quad \forall i = t+1, \dots, t+s, \forall j = 1, \dots, n$$

$$p_j + q_{ij} \geq e_{ij} (x_{ij}^m - (2\gamma_j - 1)x_{ij}^\alpha) z_i, \quad \forall i = 1, \dots, t, \forall j = 1, \dots, n$$

$$-z_i \leq w_i \leq z_i, \quad \forall i = 1, 2, \dots, t+s$$

$$p_j, q_{ij} \geq 0, \quad \forall i, j$$

$$w_i \geq 0, i = 1, \dots, t+s$$

where the free variable z_o is removed and the first constraint of models (23) and (26) are moved to objective functions.

It is notable that the objective function and constraints of the models (30) and (32) are the same except the constraints related to robust optimization. In fact, the 3th to 6th constraints are different for these models. The constraints show

that the feasible space of model (32) is greater than the feasible space of model (30). Consider the 3th constraint for the models. For the model (30), the third constraint is as $p_o + q_{io} \geq e_{io} (y_{io}^m + (1 - 2\gamma_o) y_{io}^\beta) z_i$ and for the model (32), it express as $p_o + q_{io} \geq [e_{io}^m - (1 - 2\gamma_{io}) e_{io}^\alpha] (y_{io}^m + (1 - 2\gamma_o) y_{io}^\beta) z_i$. Since the value of $[e_{io}^m - (1 - 2\gamma_{io}) e_{io}^\alpha]$ is a positive value, then the feasible space of model (32) is greater than the feasible space of model (30). Hence, the objective function of model (32) is equal or greater than the objective function model (30), namely, $\theta_o^{RCDEA-Fuzzy\ Perturbation} \geq \theta_o^{RCDEA}$. Note that we set $e_{io}^m = e_{io}$. In a similar way, for $\gamma > 0.5$, we can show that the objective function of model (33) is equal or smaller than the objective function model (31).

3.6.2. RCDEA model with fuzzy perturbation in fuzzy inputs and fuzzy outputs

In real world application, the level of uncertainty is unknown and considering certain values for uncertainty level of data (the value e) may be contributed to imprecise results. In this section, the percentage of perturbation is considered as L - R fuzzy number $\tilde{e}_{ij} = (e_{ij}^m, e_{ij}^\alpha, e_{ij}^\beta)_{LR}$. Consider the fifth constraint of RCDEA model (30). In fuzzy status, this constraint is written as $Cre(p_j + q_{ij} \geq \tilde{e}_{ij} (y_{ij}^m + (2\gamma_j - 1) y_{ij}^\beta) z_i) \geq \gamma_{ij}$. According to above lemma, the later fuzzy constraint is converted to $p_j + q_{ij} \geq [e_{ij}^m - (1 - 2\gamma_{ij}) e_{ij}^\alpha] (y_{ij}^m + (2\gamma_j - 1) y_{ij}^\beta) z_i$. In fact, in order to obtain a RCDEA model with fuzzy perturbation, it is enough to replace e_{ij} with $[e_{ij}^m - (1 - 2\gamma_{ij}) e_{ij}^\alpha]$ for $\gamma_{ij} \leq 0.5$ and $[e_{ij}^m + (2\gamma_{ij} - 1) e_{ij}^\beta]$ $\gamma_{ij} > 0.5$. Therefore, models (30) and (31) are converted to (32) and (33), respectively:

$$\theta_o^{RCDEA-Fuzzy\ Perturbation} = \max_{\gamma_o, \gamma_i \leq 0.5} \sum_{i=t+1}^{t+s} w_i (y_{io}^m + (1-2\gamma_o)y_{io}^\beta) - \theta_o^{CCR} \sum_{i=1}^t w_i (x_{io}^m - (1-2\gamma_o)x_{io}^\alpha) - \Gamma_o p_o - \sum_{i=t+1}^{t+s} q_{io} - \sum_{i=1}^t q_{io}$$

s.t.:

$$\sum_{i=t+1}^{t+s} w_i (y_{ij}^m - (1-2\gamma_j)y_{ij}^\alpha) - \theta_j^{CCR} \sum_{i=1}^t w_i (x_{ij}^m + (1-2\gamma_j)x_{ij}^\beta) + \Gamma_j p_j + \sum_{i=t+1}^{t+s} q_{ij} + \sum_{i=1}^t q_{ij} \leq 0, j=1, \dots, n$$

$$\sum_{i=1}^{m+s} w_i = 1$$

$$p_o + q_{io} \geq [e_{io}^m - (1-2\gamma_{io})e_{io}^\alpha](y_{io}^m + (1-2\gamma_o)y_{io}^\beta)z_i, \quad \forall i = t+1, \dots, t+s \quad (32)$$

$$p_o + q_{io} \geq [e_{io}^m - (1-2\gamma_{io})e_{io}^\alpha](x_{io}^m - (1-2\gamma_o)x_{io}^\alpha)z_i, \quad \forall i = 1, \dots, t$$

$$p_j + q_{ij} \geq [e_{ij}^m - (1-2\gamma_{ij})e_{ij}^\alpha](y_{ij}^m - (1-2\gamma_j)y_{ij}^\alpha)z_i, \quad \forall i = t+1, \dots, t+s, \forall j = 1, \dots, n$$

$$p_j + q_{ij} \geq [e_{ij}^m - (1-2\gamma_{ij})e_{ij}^\alpha](x_{ij}^m + (1-2\gamma_j)x_{ij}^\beta)z_i, \quad \forall i = 1, \dots, t, \forall j = 1, \dots, n$$

$$-z_i \leq w_i \leq z_i, \quad \forall i = 1, 2, \dots, t+s$$

$$p_j, q_{ij} \geq 0, \quad \forall i, j$$

$$w_i \geq 0, \quad i = 1, \dots, t+s$$

$$\theta_o^{RCDEA-Fuzzy\ Perturbation} = \max_{\gamma_o, \gamma_i > 0.5} \sum_{i=t+1}^{t+s} w_i (y_{io}^m - (2\gamma_o - 1)y_{io}^\alpha) - \theta_o^{CCR} \sum_{i=1}^t w_i (x_{io}^m + (2\gamma_o - 1)x_{io}^\beta) - \Gamma_o p_o - \sum_{i=t+1}^{t+s} q_{io} - \sum_{i=1}^t q_{io}$$

s.t.:

$$\sum_{i=t+1}^{t+s} w_i (y_{ij}^m + (2\gamma_j - 1)y_{ij}^\beta) - \theta_j^{CCR} \sum_{i=1}^t w_i (x_{ij}^m - (2\gamma_j - 1)x_{ij}^\alpha) + \Gamma_j p_j + \sum_{i=t+1}^{t+s} q_{ij} + \sum_{i=1}^t q_{ij} \leq 0, j=1, \dots, n$$

$$\sum_{i=1}^{m+s} w_i = 1$$

$$p_o + q_{io} \geq [e_{io}^m + (2\gamma_{io} - 1)e_{io}^\beta](y_{io}^m - (2\gamma_o - 1)y_{io}^\alpha)z_i, \quad \forall i = t+1, \dots, t+s \quad (33)$$

$$p_o + q_{io} \geq [e_{io}^m + (2\gamma_{io} - 1)e_{io}^\beta](x_{io}^m + (2\gamma_o - 1)x_{io}^\beta)z_i, \quad \forall i = 1, \dots, t$$

$$p_j + q_{ij} \geq [e_{ij}^m + (2\gamma_{ij} - 1)e_{ij}^\beta](y_{ij}^m + (2\gamma_j - 1)y_{ij}^\beta)z_i, \quad \forall i = t+1, \dots, t+s, \forall j = 1, \dots, n$$

$$p_j + q_{ij} \geq [e_{ij}^m + (2\gamma_{ij} - 1)e_{ij}^\beta](x_{ij}^m - (2\gamma_j - 1)x_{ij}^\alpha)z_i, \quad \forall i = 1, \dots, t, \forall j = 1, \dots, n$$

$$-z_i \leq w_i \leq z_i, \quad \forall i = 1, 2, \dots, t+s$$

$$p_j, q_{ij} \geq 0, \quad \forall i, j$$

$$w_i \geq 0, \quad i = 1, \dots, t+s$$

3.7. BWM-RCDEA model with fuzzy perturbation in fuzzy inputs and fuzzy outputs

In this section, a novel BWM-RCDEA model is proposed. Existence of mutual normalization constraint of $\sum_{i=1}^{m+s} w_i = 1$

in both RCDEA and BWM made it possible to incorporate DMs' preferences into the proposed BWM-RCDEA model.

According to models (32) and (33), the proposed BWM- RCDEA model with fuzzy perturbation for $\gamma \leq 0.5$ and $\gamma > 0.5$ can be expressed as models (34) and (35), respectively:

$$\begin{aligned}
\max_{\gamma_o, \gamma_i \leq 0.5} f_1 &= \sum_{i=t+1}^{t+s} w_i (y_{io}^m + (1-2\gamma_o) y_{io}^\beta) - \theta_o^{CCR} \sum_{i=1}^t w_i (x_{io}^m - (1-2\gamma_o) x_{io}^\alpha) - \Gamma_o p_o - \sum_{i=t+1}^{t+s} q_{io} - \sum_{i=1}^t q_{io} \\
\max f_2 &= -\xi \\
s.t.: \\
\sum_{i=t+1}^{t+s} w_i (y_{ij}^m - (1-2\gamma_j) y_{ij}^\alpha) - \theta_j^{CCR} \sum_{i=1}^t w_i (x_{ij}^m + (1-2\gamma_j) x_{ij}^\beta) + \Gamma_j p_j + \sum_{i=t+1}^{t+s} q_{ij} + \sum_{i=1}^t q_{ij} &\leq 0, j=1, \dots, n \\
\sum_{i=1}^{m+s} w_i &= 1 \\
p_o + q_{io} &\geq [e_{io}^m - (1-2\gamma_o) e_{io}^\alpha] (y_{io}^m + (1-2\gamma_o) y_{io}^\beta) z_i, \quad \forall i = t+1, \dots, t+s \\
p_o + q_{io} &\geq [e_{io}^m - (1-2\gamma_o) e_{io}^\alpha] (x_{io}^m - (1-2\gamma_o) x_{io}^\alpha) z_i, \quad \forall i = 1, \dots, t \\
p_j + q_{ij} &\geq [e_{ij}^m - (1-2\gamma_j) e_{ij}^\alpha] (y_{ij}^m - (1-2\gamma_j) y_{ij}^\alpha) z_i, \quad \forall i = t+1, \dots, t+s, \forall j = 1, \dots, n \\
p_j + q_{ij} &\geq [e_{ij}^m - (1-2\gamma_j) e_{ij}^\alpha] (x_{ij}^m + (1-2\gamma_j) x_{ij}^\beta) z_i, \quad \forall i = 1, \dots, t, \forall j = 1, \dots, n \\
-z_i &\leq w_i \leq z_i, \quad \forall i = 1, 2, \dots, t+s \\
|w_B - a_{Bi} w_i| &\leq \xi, \quad i = 1, \dots, t+s \\
|w_i - a_{iW} w_w| &\leq \xi, \quad i = 1, \dots, t+s \\
p_j, q_{ij} &\geq 0, \quad \forall i, j \\
w_i &\geq 0, \quad i = 1, \dots, t+s
\end{aligned} \tag{34}$$

$$\begin{aligned}
\max_{\gamma_o, \gamma_i > 0.5} f_1 &= \sum_{i=t+1}^{t+s} w_i (y_{io}^m - (2\gamma_o - 1) y_{io}^\alpha) - \theta_o^{CCR} \sum_{i=1}^t w_i (x_{io}^m + (2\gamma_o - 1) x_{io}^\beta) - \Gamma_o p_o - \sum_{i=t+1}^{t+s} q_{io} - \sum_{i=1}^t q_{io} \\
\max f_2 &= -\xi \\
s.t.: \\
\sum_{i=t+1}^{t+s} w_i (y_{ij}^m + (2\gamma_j - 1) y_{ij}^\beta) - \theta_j^{CCR} \sum_{i=1}^t w_i (x_{ij}^m - (2\gamma_j - 1) x_{ij}^\alpha) + \Gamma_j p_j + \sum_{i=t+1}^{t+s} q_{ij} + \sum_{i=1}^t q_{ij} &\leq 0, j=1, \dots, n \\
\sum_{i=1}^{m+s} w_i &= 1 \\
p_o + q_{io} &\geq [e_{io}^m + (2\gamma_o - 1) e_{io}^\beta] (y_{io}^m - (2\gamma_o - 1) y_{io}^\alpha) z_i, \quad \forall i = t+1, \dots, t+s \\
p_o + q_{io} &\geq [e_{io}^m + (2\gamma_o - 1) e_{io}^\beta] (x_{io}^m + (2\gamma_o - 1) x_{io}^\beta) z_i, \quad \forall i = 1, \dots, t \\
p_j + q_{ij} &\geq [e_{ij}^m + (2\gamma_j - 1) e_{ij}^\beta] (y_{ij}^m + (2\gamma_j - 1) y_{ij}^\beta) z_i, \quad \forall i = t+1, \dots, t+s, \forall j = 1, \dots, n \\
p_j + q_{ij} &\geq [e_{ij}^m + (2\gamma_j - 1) e_{ij}^\beta] (x_{ij}^m - (2\gamma_j - 1) x_{ij}^\alpha) z_i, \quad \forall i = 1, \dots, t, \forall j = 1, \dots, n
\end{aligned} \tag{35}$$

$$\begin{aligned}
& -z_i \leq w_i \leq z_i, \quad \forall i = 1, 2, \dots, t + s \\
& |w_B - a_{Bi} w_i| \leq \xi, \quad i = 1, \dots, t + s \\
& |w_i - a_{iW} w_W| \leq \xi, \quad i = 1, \dots, t + s \\
& p_j, q_{ij} \geq 0, \quad \forall i, j \\
& w_i \geq 0, \quad i = 1, \dots, t + s
\end{aligned} \tag{35}$$

In models (34) and (35), the constraints belong to both BWM and RCDEA models. Also, the constraint

$$|w_B - a_{Bi} w_i| \leq \xi \text{ is easily transformed to two linear constraints } w_B - a_{Bi} w_i \leq \xi \text{ and } a_{Bi} w_i - w_B \geq \xi.$$

Lemma: The models (34) and (35) are feasible.

Proof: Without loss of generality, it is assumed that the $w_1 = z_1 = 1$ and other variables are equal to zero except q_{1j}

. If we find a feasible value for q_{1j} , then the model (34) is feasible. Since $w_1 = z_1 = 1$, the constraints of the model

(34) is as follows:

The first constraint will be as $-\theta_j^{CCR} (x_{1j}^m + (1 - 2 \times (0.4))x_{1j}^\beta) + q_{1j} \leq 0, j = 1, \dots, n$. It is assumed that the x_{1j}^α and

x_{1j}^β are 0.05 of the x_{1j}^m . Also, the e_{1j}^α and e_{1j}^β are 0.05 of the e_{1j}^m . Hence, the first constraint can be written as

$q_{1j} \leq 1.01x_{1j}^m \times \theta_j^{CCR}$. In addition, the 4th and the 6th constraints will be $q_{1j} \geq 0.99e_{1j}^m \times 0.99x_{1j}^m$ and

$q_{1j} \geq 0.99e_{1j}^m \times 1.01x_{1j}^m$. Therefore $0.99e_{1j}^m \times 1.01x_{1j}^m \leq q_{1j} \leq 1.01x_{1j}^m \times \theta_j^{CCR}$. Since e_{1j}^m is the perturbation

percentage (for instance $e_{1j}^m = 0.05$) and it is a small value, then, we can find a feasible value for $q_{1j} \geq 0$. Other

constraints including the constraints related to BWM are feasible. In a similar way, the model (35) is feasible, too.

The models (34) and (35) are bi-objective programming models which can be solved via some multiple objective

programming models such as parametric method, constraint- ε , min-max approach, goal programming and etc. In this

paper, the min-max approach has been used for solving models (34) and (35). The min-max approach for bi-objectives

of model (34) is as follows:

$$\begin{aligned}
& \min \max \{ (f_1^* - [\sum_{i=t+1}^{t+s} w_i (y_{io}^m + (1 - 2\gamma_o) y_{io}^\beta) - \theta_o^{CCR} \sum_{i=1}^t w_i (x_{io}^m - (1 - 2\gamma_o) x_{io}^\alpha)] \\
& - \Gamma_o P_o - \sum_{i=t+1}^{t+s} q_{io} - \sum_{i=1}^t q_{io}), (f_2^* - [-\xi]) \}
\end{aligned} \tag{36}$$

where f_1^* and f_2^* are the ideal values of first and second objective functions. For calculating f_1^* and f_2^* , the objective functions f_1 and f_2 are optimized on the constraints of model (34), separately. In the other words, for obtaining the ideal value for f_1 , the objective function f_2 is eliminated from the model (34) and the model is solved by using the objective function f_1 . Also, the ideal value for f_2 is obtained by eliminating the objective function f_1 from the model (34). It is clear that the min-max (36) is easily converted to a linear model as follows:

min κ

s.t :

$$f_1^* - [\sum_{i=t+1}^{t+s} w_i (y_{io}^m + (1-2\gamma_o)y_{io}^\beta) - \theta_o^{CCR} \sum_{i=1}^t w_i (x_{io}^m - (1-2\gamma_o)x_{io}^\alpha) - \Gamma_o p_o - \sum_{i=t+1}^{t+s} q_{io} - \sum_{i=1}^t q_{io}] \leq \kappa$$

$$f_2^* - [-\xi] \leq \kappa$$

$$\sum_{i=t+1}^{t+s} w_i (y_{ij}^m - (1-2\gamma_j)y_{ij}^\alpha) - \theta_j^{CCR} \sum_{i=1}^t w_i (x_{ij}^m + (1-2\gamma_j)x_{ij}^\beta) + \Gamma_j p_j + \sum_{i=t+1}^{t+s} q_{ij} + \sum_{i=1}^t q_{ij} \leq 0, j=1, \dots, n$$

$$\sum_{i=1}^{m+s} w_i = 1$$

$$p_o + q_{io} \geq [e_{io}^m - (1-2\gamma_{io})e_{io}^\alpha](y_{io}^m + (1-2\gamma_o)y_{io}^\beta)z_i, \quad \forall i = t+1, \dots, t+s \quad (37)$$

$$p_o + q_{io} \geq [e_{io}^m - (1-2\gamma_{io})e_{io}^\alpha](x_{io}^m - (1-2\gamma_o)x_{io}^\alpha)z_i, \quad \forall i = 1, \dots, t$$

$$p_j + q_{ij} \geq [e_{ij}^m - (1-2\gamma_{ij})e_{ij}^\alpha](y_{ij}^m - (1-2\gamma_j)y_{ij}^\alpha)z_i, \quad \forall i = t+1, \dots, t+s, \forall j = 1, \dots, n$$

$$p_j + q_{ij} \geq [e_{ij}^m - (1-2\gamma_{ij})e_{ij}^\alpha](x_{ij}^m + (1-2\gamma_j)x_{ij}^\beta)z_i, \quad \forall i = 1, \dots, t, \forall j = 1, \dots, n$$

$$-z_i \leq w_i \leq z_i, \quad \forall i = 1, 2, \dots, t+s$$

$$|w_B - a_{Bi} w_i| \leq \xi, \quad i = 1, \dots, t+s$$

$$|w_i - a_{iW} w_W| \leq \xi, \quad i = 1, \dots, t+s$$

$$p_j, q_{ij} \geq 0, \quad \forall i, j$$

$$w_i \geq 0, \quad i = 1, \dots, t+s$$

The linear BWM- RCDEA model (37) is solved and the optimal weights ($W_1^*, \dots, W_m^*, W_{m+1}^*, \dots, W_{m+s}^*$) are obtained for inputs and outputs. Then, the efficiency score of DMU_o for $\gamma \leq 0.5$ is calculated by equation (38).

$$\theta_{o,\gamma_o \leq 0.5}^{BWM-RCDEA} = \frac{\sum_{i=m+1}^{m+s} w_i^* [y_{io}^m + (1-2\gamma_o)y_{io}^\beta]}{\sum_{i=1}^m w_i^* [x_{io}^m - (1-2\gamma_o)x_{io}^\alpha]} \quad (38)$$

According to the above way, for $\gamma > 0.5$, the bi-objective model (35) can be transferred to linear models and the efficiency score of DMU_o can be calculated as follows:

$$\theta_{o,\gamma_o > 0.5}^{BWM-RCDEA} = \frac{\sum_{i=m+1}^{m+s} w_i^* [y_{io}^m - (2\gamma_o - 1)y_{io}^\alpha]}{\sum_{i=1}^m w_i^* [x_{io}^m + (2\gamma_o - 1)x_{io}^\beta]} \quad (39)$$

4. Illustrative Application

In this section, data of 39 electricity distribution companies in Iran is collected and companies are evaluated through the proposed BWM-RCDEA model. Iranian distribution companies are run under supervision of TAVANIR Company (Iran Power Generation, Transmission and Distribution Management Company). TAVANIR was established in 1992 and is responsible for development of electricity power generation, transmission and distribution in Iran. Performance assessment of electricity distribution companies is very important from variety of aspects, especially service to customers and their satisfaction, selling electricity as an economic commodity at a lower cost, development of distribution network for covering the vast surface of the country and reduce electricity failure.

In order to evaluate efficiency of electricity distribution companies, first input and output variables must be selected. Since there is no firm consensus on which variables best describe the operation of distribution utilities, in literature, different variables have been selected. For instance, Petridis et al. (2019) evaluated 20 Turkish electric distribution companies using a network DEA model with criteria of number of staff, net consumption, length of cables, installed capacity as inputs, energy supply as intermediate and energy loses, annual faults and disruption, number of customers and number of towns/villages as desirable and undesirable output variables. Çelen and Aydin (2012) evaluated 21 electricity distribution companies in Turkey. They applied number of employees, length of the distribution line and transformers capacity as inputs and quality of service, number of customers and quantity delivered as outputs. Also, Çelen (2013) evaluated 21 distribution utilities in Turkey between years 2002- 2009. Variables of number of

employees, length of the distribution line, transformer capacity and quality of electricity considered as inputs and variables of electricity delivered and number of customers as outputs. Tavassoli et al. (2015) selected the number of employees, network length and transmission capacity as inputs and unit delivery and service area as outputs of their research. Moreover, Azadeh et al. (2015) evaluated Iranian distribution companies using network length, transport capacity and the number of employees as inputs while number of customers and total electricity sales are chosen as stochastic outputs. In other research for evaluating Iranian distribution units, Sadjadi and Omrani (2008) adopted number of employees, transformers capacity and network length as inputs and number of customers and total electricity sales as outputs. And finally, Omrani et al., (2020) for evaluating 39 electricity distribution companies in Iran, used four inputs of number of employees, service area network length, and, transformers capacity and three outputs of number of customers, energy delivery, and total electricity sales.

As it can be seen, different variables have been selected by different researchers. In this study, the most frequent variables in the literature are selected. For this purpose, four inputs including service area (km^2), number of employees, transformers capacity (Mega Volt Amp) and network length (km) and three outputs including energy delivery (million KWh), number of customers (*1000) and total electricity sales (million KWh) are selected. Our data series involves annual data on 39 companies observed in 2017. These data are retrieved from Iran Ministry of Energy, TAVANIR Company (www.tavanir.org.ir) and presented in Table 2. It should be noted that since robust optimization is sensitive to normalization, data are normalized and presented in interval [0, 1].

-----[Table 2 about here] -----

5. Results and Discussions

In this section the result of applying the proposed model has been presented. Indeed, electricity companies' efficiency scores evaluated in three phases. In the first phase, a DEA model has been applied to estimate the efficiency scores of companies. Then considering data as fuzzy sets, and considering perturbation in constructing fuzzy sets, RCDEA model with and without fuzzy perturbation in fuzzy inputs and fuzzy outputs results are calculated. Finally, in the third phase, DMs' preferences and judgments on input and output variables are incorporated into the RCDEA models and BWM-RCDEA models are generated. The results of implementing each phase of the proposed model are discussed below.

5.1. DEA Results

The results of conventional DEA model are presented in Table 3 and Figure 1. According to the results, the efficiencies of 10 companies are equal to one. In other words, these 10 companies technically are efficient in converting inputs to outputs and form the efficient frontier. On the other hand, companies with efficiency scores less than 1 are inefficient and among inefficient units, Ilam has the worst performance with the score of 0.6351. Also, Gharb-e-Mazandaran with the score of 0.6685 and Golestan with the score of 0.6750 are the second and the third worst performance units respectively. As it can be observed from the Table 3, DEA is unable to rank all DMUs and low distinguish power of DEA is the weakness of this popular model to evaluate efficiency scores. Moreover, DEA uses only data to calculate efficiencies, and DMUs are free to find the best combination of input and output variables for reaching the efficient frontier. Finally, the mean of efficiency scores is 0.8681 which is more than both RCDEA and BWM-RCDEADEA results.

-----[Table 3 about here] -----

-----[Fig 1 about here] -----

5.2. RCDEA Results

RCDEA results has been shown in the Table 3. Initially, in order to obtain RCDEA results, some parameters should be set. First, without losing any generality, it is assumed that $\gamma = \gamma_i = \gamma_r$. So, for different γ ($\gamma \leq 0.5$ and $\gamma > 0.5$) either model (23) or (26) can be applied. In this paper, proposed model is implemented for $\gamma = 0.4$ and $\gamma = 0.8$ respectively. The mean value of *L-R* fuzzy numbers (*m*) are considered as data reported in Table 2 after normalizing. In addition, the left and right spreads (α, β) are 0.05 of the mean value. It is assumed that in determining mean values, there is 0.03 perturbation, and this uncertainty degree is exact. Furthermore, to ensure full protection of model, budget of uncertainty parameter, Γ , is set equal 7 for all constraints. Indeed, model is immune if 100% of uncertain parameters take their worst-case value.

According to the results for $\gamma = 0.4$, although all scores get reduced than DEA model but ranking has not change tangibly and is almost the same with the DEA model. Since it is considered the same level of robustness and fuzziness for all data, hence, efficiency scores decreased by almost the same proportion. For example, units with the score of 1, in a new evaluation get the score of 0.9802 and like DEA model 10 companies have the best performance. On the

other hand, on the bottom of the Table 3, Ilam, Gharb-e-Mazandaran and Golestan with the scores of 0.6115, 0.6452 and 0.6518 have the weakest performance, respectively. Moreover, the mean of all scores is 0.8425, which implies that model in order to keep constraints immune against uncertainty, releases the optimal solutions and scores get lower values to keep the model feasible.

Also, RCDEA model is implemented for $\gamma = 0.8$ and efficiencies are evaluated using model (31). According to the results, scores are lower than both DEA and RCDEA for $\gamma = 0.4$. However, units' ranking has not changed significantly and is almost the same with previous two models. For example, top efficiency score is 0.8352 for 10 companies and the lowest score is 0.5210 which belongs to the Ilam. The mean of scores is 0.7173, which is lower than both DEA and RCDEA for $\gamma = 0.4$.

5.3. RCDEA with Fuzzy Perturbation Results

To observe the impact of uncertainty on the perturbation level, the results of RCDEA model are recalculated considering fuzzy perturbation. The results are presented in the Table 3 and Figure 1. According to the results, for $\gamma \leq 0.5$ with fuzzy perturbation, efficiency scores show slight increase, and for $\gamma > 0.5$ with fuzzy perturbation scores indicate slight decrease in comparison with RCDEA models with exact perturbation. For instance, for $\gamma \leq 0.5$ with fuzzy perturbation, best companies get the score of 0.9808 which is more than $\gamma \leq 0.5$ with exact perturbation model results. Also, the mean of efficiency scores is 0.8467. The reason for increasing efficiency scores is that considering fuzzy data for perturbation values lead to decrease in value of perturbation and resulted in increase in efficiency scores. In the other words, when uncertainty level decreases, model keeps optimal solutions and is closer to conventional deterministic DEA model results.

In contrast, considering fuzzy sets for perturbation value for $\gamma > 0.5$, resulted in decreasing efficiency scores and consequently decrease in the mean of scores to 0.7158. Indeed, model to keep constraints against increase in uncertainty level releases optimal solutions. It should be noted that since uncertainty considered the same for all inputs, outputs and perturbation values, rankings remained almost unchanged in models.

5.4. BWM- RCDEA Results

In order to reduce input and output weights flexibility and to incorporate DMs' judgement into the evaluations, a novel BWM- RCDEA model is applied and efficiency scores for electricity distribution companies are re-calculated. In this regard, a preference vector is designed according to the opinions of a team of experts consist of three members with sufficient experiences at the Azarbaijan Gharbi company. The values are outcome of the mentioned team closely working and are acceptable for all team members. Table 4 presents the best, the worst criteria and preference values of the best criterion over all criteria and preference of all criteria over the worst criterion.

-----[Table 4 about here] -----

After determination of preferences' vector, obtained BWM model is incorporated into the RCDEA model and a bi-objective BWM- RCDEA models (34) and (35) are obtained. Then the bi-objective models are solved with pre-defined parameters (section 4.2, first paragraph) using min-max approach (37) and results are shown in the Table 5 and Figure (2).

-----[Table 5 about here] -----

-----[Fig 2 about here] -----

According to the results for $\gamma = 0.4$ with fuzzy perturbation, Tehran Bozorg and Ahwaz units with the score of 0.9802 have the best performance and are followed by Alborz and Mashhad with efficiency scores of 0.8371 and 0.7814 respectively. In contrary, Khorasan-e- Jonubi with the score of 0.1801 has the worst performance in converting inputs to outputs. Khorasan-e-Shomali with the score of 0.2260 is the second and Jonub-e-Kerman with the score of 0.2554 is the third worst performance among 39 companies. As it can be seen, the proposed model is able to rank DMUs almost fully and has improved the distinguish power of conventional DEA dramatically. Furthermore, the efficiency scores get reduced meaningfully, because we have added new constraints into the proposed BWM- RCDEA model and under such circumstance, results will not improve. Therefore, the mean of scores is 0.4510, which is less than RCDEA model results for $\gamma = 0.4$. Results imply that the preferences of DMs have a significant impact on the efficiencies, since data are normalized and higher weights for variables can subjectively modify the efficiencies.

According to the BWM- RCDEA results for $\gamma = 0.8$, again, Tehran Bozorg and Ahwaz with the efficiency score of 0.8352 has the best performance among 39 companies. Alborz unit with the score of 0.7161 is ranked second and is followed by Mashhad with the score of 0.6703. At the bottom of the ranking Khorasan-e- Jonubi with the score of

0.1539 has the weakest performance. Khorasan-e- Shomali with the score of 0.1923 has the second worst performance and Jonub-e-Kerman with the score of 0.2183 has the third weakest performance among 39 units. The other units' performances get reduced than BWM- RCDEA results for $\gamma = 0.8$ with fuzzy perturbation, however, ranking has not changed dramatically in comparison with BWM-RCDEA model with $\gamma = 0.4$. In overall, distinguish power of DEA has been improved significantly and model is able to immune constraints and objective function against uncertainty obviously. Furthermore, using both data and DMs' preferences in evaluations leads to a more precise and reliable ranking. Like RCDEA model, to observe the fuzziness impact on the perturbation degree results, we have implemented BWM-RCDEA with considering exact values for perturbation and reported results in the Table 5. Results approve that ignoring fuzziness of perturbation lead decrease in scores for $\gamma \leq 0.5$ model in comparison with BWM- RCDEA model with fuzzy perturbation in fuzzy inputs and outputs. Also considering exact value for perturbation for $\gamma > 0.5$ resulted in increase in scores and the mean of scores in comparison with BWM- RCDEA model with fuzzy perturbation. To present the differences of scores generated by different models without and with fuzzy perturbation, the scores are depicted in Figures 3 and 4, respectively.

-----[Fig 3 about here] -----

-----[Fig 4 about here] -----

For investigating the relationship between the results of the DEA, RCDEA, RCDEA with fuzzy perturbation, BWM- RCDEA and BWM-RCDEA with fuzzy perturbation, we perform Pearson correlation test between efficiency scores and Spearman correlation test between ranks. The results for $\gamma = 0.4$ are presented in the Tables (6) and (7). As can be seen in Tables (6) and (7), the both correlations are significant at the 0.01 level.

-----[Table 6 about here] -----

-----[Table 7 about here] -----

5.5. Sensitivity analysis

Here, sensitivity analysis is performed on parameters such as γ , Γ and ℓ . For sensitivity analysis, we consider two scenarios for each parameter and compare the results. For example, for perturbation degree, ℓ , we consider the perturbation equal to 0.05, solve the model and compare the results with our initial 0.03. Likewise, we run the model

for different values of the, γ and Γ . The results are presented in the Tables 8 and 9, respectively. The results indicate that by increasing perturbation degree (ℓ), efficiency scores get lower. Also, by decreasing the budget of uncertain parameters in each constraints (Γ), efficiencies get higher scores and by increasing value for γ in the model (37), scores take lower values. Moreover, by decreasing values for γ in the case $\gamma > 0.5$, for example $\gamma = 0.6$, scores take higher values than RCDEA model for $\gamma = 0.8$.

-----[Table 8 about here] -----

-----[Table 9 about here] -----

6. Recommendations for Policy Makers

In this section, according to the BWM- RCDEA results recommendations have been made for decision makers. In this regard, the average weight of criteria has been calculated after implementing BWM- RCDEA model and results are indicated in the Table 10.

-----[Table 10 about here] -----

According to the Table 10, energy delivery has the least impact on the evaluations and in contrary, total electricity sale is the most important criterion. Units, in order to increase their efficiencies are better to put their attention to sale energy and generate energy from available and cheap resources such as renewable sources. For example, small sized power plants which are closer to customers, not only needs less network length, but also increases the outputs simultaneously. Also, renewing distribution infrastructures to improve energy delivery will lead to increase efficiencies of electricity distribution companies. Specifically, for Khorasan-e-Jonubi, Khorasan-e-Shomali and Jonub-e-Kerman that sale less electricity for customers because of small populations (located in desert areas with less population density), and it is almost impossible to increase their outputs substantially, decreasing the number of employers and network length by constructing small sized renewable resources may improve their efficiency.

7. Conclusion

In real competitive markets, performance evaluation is crucial for companies to know their strengths and weaknesses to set appropriate policies for improving their efficiencies. Distribution companies' performance evaluation is considered crucial issue for regulators. In this study, we applied a novel approach of BWM- RCDEA model to

investigate efficiency scores of 39 electricity distribution companies in Iran. Indeed, beside of considering uncertainty in data as fuzzy sets and uncertainty in constructing fuzzy sets as robust optimization with fuzzy perturbation, DMs' preferences were incorporated into DEA model, too. The proposed bi-objective model was solved using min-max approach. As results, the proposed model improves the distinguish power of DEA and results are more reliable and precise than conventional DEA. Also, recommendations were proposed for decision makers to improve units' efficiencies. In summary, due to the importance of total electricity sale criterion on the efficiency evaluations, generating energy from convenient and cheap resources may lead to increase in electricity sale and consequently improving performances. For future studies, the BWM model can be combined with other DEA models such as super-efficiency DEA, additive models and etc. Also, researchers can consider DMs' judgments as fuzzy numbers and fuzzy BWM model can be combined with the DEA models. Also, developing group BWM-DEA for considering the preferences of multiple DMs is very important in this field. Finally, for dealing with real worlds problem with integer variables, such as number of customers, the mixed integer model of BWM-RCDEA could be applied.

Acknowledgement

The article was prepared within the framework of the Basic Research Program at HSE University. The authors would like to thank the Editor and the Guest Editor of *Annals of Operations Research*, and three anonymous reviewers for their insightful comments and suggestions. As results this paper has been improved substantially.

References

- Alizadeh, A., & Omrani, H. (2019). An integrated multi response Taguchi-neural network-robust data envelopment analysis model for CO2 laser cutting. *Measurement*, *131*, 69-78.
- Allen, R., Athanassopoulos, A., Dyson, R. G., & Thanassoulis, E. (1997). Weights restrictions and value judgments in data envelopment analysis: evolution, development and future directions. *Annals of Operations Research*, *73*, 13-34.
- Amini, M., Dabbagh, R., Omrani, H., (2019). A fuzzy data envelopment analysis based on credibility theory for estimating road safety. *Decision Science Letter*. DOI: 10.5267/j.dsl.2019.1.001

- Ashrafi, A., & Kaleibar, M. M. (2017). Cost, revenue and profit efficiency models in generalized fuzzy data envelopment analysis. *Fuzzy Information and Engineering*, 9(2), 237-246.
- Azadeh, A., Haghighi, S. M., Zarrin, M., & Khaefi, S. (2015). Performance evaluation of Iranian electricity distribution units by using stochastic data envelopment analysis. *International Journal of Electrical Power & Energy Systems*, 73, 919-931.
- Banker, R.D. Charnes, A. Cooper W.W. (1984). Some methods for estimating technical and scale inefficiencies in data envelopment analyses *Management Science*, 30(9): 1078-1092.
- Ben-Tal, A. Nemirovski, A. (2000). Robust solutions of linear programming problems contaminated with uncertain data. *Mathematical Programming*, 88(3):411–424.
- Bertsimas, D. Sim, M. (2004). The price of robustness, *Operations Research*, Vol.52, No. 1, pp.35–53.
- Çelen, A. (2013). Efficiency and productivity (TFP) of the Turkish electricity distribution companies: An application of two-stage (DEA&Tobit) analysis. *Energy Policy*, 63, 300-310.
- Çelen, A., & Yalçın, N. (2012). Performance assessment of Turkish electricity distribution utilities: An application of combined FAHP/TOPSIS/DEA methodology to incorporate quality of service. *Utilities Policy*, 23, 59-71.
- Charnes, A., Cooper, W. W., & Rhodes, E. (1978). Measuring the efficiency of decision making units. *European journal of operational research*, 2(6), 429-444.
- Charnes, A., Cooper, W. W., Huang, Z. M. (1990). Polyhedral cone-ratio DEA models with an illustrative application to large commercial banks. *Journal of econometrics*, 46(1-2), 73-91.
- Chen, Y., Cook, W. D., & Lim, S. (2019). Preface: DEA and its applications in operations and data analytics. *Annals of Operations Research*, 278(1), 1-4.
- Dyson, R. G., & Thanassoulis, E. (1988). Reducing weight flexibility in data envelopment analysis. *Journal of the Operational Research Society*, 39(6), 563-576.
- Degl'Innocenti, M., Kourtzidis, S. A., Sevic, Z., & Tzeremes, N. G. (2017). Investigating bank efficiency in transition economies: A window-based weight assurance region approach. *Economic Modelling*, 67, 23-33.
- Dubois, D., & Prade, H. (1978). Operations on fuzzy numbers. *International Journal of Systems Sciences* 9(6):613–626

- Ebrahimi, B., & Khalili, M. (2018). A new integrated AR-IDEA model to find the best DMU in the presence of both weight restrictions and imprecise data. *Computers & Industrial Engineering*, 125, 357-363.
- Emrouznejad, A., & Yang, G. L. (2018). A survey and analysis of the first 40 years of scholarly literature in DEA: 1978–2016. *Socio-Economic Planning Sciences*, 61, 4-8.
- Emrouznejad, A and M. Tavana (2014) Performance Measurement with Fuzzy Data Envelopment Analysis, *Studies in Fuzziness and Soft Computing* 309, Springer-Verlag.
- Emrouznejad, A., M. Tavana, A. Hatami-Marbini (2014) The State of the Art in Fuzzy Data Envelopment Analysis”, in Performance Measurement with Fuzzy Data Envelopment Analysis published in *Studies in Fuzziness and Soft Computing* 309: 1:48, Springer-Verlag.
- Fasanghari, M., Amalnick, M. S., Anvari, R. T., & Razmi, J. (2015). A novel credibility-based group decision making method for Enterprise Architecture scenario analysis using Data Envelopment Analysis. *Applied Soft Computing*, 32, 347-368.
- Guo, P. (2009). Fuzzy data envelopment analysis and its application to location problems. *Information Sciences*, 179(6), 820-829.
- Guo, P., Tanaka, H., 2001. Fuzzy DEA: a perceptual evaluation method. *Fuzzy Sets and Systems* 119 (1), 149-160.
- Hatami-Marbini, A., Emrouznejad, A., & Tavana, M. (2011). A taxonomy and review of the fuzzy data envelopment analysis literature: two decades in the making. *European journal of operational research*, 214(3), 457-472.
- Hatami-Marbini, A., M. Tavana, A. Emrouznejad and S. Saati (2012). Efficiency measurement in fuzzy additive Data Envelopment Analysis. *International Journal of Industrial and Systems Engineering* 10(1): 1-20.
- Hatami-Marbini, A., & Saati, S. (2018). Efficiency evaluation in two-stage data envelopment analysis under a fuzzy environment: A common-weights approach. *Applied Soft Computing*, 72, 156-165.
- Ignatius, J., Ghasemi, M. R., Zhang, F., Emrouznejad, A., & Hatami-Marbini, A. (2016). Carbon efficiency evaluation: An analytical framework using fuzzy DEA. *European Journal of Operational Research*, 253(2), 428-440.

- Khalili, M., Camanho, A. S., Portela, M. C. A. S., & Alirezaee, M. R. (2010). The measurement of relative efficiency using data envelopment analysis with assurance regions that link inputs and outputs. *European Journal of Operational Research*, 203(3), 761-770.
- Kim, J. H., Kim, W. C., & Fabozzi, F. J. (2018). Recent advancements in robust optimization for investment management. *Annals of Operations Research*, 266(1), 183-198.
- Lai, P. L., Potter, A., Beynon, M., & Beresford, A. (2015). Evaluating the efficiency performance of airports using an integrated AHP/DEA-AR technique. *Transport Policy*, 42, 75-85.
- Lee, S., Cho, Y., & Ko, M. (2020). Robust Optimization Model for R&D Project Selection under Uncertainty in the Automobile Industry. *Sustainability*, 12(23), 10210.
- Lertworasirikul S., Fang SC., Joines J.A., Nuttle H.L.W. (2003a) Fuzzy Data Envelopment Analysis: A Credibility Approach. In: Verdegay JL. (eds) Fuzzy Sets Based Heuristics for Optimization. Studies in Fuzziness and Soft Computing, vol 126. Springer, Berlin, Heidelberg
- Lertworasirikul, S., Fang, S. C., Joines, J. A., & Nuttle, H. L. (2003b). Fuzzy data envelopment analysis (DEA): a possibility approach. *Fuzzy sets and Systems*, 139(2), 379-394.
- Li, X., & Liu, B. (2006). A sufficient and necessary condition for credibility measures. *International Journal of Uncertainty, Fuzziness and Knowledge-Based Systems*, 14(5), 527-535.
- Liu, S. T. (2014). Restricting weight flexibility in fuzzy two-stage DEA. *Computers & Industrial Engineering*, 74, 149-160.
- Liu, S. T. (2008). A fuzzy DEA/AR approach to the selection of flexible manufacturing systems. *Computers & Industrial Engineering*, 54(1), 66-76.
- Liu, B., & Liu, Y. K. (2002). Expected value of fuzzy variable and fuzzy expected value models. *IEEE Transactions on Fuzzy Systems*, 10(4), 445-450.
- Liu, Y. K., & Liu, B. (2003). Fuzzy random variables: A scalar expected value operator. *Fuzzy Optimization and Decision Making*, 2(2), 143-160.

- Ma, Z., & Cui, W. (2014). Generalized fuzzy data envelopment analysis methods. *Applied Soft Computing*, 19, 215-225.
- Mi, X., Tang, M., Liao, H., Shen, W., & Lev, B. (2019). The state-of-the-art survey on integrations and applications of the best worst method in decision making: Why, what, what for and what's next?. *Omega*. <https://doi.org/10.1016/j.omega.2019.01.009>
- Omrani, H. (2013). Common weights data envelopment analysis with uncertain data: A robust optimization approach. *Computers & Industrial Engineering*, 66(4), 1163-1170.
- Omrani, H., Adabi, F., & Adabi, N. (2017). Designing an efficient supply chain network with uncertain data: a robust optimization—data envelopment analysis approach. *Journal of the Operational Research Society*, 68(7), 816-828.
- Omrani, H., Alizadeh, A., & Emrouznejad, A. (2018). Finding the optimal combination of power plants alternatives: A multi response Taguchi-neural network using TOPSIS and fuzzy best-worst method. *Journal of Cleaner Production*, 203, 210-223.
- Omrani, H., Alizadeh, A., & Naghizadeh, F. Incorporating decision makers' preferences into DEA and common weight DEA models based on the best–worst method (BWM). *Soft Computing*, In press. DOI: 10.1007/s00500-019-04168-z.
- Peykani, P., E. Mohammadi, A. Emrouznejad, M. S. Pishvae, M. Rostamy-Malkhalifeh (2019). Fuzzy Data Envelopment Analysis: An Adjustable Approach, *Expert Systems with Applications*, In Press. DOI: 10.1016/j.eswa.2019.06.039
- Petridis, K., Ünsal, M. G., Dey, P., & Örkücü, H. H. (2019). A novel network data envelopment analysis model for performance measurement of Turkish electric distribution companies. *Energy*.
- Rezaei, J. (2015). Best-worst multi-criteria decision-making method. *Omega*, 53, 49-57.
- Rezaei, J. (2016). Best-worst multi-criteria decision-making method: Some properties and a linear model. *Omega*, 64, 126-130.
- Roll, Y., Cook, W. D., & Golany, B. (1991). Controlling factor weights in data envelopment analysis. *IIE transactions*, 23(1), 2-9.

- Sadjadi, S. J., & Omrani, H. (2008). Data envelopment analysis with uncertain data: An application for Iranian electricity distribution companies. *Energy Policy*, 36(11), 4247-4254.
- Sadjadi, S. J., Omrani, H., Abdollahzadeh, S., Alinaghian, M., & Mohammadi, H. (2011a). A robust super-efficiency data envelopment analysis model for ranking of provincial gas companies in Iran. *Expert Systems with Applications*, 38(9), 10875-10881.
- Sadjadi, S. J., Omrani, H., Makui, A., & Shahanaghi, K. (2011b). An interactive robust data envelopment analysis model for determining alternative targets in Iranian electricity distribution companies. *Expert Systems with Applications*, 38(8), 9830-9839.
- Saeidi-Mobarakeh, Z., Tavakkoli-Moghaddam, R., Navabakhsh, M., & Amoozad-Khalili, H. (2020). A bi-level and robust optimization-based framework for a hazardous waste management problem: A real-world application. *Journal of Cleaner Production*, 252, 119830.
- Salahi, M., Torabi, N., & Amiri, A. (2016). An optimistic robust optimization approach to common set of weights in DEA. *Measurement*, 93, 67-73.
- Sarıca, K., & Or, I. (2007). Efficiency assessment of Turkish power plants using data envelopment analysis. *Energy*, 32(8), 1484-1499.
- Sengupta, J. K. (1992). A fuzzy systems approach in data envelopment analysis. *Computers & Mathematics with Applications*, 24(8-9), 259-266.
- Shabanpour, H., Yousefi, S., & Saen, R. F. (2017). Future planning for benchmarking and ranking sustainable suppliers using goal programming and robust double frontiers DEA. *Transportation Research Part D: Transport and Environment*, 50, 129-143.
- Soleimani-Damaneh, M. (2008). Fuzzy upper bounds and their applications. *Chaos, Solitons & Fractals*, 36(2), 217-225.
- Soltanzadeh, E., & Omrani, H. (2018). Dynamic network data envelopment analysis model with fuzzy inputs and outputs: an application for Iranian Airlines. *Applied Soft Computing*, 63, 268-288.

- Soyster, A.L.(1973). Convex programming with set-inclusive constraints and applications to inexact linear programming. *Operation Research*, 21, 1154–1157.
- Tavana, M., Khanjani Shiraz, R., Hatami-Marbini, A., Agrell, P.J., & Paryab, K. (2012). Fuzzy stochastic data envelopment analysis with application to base realignment and closure (BRAC). *Expert Systems with Applications*, 39, 12247-12259.
- Tavassoli, M., Faramarzi, G. R., & Saen, R. F. (2015). Ranking electricity distribution units using slacks-based measure, strong complementary slackness condition, and discriminant analysis. *International Journal of Electrical Power & Energy Systems*, 64, 1214-1220.
- Thompson, R. G., Singleton Jr, F. D., Thrall, R. M., & Smith, B. A. (1986). Comparative site evaluations for locating a high-energy physics lab in Texas. *Interfaces*, 16(6), 35-49.
- Toloo, M., & Mensah, E. K. (2019). Robust optimization with nonnegative decision variables: a DEA approach. *Computers & Industrial Engineering*, 127, 313-325.
- Triantis, K., & Girod, O. (1998). A mathematical programming approach for measuring technical efficiency in a fuzzy environment. *Journal of Productivity Analysis*, 10(1), 85-102.
- Vafadarnikjoo, A., Tavana, M., Botelho, T., & Chalvatzis, K. (2020). A neutrosophic enhanced best–worst method for considering decision-makers’ confidence in the best and worst criteria. *Annals of Operations Research*, 289(2), 391-418.
- Wang, Y. M., Chin, K. S., & Poon, G. K. K. (2008). A data envelopment analysis method with assurance region for weight generation in the analytic hierarchy process. *Decision Support Systems*, 45(4), 913-921.
- Yin, H., Yu, D., Yin, S., & Xia, B. (2018). Possibility-based robust design optimization for the structural-acoustic system with fuzzy parameters. *Mechanical Systems and Signal Processing*, 102, 329-345.
- Zadeh, L. A. (1978). Fuzzy sets as a basis for a theory of possibility. *Fuzzy sets and systems*, 1(1), 3-28.
- Zhao, H., Guo, S., & Zhao, H. (2019). Comprehensive assessment for battery energy storage systems based on fuzzy-MCDM considering risk preferences. *Energy*, 168, 450-461.
- Zimmermann, H. J. (1978). Fuzzy programming and linear programming with several objective functions. *Fuzzy sets and systems*, 1(1), 45-55.

Zimmermann, H. J. (2001). *Fuzzy sets theory and its applications* (fourth ed.). Dordrecht, The Netherlands: Kluwer Academic Publishers.

Zohrehbandian, M., Makui, A., & Alinezhad, A. (2010). A compromise solution approach for finding common weights in DEA: An improvement to Kao and Hung's approach. *Journal of the Operational Research Society*, 61(4), 604-610.




Figure1. The scores generated by RCDEA model




Figure 2. The score generated by BWM- RCDEA model




Figure 3. The scores generated by different models with exact perturbation value ($e=0.03$)




Figure 4. The scores generated by different models with fuzzy perturbation value

Table 1. Parameters applied for the models in this study

Parameter	Definition
w_i	Weight of input and output variables
$(x_{ij}^m, x_{ij}^\alpha, x_{ij}^\beta), (y_{rj}^m, y_{rj}^\alpha, y_{rj}^\beta)$	i^{th} fuzzy input and r^{th} fuzzy output for j^{th} DMU, respectively
Γ_0, Γ_j	Budget of uncertainty for objective function and j^{th} constraint respectively
p_0, p_j	Dual variable for objective function and constraints in robust counterpart model
q_{ij}	Dual variables for constructing robust model in Bertsimas and Sim (2004) approach
z_i	Linearization variable in Bertsimas and Sim (2004) approach
$(e_{ij}^m, e_{ij}^\alpha, e_{ij}^\beta)$	Fuzzy perturbation percentage
a_{Bi}, a_{iW}	The best and the worst preference values of DM
γ	The confidence level for credibility theory $\gamma \in [0, 1]$

Table 2. Data of electricity distribution companies for year 2017

DMU	Company	Inputs				Outputs		
		no.	name	Service area	Number of employees	Transformer capacity	Network length	Energy delivery
1	Tabriz	4770	568	1864.3	9085.2	4321	957.8	3962.9
2	Azarbayejan Sharghi	40722	619	1908.0	24075	3613	815.7	3255.9
3	Azarbayejan Gharbi	37412	876	2535.2	27159.1	5564	1207.6	4905.7
4	Ardabil	17880	364	920.3	13482	1932	532.7	1682.1
5	Esfahan	91000	488	5501.0	38584.8	10216	1400.5	9424.3
6	Ostan-e-Esfahan	15706	304	3121.0	14769	5892	1135.6	5499.2
7	Chaharmahal-o-Bakhtiari	16411	180	1059.0	11448	1878.5	350.8	1654.2
8	Markazi	29127	352	2371.3	19990.2	5093.2	697.2	4605.1
9	Hamedan	19368	336	2234.1	18353.9	3613	718.9	3230.2
10	Lorestan	28306	313	1849.3	15766	3250.6	616.2	2935.0
11	Alborz	5142	355	2949.6	12723.6	7193	1276.0	6252.9
12	Tehran Bozorg	1011	1601	11549.0	32061	21886.1	4530.2	20381.1
13	Ostan-e-Tehran	13029	637	9284.8	33802.1	14473	2200.5	12167.5
14	Ghom	11240	230	1870.2	7687	3717	541.5	3337.0
15	Mashhad	3863	347	3655.1	15946	7247	1498.6	6797.4
16	Khorasan-e-Razavi	116794	579	3262.2	41129.9	7975	1224.1	7109.7
17	Khorasan-e-Jonubi	150800	268	1002.0	18174	1673	355.2	1551.9
18	Khorasan-e-Shomali	28434	186	814.6	11255.3	1351	338.9	1225.6
19	Ahwaz	7174	430	4585.4	10259	10774	560.1	9207.3
20	Khouzestan	57945	457	8088.0	31223	18530	1031.6	16075.1
21	Kohkiluye-o-Boyerahmad	16249	186	1168.8	8455.5	1998	242.0	1730.5
22	Zanjan	22164	259	1472.2	13922.9	3697	428.3	3395.6
23	Ghazvin	15627	248	1918.8	12236.4	4573	581.2	4241.1
24	Semnan	97491	143	1355.9	11285.1	2686	365.9	2510.3
25	Sistan-o-Balouchestan	187502	579	2785.1	36774.5	6146	777.3	5298.4
26	Kermanshah	25009	337	2040.8	18406.9	3563	743.0	3020.0
27	Kordestan	29137	271	1398.0	16010	2357	621.8	2123.2
28	Ilam	20150	107	1034.0	7558	1576	214.1	1335.5
29	Shiraz	20148	480	4261.7	25061	6750	1058.6	6188.8
30	Fars	102568	393	4224.0	36435	8260	915.3	7614.5
31	Boushehr	23168	187	3776.0	14583	6905	443.8	6109.6
32	Shomal-e-Kerman	91193	394	2130.2	19545.3	4407	601.3	3915.7
33	Jonub-e-Kerman	95887	415	3049.0	33572.3	6325	531.2	5538.0
34	Gilan	14044	675	3161.3	28442.73	5839	1378.6	5200.6
35	Mazandran	14732	586	3873.0	27648.7	6870	1328.6	5952.2
36	Gharb-e-Mazandaran	9024	281	1904.5	11349.5	2541	569.2	2100.5
37	Golestan	20437	486	2318.2	15025.5	3801	700.6	3316.3
38	Hormozgan	70607	354	4409.9	23988	9849	658.4	8621.7
39	Yazd	73873	437	1951.7	18573.2	5211	643.5	4845.7

Table 3. Results of DEA and RCDEA with and without fuzzy perturbation in fuzzy inputs and outputs

DMU	DEA	$\gamma = 0.4$		$\gamma = 0.8$	
		RCDEA with fuzzy perturbation	RCDEA	RCDEA with fuzzy perturbation	RCDEA
Tabriz	1	0.9808	0.9802	0.8337	0.8352
Azarbajejan Sharghi	0.8234	0.8023	0.8017	0.6818	0.6832
Azarbajejan Gharbi	0.941	0.9189	0.9183	0.7809	0.7825
Ardabil	1	0.9808	0.9802	0.8337	0.8352
Esfahan	0.8497	0.8288	0.8283	0.7044	0.7058
Ostan-e-Esfahan	0.9183	0.8982	0.8976	0.7634	0.7649
Chaharmahal-o-Bakhtiari	0.7349	0.7128	0.7123	0.6057	0.6070
Markazi	0.868	0.8473	0.8468	0.7172	0.7185
Hamedan	0.7173	0.6950	0.6945	0.5905	0.5918
Lorestan	0.7431	0.7211	0.7206	0.6127	0.6140
Alborz	1	0.9808	0.9802	0.8337	0.8352
Tehran Bozorg	1	0.9808	0.9802	0.8337	0.8352
Ostan-e-Tehran	0.9807	0.9613	0.9607	0.6907	0.6919
Ghom	0.8438	0.8229	0.8223	0.6938	0.6950
Mashhad	1	0.9808	0.9802	0.8337	0.8352
Khorasan-e-Razavi	0.963	0.9424	0.9417	0.8001	0.8016
Khorasan-e-Jonubi	0.7231	0.7009	0.7004	0.5955	0.5968
Khorasan-e-Shomali	0.8491	0.8253	0.8247	0.7013	0.7028
Ahwaz	1	0.9808	0.9802	0.8337	0.8352
Khouzestan	1	0.9808	0.9802	0.8337	0.8352
Kohkiluyeh-o-Boyerahmad	0.6792	0.6565	0.6560	0.5577	0.5590
Zanjan	1	0.9808	0.9802	0.8337	0.8352
Ghazvin	1	0.9808	0.9802	0.8337	0.8352
Semnan	0.8768	0.8523	0.8517	0.7242	0.7257
Sistan-o-Balouchestan	0.8377	0.8161	0.8156	0.6919	0.6931
Kermanshah	0.7976	0.7731	0.7725	0.6558	0.6573
Kordestan	0.943	0.9140	0.9133	0.7761	0.7778
Ilam	0.6351	0.6119	0.6115	0.5198	0.5210
Shiraz	0.6856	0.6630	0.6625	0.5600	0.5610
Fars	0.8646	0.8437	0.8431	0.7170	0.7185
Boushehr	0.9688	0.9492	0.9487	0.8069	0.8084
Shomal-e-Kerman	0.8009	0.7795	0.7790	0.6573	0.6628
Jonub-e-Kerman	0.8385	0.8129	0.8123	0.6907	0.6922
Gilan	0.9139	0.8908	0.8901	0.7569	0.7585
Mazandran	0.7639	0.7421	0.7416	0.6306	0.6319
Gharb-e-Mazandaran	0.6685	0.6457	0.6452	0.5485	0.5498
Golestan	0.675	0.6523	0.6518	0.5541	0.5554
Hormozgan	0.9523	0.9326	0.9320	0.7927	0.7942
Yazd	1	0.9808	0.9802	0.8337	0.8352
Mean	0.8681	0.8466	0.8426	0.7158	0.7173

Table 4. Preference values of best criterion over all criteria and all criteria over worst criterion

Criteria	Area service	Number of employees	Transformer capacity	Network length	Energy delivery	Number of customers	Total electricity sale
Best criterion (Total electricity sale)	8	6	4	5	3	2	1
Worst criterion (Area service)	1	3	5	4	6	7	8

Table 5. Results of DEA and BWM- RCDEA models with and without fuzzy perturbation

DMU	DEA	$\gamma = 0.4$		$\gamma = 0.8$	
		BWM- RCDEA with fuzzy perturbation	BWM- RCDEA	BWM- RCDEA with fuzzy perturbation	BWM- RCDEA
Tabriz	1.0000	0.6920	0.6916	0.5903	0.5913
Azarbayejan Sharghi	0.8234	0.2666	0.2665	0.2272	0.2276
Azarbayejan Gharbi	0.9410	0.3434	0.3432	0.2928	0.2933
Ardabil	1.0000	0.2726	0.2724	0.2320	0.2324
Esfahan	0.8497	0.4185	0.4182	0.3575	0.3581
Ostan-e-Esfahan	0.9183	0.6591	0.6587	0.5616	0.5626
Chaharmahal-o-Bakhtiari	0.7349	0.2749	0.2748	0.2340	0.2344
Markazi	0.8680	0.3893	0.3890	0.3322	0.3328
Hamedan	0.7173	0.3394	0.3392	0.2895	0.2900
Lorestan	0.7431	0.3396	0.3394	0.2889	0.2894
Alborz	1.0000	0.8376	0.8371	0.7149	0.7161
Tehran Bozorg	1.0000	0.9808	0.9802	0.8337	0.8352
Ostan-e-Tehran	0.9807	0.6325	0.6322	0.5424	0.5433
Ghom	0.8438	0.6667	0.6663	0.5564	0.5574
Mashhad	1.0000	0.7819	0.7814	0.6692	0.6703
Khorasan-e-Razavi	0.9630	0.3541	0.3539	0.2737	0.2742
Khorasan-e-Jonubi	0.7231	0.1802	0.1801	0.1537	0.1539
Khorasan-e-Shomali	0.8491	0.2262	0.2260	0.1920	0.1923
Ahwaz	1.0000	0.9808	0.9802	0.8337	0.8352
Khouzestan	1.0000	0.6832	0.6828	0.5837	0.5847
Kohkiluye-o-Boyerahmad	0.6792	0.3142	0.3140	0.2669	0.2674
Zanjan	1.0000	0.3865	0.3863	0.3294	0.3299
Ghazvin	1.0000	0.5455	0.5452	0.4649	0.4657
Semnan	0.8768	0.3832	0.3830	0.3264	0.3270
Sistan-o-Balouchestan	0.8377	0.2703	0.2702	0.2319	0.2322
Kermanshah	0.7976	0.3302	0.3300	0.2814	0.2819
Kordestan	0.9430	0.2911	0.2909	0.2477	0.2482
Ilam	0.6351	0.3044	0.3042	0.2451	0.2456
Shiraz	0.6856	0.4257	0.4254	0.3638	0.3644
Fars	0.8646	0.3705	0.3703	0.2920	0.2925
Boushehr	0.9688	0.5694	0.5691	0.4850	0.4858
Shomal-e-Kerman	0.8009	0.3438	0.3436	0.2934	0.2939
Jonub-e-Kerman	0.8385	0.2556	0.2554	0.2180	0.2183
Gilan	0.9139	0.3892	0.3890	0.3332	0.3338
Mazandran	0.7639	0.4181	0.4178	0.3578	0.3584
Gharb-e-Mazandaran	0.6685	0.3655	0.3653	0.3110	0.3116
Golestan	0.6750	0.3746	0.3744	0.3185	0.3191
Hormozgan	0.9523	0.5213	0.5210	0.4457	0.4464
Yazd	1.0000	0.4233	0.4230	0.3615	0.3621
Mean	0.8681	0.4513	0.4510	0.3829	0.3835

Table 6. Pearson correlation test among different models

	DEA	RCDEA	RCDEA with fuzzy perturbation	BWM-RCDEA	BWM-RCDEA with fuzzy perturbation
DEA	1.000	1.000 ^(*)	1.000 ^(*)	0.513 ^(*)	0.516 ^(*)
RCDEA		1.000	1.000 ^(*)	0.515 ^(*)	0.518 ^(*)
RCDEA with fuzzy perturbation			1.000	0.514 ^(*)	0.517 ^(*)
BWM-RCDEA				1.000	1.000 ^(*)
BWM-RCDEA with fuzzy perturbation					1.000

*. Correlation is significant at the 0.01 level (2-tailed).

Table 7. Spearman correlation test among different models

	DEA	RCDEA	RCDEA with fuzzy perturbation	BWM-RCDEA	BWM-RCDEA with fuzzy perturbation
DEA	1.000	1.000 ^(*)	1.000 ^(*)	0.505 ^(*)	0.502 ^(*)
RCDEA		1.000	1.000 ^(*)	0.505 ^(*)	0.502 ^(*)
RCDEA with fuzzy perturbation			1.000	0.505 ^(*)	0.502 ^(*)
BWM-RCDEA				1.000	1.000 ^(*)
BWM-RCDEA with fuzzy perturbation					1.000

*. Correlation is significant at the 0.01 level (2-tailed).

Table 8. Sensitivity analysis for RCDEA with $\gamma \leq 0.5$

DMU	RCDEA			
	Initial RCDEA	$e = 0.05$	$\Gamma = 5$	$\gamma = 0.3$
Tabriz	0.9802	0.9417	0.9802	1.0202
Azarbayejan Sharghi	0.8017	0.7666	0.8017	0.8345
Azarbayejan Gharbi	0.9183	0.8781	0.9183	0.9558
Ardabil	0.9802	0.9417	0.9802	1.0202
Esfahan	0.8283	0.7927	0.8283	0.8621
Ostan-e-Esfahan	0.8976	0.8607	0.8976	0.9343
Chaharmahal-o-Bakhtiari	0.7123	0.6789	0.7123	0.7414
Markazi	0.8468	0.8108	0.8478	0.8814
Hamedan	0.6945	0.6615	0.6946	0.7229
Lorestan	0.7206	0.6870	0.7206	0.7500
Alborz	0.9802	0.9417	0.9868	1.0202
Tehran Bozorg	0.9802	0.9417	0.9802	1.0202
Ostan-e-Tehran	0.9607	0.9226	0.9607	0.9999
Ghom	0.8223	0.7869	0.8226	0.8559
Mashhad	0.9802	0.9417	0.9884	1.0202
Khorasan-e-Razavi	0.9417	0.9010	0.9417	0.9802
Khorasan-e-Jonubi	0.7004	0.6672	0.7004	0.7290
Khorasan-e-Shomali	0.8247	0.7872	0.8247	0.8584
Ahwaz	0.9802	0.9417	0.9868	1.0202
Khouzestan	0.9802	0.9417	0.9875	1.0202
Kohkiluyeh-o-Boyer-Ahmad	0.6560	0.6237	0.6571	0.6828
Zanjan	0.9802	0.9417	0.9814	1.0202
Ghazvin	0.9802	0.9417	0.9815	1.0202
Semnan	0.8517	0.8120	0.8517	0.8864
Sistan-o-Baluchestan	0.8156	0.7808	0.8156	0.8495
Kermanshah	0.7725	0.7359	0.7725	0.8041
Kordestan	0.9133	0.8700	0.9133	0.9506
Ilam	0.6115	0.5800	0.6116	0.6364
Shiraz	0.6625	0.6300	0.6630	0.6895
Fars	0.8431	0.7889	0.8431	0.8776
Boushehr	0.9487	0.9108	0.9488	0.9874
Shomal-e-Kerman	0.7790	0.7443	0.7790	0.8108
Jonub-e-Kerman	0.8123	0.7756	0.8123	0.8466
Gilan	0.8901	0.8470	0.8901	0.9265
Mazandran	0.7416	0.7077	0.7417	0.7719
Gharb-e-Mazandaran	0.6452	0.6131	0.6457	0.6715
Golestan	0.6518	0.6195	0.6518	0.6784
Hormozgan	0.9320	0.8944	0.9320	0.9700
Yazd	0.9802	0.9417	0.9802	1.0202

Table 9. Sensitivity analysis for RCDEA with $\gamma > 0.5$

DMU	RCDEA			
	Initial RCDEA	$e = 0.05$	$\Gamma = 5$	$\gamma = 0.6$
Tabriz	0.8352	0.8024	0.8352	0.9048
Azarbajejan Sharghi	0.6832	0.6533	0.6832	0.7401
Azarbajejan Gharbi	0.7825	0.7483	0.7825	0.8477
Ardabil	0.8352	0.8024	0.8352	0.9048
Esfahan	0.7058	0.6749	0.7055	0.7646
Ostan-e-Esfahan	0.7649	0.7116	0.7649	0.8286
Chaharmahal-o-Bakhtiari	0.6070	0.5785	0.6072	0.6575
Markazi	0.7185	0.6909	0.7224	0.7817
Hamedan	0.5918	0.5636	0.5919	0.6411
Lorestan	0.6140	0.5854	0.6143	0.6652
Alborz	0.8352	0.8024	0.8373	0.9048
Tehran Bozorg	0.8352	0.8024	0.8352	0.9048
Ostan-e-Tehran	0.6919	0.6647	0.6919	0.8868
Ghom	0.6950	0.6705	0.7009	0.7591
Mashhad	0.8352	0.8024	0.8352	0.9048
Khorasan-e-Razavi	0.8016	0.7677	0.8025	0.8693
Khorasan-e-Jonubi	0.5968	0.5685	0.5968	0.6465
Khorasan-e-Shomali	0.7028	0.6708	0.7028	0.7613
Ahwaz	0.8352	0.8024	0.8352	0.9048
Khouzestan	0.8352	0.8024	0.8352	0.9048
Kohkiluy-e-o-Boyerahmad	0.5590	0.5315	0.5596	0.6056
Zanjan	0.8352	0.8024	0.8362	0.9048
Ghazvin	0.8352	0.8005	0.8359	0.9048
Semnan	0.7257	0.6805	0.7158	0.7862
Sistan-o-Balouchestan	0.6931	0.6653	0.6950	0.7534
Kermanshah	0.6573	0.6271	0.6583	0.7131
Kordestan	0.7778	0.7406	0.7782	0.8431
Ilam	0.5210	0.4942	0.5160	0.5590
Shiraz	0.5610	0.5196	0.5647	0.6115
Fars	0.7185	0.6722	0.7061	0.7783
Boushehr	0.8084	0.7761	0.8084	0.8757
Shomal-e-Kerman	0.6628	0.6343	0.6638	0.7191
Jonub-e-Kerman	0.6922	0.6609	0.6931	0.7509
Gilan	0.7585	0.6741	0.7554	0.8210
Mazandran	0.6319	0.5442	0.6319	0.6846
Gharb-e-Mazandaran	0.5498	0.5215	0.5483	0.5956
Golestan	0.5554	0.5279	0.5558	0.6017
Hormozgan	0.7942	0.7189	0.7940	0.8602
Yazd	0.8352	0.8024	0.8352	0.9048

Table 10. Weights of criteria

Variables	BWM-RCDEA Weight $\gamma = 0.4$	BWM-RCDEA Weight $\gamma = 0.8$
Service area	0.0670	0.0753
Number of employees	0.1290	0.1351
Transformer capacity	0.1953	0.2050
Network length	0.1198	0.1195
Energy delivery	0.00001	0.00001
Number of customers	0.2001	0.1861
Total electricity sale	0.2965	0.2790