

# Effects of Low-Level Laser Irradiation on Mammalian Cell Cultures: Comparative Experimental Studies with Different Types of Lasers at 1260-1270 nm

A Khokhlova<sup>1</sup>, I Zolotovskii<sup>1</sup>, S Sokolovski<sup>2</sup>, Yu Saenko<sup>3</sup>, E Rafailov<sup>2</sup>, D Stoliarov<sup>1,2</sup>, E Pogodina<sup>3</sup>, A Gilmutdinova<sup>1</sup>, V Svetukhin<sup>4</sup> and A Fotiadi<sup>1,5,6</sup>

<sup>1</sup>Laboratory of Nonlinear and Microwave Photonics, Ulyanovsk State University, Ulyanovsk, Russia

<sup>2</sup>Aston Institute of Photonic Technologies, Aston University, Birmingham, UK

<sup>3</sup>Laboratory of Molecular and Cell Biology, Ulyanovsk state university, Ulyanovsk, Russia

<sup>4</sup>Scientific-Manufacturing Complex "Technological Centre", Moscow, Russia

<sup>5</sup>Université de Mons, Mons, Belgium

<sup>6</sup>Ioffe Physical-Technical Institute, St.-Petersburg, Russia

E-mail: avhohlova@gmail.com

**Abstract.** The effects observed under near-infrared laser irradiation of mammalian cells have been demonstrated to depend to a large extent on the type of irradiation source. In our experiments, we have measured concentration of reactive oxygen species in cell cultures of different origin (rodent and human, cancerous and non-cancerous) exposed to the radiation of low-level lasers at 1265 nm. Surprisingly, the radiation effects of narrowband laser occur to be more pronounced compared with those provided by the lasers of broader linewidth. Also, we have found that the aggressive types of cancer require a more accurate selection of irradiation parameters and laser operation regime.

## 1. Introduction

In recent years, the laser sources emitting in the near infrared range have been enthusiastically exploited in medical applications. An infrared light exhibits ability essential for medical needs to penetrate tissues deeper than the visible light [1-3].

The laser sources operating 1265-1270 nm range are of particular interest. Such sources enable singlet oxygen generation in inhomogeneous media and, in particular, in biological objects [4-8].

Within these wavelengths, the effects in biological objects have been observed at very low power. Therefore, the 1265-1270 nm radiation can be used in low-level laser therapy or in the so-called photobiomodulation therapy (PBMT) allowing minimal tissue damages [9-11].

Photobiomodulation applications range from regenerative medicine to cancer treatment [12-16]. It is very important to study the radiation-induced effects provided by different laser sources. The radiation emitted by different sources may possess different physical parameters thus having different effects on the biological objects.

In our study, the 1265 nm laser effects on mammalian cell cultures are explored. The performed in vitro studies have provided simple and convenient preclinical models that allow eliminating involvement of laboratory animals.



## 2. Methods

### 2.1. Cell cultures

The research has been performed on mammalian cell cultures.

1. CHO-K1 (ATCC® CCL-61™) is the Chinese hamster ovary epithelial cell line. The cells are not cancerous but immortalized.

2. HCT116 (ATCC® CCL-247™) is the human colon cancer cell line.

3. B16-F10 (ATCC® CRL-6475™) is the murine melanoma cell line. This transplantable cell culture is the most commonly exploited to study melanoma and to develop the laboratory animal models.

4. A875 (ATCC) is the human melanoma cell line.

### 2.2. Laser Sources

A semiconductor narrow-band (highly coherent) laser (Yenista Optics (Osics T100)) and a laser diode (Innolume, ThorLabs) operating at the wavelength of 1265 nm are used in the experiments. To adjust the diameter of the laser beam, the two collimators with waists of 3.5 and 7 mm are used.

### 2.3. Experiment

Twenty-four hours before the experiment, the cells were seeded in a slide chamber for adhesion and growth to the exponential phase. During the experiment, the cells were kept in a general or benchtop incubator in a humidified atmosphere, with 5% CO<sub>2</sub> and at a temperature of 37°C.

Since the laser irradiation at a higher power is accompanied by the heating effect, the culture medium temperature was measured in the well of the slide chamber before the experiment in order to keep the temperature inside the incubator at 37°C.

To avoid heating under the irradiation at higher powers, the control well was placed in the general incubator at 37°C to provide the same temperature. The slide-chamber was placed on the stand; the collimator was fixed at the bottom at a distance of 1 cm from the sample. During the irradiation of the experimental well, the control well was shielded with a metal foil.

The energy density of laser radiation is calculated as follows:

$$E = \frac{P \times t}{S}, \quad (1)$$

where  $P$  is the average output power ( $W$ ),  $t$  is the exposure time ( $sec$ ),  $S$  is the laser spot area on the cell culture ( $cm^2$ ) [4].


The irradiation time in all experiments was 30 minutes, unless otherwise indicated in Results and Discussion. Each experiment was performed in triplicate.

### 2.4. Fluorescent staining of cells

After irradiation, the cells were stained for fluorescence microscopy with dichloro-dihydro-fluorescein diacetate to evaluate intracellular concentration of the reactive oxygen species as described [17].


## 3. Results and Discussion

Figure 1 shows the concentration of reactive oxygen species in cell cultures immediately after irradiation with a low-power narrowband laser. The dotted line is the baseline (hereinafter) where the experimental and control values are nearly the same.



The ROS level in the Chinese hamster ovary and human colon cancer cells significantly increases after the laser irradiation. However, in melanoma cells, no increase of the oxidative stress is registered even after the exposure for 60 minutes (energy density is  $18.9\text{ J/cm}^2$ ). It could be explained by the different intracellular mechanisms.

Figure 2 shows the ROS concentration in Chinese hamster ovary cells and human colon cancer cells after the exposure to laser sources of different linewidths. One can see that the effects of narrowband laser irradiation are more pronounced. Noteworthy, the power of the laser diode was more than twice as high in these experiments than the power of narrowband laser.



It can be explained by the fact that the power spectral density of the narrowband low-power laser radiation at the point of its maximum is higher than that of the radiation provided by the high-power (broadband) laser diode: A ratio describing the power spectral densities for these laser radiation sources is as follows:


$$\frac{P_{\lambda l}}{P_{\lambda h}} \sim \frac{\Delta\lambda_h E_l}{\Delta\lambda_l E_h} \sim 2 \times 10^3 \frac{E_l}{E_h}, \quad (2)$$

where  $P_{\lambda l}$  is the power spectral density of the narrowband source ( $W$ ),  $P_{\lambda h}$  is the power spectral density of the broadband source ( $W$ ),  $E_l$  is the surface energy density of PBM ( $J/cm^2$ ),  $E_h$  is the surface energy density of high-power laser irradiation ( $J/cm^2$ ),  $\Delta\lambda_l$  is the linewidth of PBMT source ( $nm$ ),  $\Delta\lambda_h$  is the linewidth of high-power laser source ( $nm$ ).

Low-power and high-power laser linewidths are 0.001 nm and 1.5 nm, respectively. Therefore, in the present study, the power spectral density of the low-power laser radiation at the point of its maximum is higher than that of the radiation provided by the high-power laser. All this allows us to conclude existence of the narrowband resonance effect presumably on the surface of mitochondrial membranes [18-19].

Further experiments were performed with the 1265 nm laser diode to provide higher powers.

Figure 3 demonstrates resistance of the human melanoma cell line to laser irradiation. The power of 10 mW is required for normal cells or non-aggressive cancer to develop oxidative stress (Fig. 2), whereas melanoma demonstrates this effect at 250 mW.



**Figure 3.** Level of ROS in the A875 (human melanoma) cell line exposed to 1265 nm laser diode irradiation ( $t=30\text{min}$ );  $P=10\text{ mW}$ ,  $100\text{ mW}$ ,  $250\text{ mW}$ ;  $S=0.8\text{ cm}^2$ ;  $E_1=22.5\text{ J/cm}^2$ ,  $E_2=225\text{ J/cm}^2$ ,  $E_3=562.5\text{ J/cm}^2$ . \* - statistically significant difference between experiment and control.

Figure 4 shows the ROS concentration at different intensities of laser irradiation. A significant difference is registered in the oxidative stress level of human and murine melanoma cells. This can be explained by different origin of species (human and murine) or mutation and requires further research.





Figure 5 shows the concentration of reactive oxygen species 1 hour after irradiation. One can see that in the Chinese hamster ovary cells the oxidative stress is still increasing. It may indicate the antioxidant system failure leading to cell death. In melanoma cells, the level of oxidative stress decreases. Moreover, in the murine melanoma cells 1 hour after laser irradiation exposure there is no statistical significance between experiment and control groups.



#### 4. Conclusions

Our experiments have shown that the effects of the 1265 nm laser irradiation on biological cells depend not only on the irradiation dose but also on the laser linewidth, power, and intensity, as well as on the cell type.

#### Acknowledgments

The reported melanoma research was funded by Russian Foundation for Basic Research (RFBR) and Ulyanovsk Region, project number 19-415-730003 (to A. Khokhlova).

The reported non-melanoma research was funded by THE MINISTRY OF SCIENCE AND HIGHER EDUCATION OF THE RUSSIAN FEDERATION (MEGA-GRANT PROGRAM), grant number 075-15-2021-581 (to I. Zolovovskii, D. Stoliarov, A. Gilmutdinova, and A. Fotiadi).

#### Conflict of interests

The authors declare no conflict of interests.

#### References

- [1] Holick M F. Biological Effects of Sunlight, Ultraviolet Radiation, Visible Light, Infrared Radiation and Vitamin D for Health 2016 *Anticancer Research* **36** (3) 1345-1356
- [2] Martignago C C S, Tim C R, Assis L, Da Silva V R, Santos E C B D, Vieira F N, Parizotto N A, and Liebano R E. Effects of red and near-infrared LED light therapy on full-thickness skin graft in rats. 2020 *Lasers in Medical Science* **35** (1) 157-164

- [3] Barolet D, Christiaens F, and Hamblin M R. Infrared and skin: Friend or foe. *Journal of Photochemistry and Photobiology B: Biology* 2016 **155** 78-85
- [4] Saenko Y V, Glushchenko ES, Zolotovskii I O, Sholokhov E, and Kurkov A. Mitochondrial dependent oxidative stress in cell culture induced by laser radiation at 1265 nm 2016 *Lasers in Medical Science* **31** (3) 405-13
- [5] Sokolovski S G, Zolotovskaya S A, Goltsov A, Purreyron C, South A P, and Rafailov E U. Infrared laser pulse triggers increased singlet oxygen production in tumour cells 2013 *Scientific Reports* **12** **3** 3484
- [6] Anquez F, El Yazidi-Belkoura I, Randoux S, Suret P, and Courtade E. Cancerous cell death from sensitizer free photoactivation of singlet oxygen 2012 *Photochemistry and Photobiology* **88**(1):167-74
- [7] Yusupov A S, Goncharov S E, Zalevskii I D, Paramonov V M, and Kurkov A S. Raman fiber laser for the drug-free photodynamic therapy 2010 *Laser Physics* **20**: 357–359
- [8] Oliveira C S, Turchiello R, Kowaltowski A J, Indig G L, and Baptista M S. Major determinants of photoinduced cell death: Subcellular localization versus photosensitization efficiency 2011 *Free Radical Biology and Medicine* **51**(4) 824-33
- [9] Hennessy M and Hamblin M R. Photobiomodulation and the brain: a new paradigm 2017 *Journal of Optics* **19**(1) 013003
- [10] Hamblin M R, Nelson S T, and Strahan J R. Photobiomodulation and Cancer: What Is the Truth? 2018 *Photomedicine and Laser Surgery* **36**(5) 241-245
- [11] Glass G E. Photobiomodulation: A review of the molecular evidence for low level light therapy 2021 *Journal of Plastic, Reconstructive and Aesthetic Surgery* **74**(5) 1050-1060
- [12] Awotidebe A W, Inglis-Jassiem G, and Young T. Low-level laser therapy and exercise for patients with shoulder disorders in physiotherapy practice (a systematic review protocol) 2015 *Systematic Reviews* **4** 60
- [13] Ramos G A, Arliani G G, Astur D C, Pochini A C, Ejnisman B, and Cohen M. Rehabilitation of hamstring muscle injuries: a literature review 2016 *Revista Brasileira de Ortopedia* **52**(1) 11-16
- [14] Ash C, Town G, Whittall R, Tooze L, and Phillips J. Lasers and intense pulsed light (IPL) association with cancerous lesions 2017 *Lasers in Medical Science* **32**(8) 1927-1933
- [15] Błochowiak K, Andrysiak P, Sidorowicz K, Witmanowski H, Hędzielek W, and Sokalski J. Selected applications of Er:YAG and CO<sub>2</sub> lasers for treatment of benign neoplasms and tumorous lesions in the mouth 2015 *Postępy Dermatologii I Alergologii* **32**(5) 337-43
- [16] Schomaker M, Heinemann D, Kalies S, Willenbrock S, Wagner S, Nolte I, Ripken T, Murua Escobar H, Meyer H, and Heisterkamp A. Characterization of nanoparticle mediated laser transfection by femtosecond laser pulses for applications in molecular medicine 2015 *Journal of Nanobiotechnology* **13** 10
- [17] Khokhlova A, Zolotovskii I, Stoliarov D, Vorsina S, Liamina D, Pogodina E, Fotiadi A, Sokolovski S, Saenko Y, and Rafailov E. The Photobiomodulation of Vital Parameters of the Cancer Cell Culture by Low Dose of Near-IR Laser Irradiation 2019 *IEEE Journal of Selected Topics in Quantum Electronics* **1**(25) 1-10
- [18] Khokhlova A, Zolotovskii I, Sokolovski S, Saenko Y, Rafailov E, Stoliarov D, Pogodina E, Svetukhin V, Sibirny V, and Fotiadi A. The light-oxygen effect in biological cells enhanced by highly localized surface plasmon-polaritons 2019 *Scientific Reports* **9**(1) 18435
- [19] Khokhlova A, Zolotovskii I, Pogodina E, Saenko Yu, Stoliarov D, Vorsina S, Fotiadi A, Liamina D, Sokolovski S, and Rafailov E. Effects of high and low level 1265 nm laser irradiation on HCT116 cancer cells 2019 *Proceedings SPIE* **10861** Mechanisms of Photobiomodulation Therapy XIV 108610L